



**SCIENTIFIC COMMITTEE  
EIGHTEENTH REGULAR SESSION**

**ELECTRONIC MEETING**  
10-18 August 2022

---

**ANNUAL REPORT TO THE COMMISSION  
PART 1: INFORMATION ON FISHERIES, RESEARCH AND STATISTICS**

---

**WCPFC-SC18-AR/CCM-01**

**AUSTRALIA**



**Australian Government**  
**Department of Agriculture,**  
**Fisheries and Forestry**  
 ABARES

# Annual report to the Western and Central Pacific Fisheries Commission

## Part 1: Information on fisheries, research and statistics 2021 Australia

**S. Blake, B. D’Alberto, H. Patterson**

Research by the Australian Bureau of Agricultural and Resource Economics and Sciences

Technical Report 22.05

July 2022



<p>Scientific data were provided to the Commission in accordance with the decision relating to the provision of scientific data to the Commission on the 30th April 2022</p>	<p>No</p>
<p>If no, please indicate the reason(s) and intended actions: All data have been submitted except the observer/EM data. The observer/EM data submission was delayed due to database issues. These data will be submitted once full and complete data are available.</p>	

© Commonwealth of Australia 2022

### **Ownership of intellectual property rights**

Unless otherwise noted, copyright (and any other intellectual property rights, if any) in this publication is owned by the Commonwealth of Australia (referred to as the Commonwealth).

### **Creative Commons licence**

All material in this publication is licensed under a [Creative Commons Attribution 4.0 International Licence](#) except content supplied by third parties, logos and the Commonwealth Coat of Arms.

Inquiries about the licence and any use of this document should be emailed to [copyright@aff.gov.au](mailto:copyright@aff.gov.au).



### **Cataloguing data**

This publication (and any material sourced from it) should be attributed as: Blake, S, D'Alberto, B & Patterson, H 2022, *Annual report for the Western and Central Pacific Fisheries Commission Part 1: Information on fisheries, research and statistics 2021*, ABARES, Canberra, July. DOI: <https://doi.org/10.25814/02hj-4202>. CC BY 4.0.

ISSN 189-3128

Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES)

GPO Box 858 Canberra ACT 2601

Telephone 1800 900 090

Web [agriculture.gov.au](http://agriculture.gov.au)

### **Disclaimer**

The Australian Government acting through the Department of Agriculture, Fisheries and Forestry, represented by the Australian Bureau of Agricultural and Resource Economics and Sciences, has exercised due care and skill in preparing and compiling the information and data in this publication. Notwithstanding, the Department of Agriculture, Fisheries and Forestry, ABARES, its employees and advisers disclaim all liability, including liability for negligence and for any loss, damage, injury, expense or cost incurred by any person as a result of accessing, using or relying on any of the information or data in this publication to the maximum extent permitted by law.

### **Professional independence**

The views and analysis presented in ABARES publications reflect ABARES professionally independent findings, based on scientific and economic concepts, principles, information and data. These views, analysis and findings may not reflect or be consistent with the views or positions of the Australian Government or of organisations or groups that have commissioned ABARES reports or analysis. Learn more about ABARES [professional independence](#).

### **Acknowledgements**

The authors wish to thank Rupert Summerson (ABARES) for creating the maps. The authors also acknowledge and appreciate the input of Michael Dylewski (ABARES); Patty Hobsbawn (ABARES); Ashley Williams (CSIRO); Fiona Hill (AFMA); Kate Martin (AFMA); and Neil Hughes and Mat Kertesz (Department of Agriculture, Fisheries and Forestry).

### **Acknowledgement of Country**

We acknowledge the Traditional Custodians of Australia and their continuing connection to land and sea, waters, environment and community. We pay our respects to the Traditional Custodians of the lands we live and work on, their culture, and their Elders past and present.

# Contents

<b>Summary .....</b>	<b>1</b>
<b>1 Background .....</b>	<b>3</b>
1.1 Longline .....	3
1.2 Pole-and-line, purse seine and minor line.....	3
1.3 Recreational fishing .....	4
<b>2 Flag state reporting .....</b>	<b>5</b>
2.1 Domestic longlining catch and effort.....	5
2.2 Pole-and-line, purse seine and minor line catch and effort.....	11
2.3 Fishing patterns.....	14
2.4 Fleet operations.....	15
2.5 Species of special interest .....	16
2.6 Trends in size composition of retained catch .....	21
<b>3 Coastal state reporting .....</b>	<b>25</b>
<b>4 Socio-economic factors.....</b>	<b>26</b>
<b>5 Dispatch of catch.....</b>	<b>27</b>
<b>6 Onshore developments .....</b>	<b>28</b>
<b>7 Future prospects for the fishery .....</b>	<b>29</b>
<b>8 Status of data collection systems .....</b>	<b>30</b>
8.1 Logbook data collection and verification.....	30
8.2 Observer program .....	30
8.3 Port sampling program .....	30
8.4 Unloading/transhipment.....	31
8.5 Other.....	32
<b>9 Research activities.....</b>	<b>34</b>
9.1 Biological research projects .....	34
9.2 Assessment-related research projects.....	34
9.3 Ecological research projects.....	36
9.4 Bycatch research projects .....	36
<b>Appendix A: Common and Scientific Names.....</b>	<b>37</b>
<b>Appendix B: Mandatory mitigation measures in the ETBF 2022.....</b>	<b>39</b>
Seabirds .....	39
Sea turtles .....	40
Circle hooks.....	40
De-hooking device.....	41

Line cutting device .....	41
<b>References .....</b>	<b>42</b>

## Tables

Table 1. Annual catch and effort estimates (whole weight) for the Australian fleet, by gear and primary species, for the WCPFC Convention Area, 2017–21. ....	6
Table 2. Annual retained catch estimates (tonnes, t) of major non-target, associated and dependent species, including sharks, by the Australian fleet, by gear (longliners and other methods combined), in the WCPFC Convention Area, for 2017–21. ....	12
Table 3. Annual longline discard estimates (numbers) of major non-target, associated and dependent species, including sharks, by the Australian fleet in the WCPFC Convention Area, for 2017–21. ....	13
Table 4. Effort by Australian longliners and catch by primary species, within the AFZ and on the high seas, 2017–21. The percentage of catch or effort within the AFZ or on the high seas is provided in parentheses. ....	14
Table 5. Number of Australian vessels, by gear and size category, active in the WCPFC Convention Area, targeting WCPFC species, for 2017 to 2021. Gross registered tonnes (GRT) is the unit for vessel size. ....	16
Table 6. Interactions with species of special interest recorded in logbooks for the Australian longline fleet in the WCPFC Convention Area, 2017–21. Interactions not identified to species level are noted as unspecified (unspec). ....	17
Table 7. Observed captures of species of special interest for the Australian longline fleet in the WCPFC Convention Area, 2017–21. Interactions not identified to species level are noted as unspecified (unspec). ....	19
Table 8. Effort and observed seabird captures by fishing year for the ETBF a) south of 30°S, b) for 25°S – 30°S and c) for 23°N – 25°S. No seabird captures have been observed north of 23°N as there was no effort. For each year, the table gives number of longline vessels, total number of hooks (000's), number of observed hooks (000's), observer coverage (percentage of hooks that were observed), number of observed captures and the capture rate (captures per thousand hooks). Mitigation methods provided in Appendix B. ....	20
Table 9. Number of observed seabird captures in the ETBF, 2021, by species and area. ....	21
Table 10. Proportion of mitigation types used by the fleet 2017 to 2021. ....	21
Table 11. Summary of longline observer coverage (by hooks) for 2021. ....	30
Table 12. Estimated annual coverage of operational catch and effort (logbooks), observer data (% hooks) and port sampling (% coverage rate for the five main target species of individual fish weights collected from processors receiving longline-caught fish in the ETBF) for the Australian fleet active in the WCPFC Convention Area, 2017–21. ....	31
Table 13. Annual catch estimates (converted whole weights) for the ETBF for 2017–21 derived from catch disposal records. Estimates are in tonnes. ....	32
Table 14. Transshipment quantities by weight for Australian longline vessels in 2021 within the WCPFC Convention Area. ....	32
Table 15. Number of transshipment for Australian longline vessels in 2021. ....	32

## Figures

- Figure 1. Historical annual vessel numbers for the Australian fleet, by gear (longline, purse seine, pole-and-line and other commercial methods [minor line including trolling, rod-and-reel and handline]) for the WCPFC Convention Area. ....5
- Figure 2. Historical annual longline catch in the WCPFC Convention Area for (a) albacore, bigeye tuna and yellowfin tuna, as well as striped marlin and swordfish, and (b) purse seine catch skipjack tuna. ....7
- Figure 3 (a–e). Annual distributions of target species catch by the Australian longline fleet active in the WCPFC Convention Area, for 2017 to 2021. Catches have been aggregated to 5-degree blocks to address issues of confidentiality. The diameter of the pie chart in each grid cell is proportional to the total catch of that grid cell. The diameter to catch relationship is provided in the legend. The pie charts also show the proportions of catch of each of the target species. The segments of the pie chart in the legend are for illustrative purposes only. Fishing footprint shows the total extent of waters fished at a spatial resolution of 5-degree square. .... 10
- Figure 4. Longline effort distribution in the Eastern Tuna and Billfish Fishery (2021). Fishing footprint shows the total extent of waters fished at a spatial resolution of one degree square.... 15
- Figure 5 (a–e). Time series of 4-quarter and 8-quarter running means from 1997 to 2021, by quarter, of processed fish weight (in kg) of a) albacore, b) bigeye tuna, c) yellowfin tuna, d) striped marlin and e) swordfish sampled across the entire ETBF based on the data collected from the port sampling program. .... 24

# Summary

Australian commercial fisheries for highly migratory species in the Western and Central Pacific Fisheries Commission (WCPFC) Convention Area are managed as part of the Eastern Tuna and Billfish Fishery (ETBF) (a mainly longline fishery with a smaller minor line component) and Eastern Skipjack Fishery (a purse seine fishery). Most fishing occurs in the longline sector of the ETBF and, as such, this is the primary focus of the annual report. There was no activity in the Eastern Skipjack Fishery in 2021.

Total catches of WCPFC species of interest reported in logbooks for the ETBF decreased from 3,778 t (including <1 t minor line) in 2020 to 3,514 t (including <1 t minor line) in 2021. This is a decline from a peak of 8,229 t in 2002. Longline fishing effort in the ETBF has fallen from a peak of 12.40 million hooks in 2003 to 7.44 million total hooks in 2021. Of this, 7.42 million hooks, 99.8% of effort, was within the Australian exclusive economic zone (EEZ). A further 0.2% of effort occurred in high seas areas adjacent to the Australian EEZ. Thirty-five Australian vessels reported longlining in the WCPFC Convention Area during 2021.

The ETBF is a multi-species longline fishery, targeting albacore, bigeye tuna, yellowfin tuna and swordfish. The profitability and viability of the fishery is dependent on this suite of species, each of which fluctuate in availability in the Australian region through time. Longline logbook catch of albacore decreased from 1,105 t in 2020 to 1,020 t in 2021. Longline catch of bigeye tuna increased from 283 t in 2020 to 349 t in 2021. Longline catch of yellowfin tuna decreased from 1,670 t in 2020 to 1,377 t in 2021. Longline catch of swordfish increased from 536 t in 2020 to 589 t in 2021. Longline catch of striped marlin was 163 t in both 2020 and 2021. Longline catch of skipjack decreased from 2.5 t in 2020 to 2.2 t in 2021.

There were six dedicated minor line (including trolling, rod-and-reel and handline) vessels in the ETBF; most minor line catches are reported by vessels (e.g. longline vessels) on their way to and from fishing grounds. In 2021, seven vessels actively targeted tuna and billfish species using minor line in the ETBF, including one vessel which employed both longline and minorline. The number of vessels reporting using minor line has decreased in the ETBF from a peak of 52 vessels in 2001. There were no vessels active in the Eastern Skipjack Fishery (purse seine) in 2021.

The Australian Fisheries Management Authority (AFMA) observer program began deploying observers on domestic longliners in 2001, initially as part of a program to test the efficacy of seabird mitigation devices. From July 2003, observers were deployed more broadly across the fishery with the aim of collecting additional fishery data, including information on fishing gear and the size and species composition of catches. AFMA conducted a trial of electronic monitoring (i.e. e-monitoring; on-board, fixed-mount digital video cameras and other systems) in 2009–10 to evaluate the effectiveness of this technology for a range of fishery monitoring purposes and to conduct a cost–benefit analysis. E-monitoring of the fishery became compulsory from 1 July 2015 for vessels operating within the Australian Exclusive Economic Zone. As a minimum, 10% of the hauls are reviewed and used to acquit information provided in logbooks. The total number of longline hooks observed from the e-monitoring system in 2021 was 743,779, which is 9.9% of the hooks deployed.

The AFMA Commission agreed on the total allowable commercial catches (TACCs) for the ETBF. These apply to the 2021 season which commenced on 1 January 2021. The TACCs for the five main target species are: albacore (2,500 t); bigeye tuna (1,056 t); swordfish (1,163 t); striped marlin (351 t); and yellowfin tuna (2,400 t).

# 1 Background

Australian commercial fisheries for highly migratory species in the Western and Central Pacific Fisheries Commission (WCPFC) Convention Area are managed as part of the Eastern Tuna and Billfish Fishery (ETBF) (a mainly longline fishery with a small minor line component) and Eastern Skipjack Fishery (a purse seine fishery). Scientific and common names of relevant species are provided in Appendix A.

## 1.1 Longline

Japanese longliners began fishing off Australia's east coast in the late 1950s. Sporadic domestic longlining for yellowfin tuna commenced soon after, in the early 1960s. The declaration of the Australian Fishing Zone (AFZ) in 1979 resulted in Japanese longliners being licensed to fish in Australian waters under bilateral agreements. In the early 1980s, longlining increased markedly after successful air-freighting of fresh-chilled tuna to Japan. In the 1990s, effort expanded in the waters off northern Queensland, resulting in high catch rates of yellowfin and bigeye tuna.

In the mid-1990s, improved access to swordfish markets in the United States of America prompted many ETBF fishers to move to southern Queensland ports such as Mooloolaba to target swordfish. Japanese longliners were excluded from the AFZ from 1997. Longlining for swordfish has declined since early 2005 because of high fuel and bait costs, the introduction of a competitive total allowable catch (TAC) in 2006 (now an individual transferable quota system), and changes in the currency exchange rate.

The Securing our Fishing Future structural adjustment package (between 2005 and 2006) likely increased the exit of less-efficient vessels from the fishery. However, the number of longline vessels was already declining from a peak of 152 vessels in 1999; by 2007, 58 longline vessels remained, and in 2021 a total of 35 longline vessels were active.

Management through total allowable commercial catch (TACC) limits and individual transferable quotas (ITQs) commenced in 2011. Quota species are albacore, bigeye tuna, yellowfin tuna, swordfish and striped marlin. The level of latency in the ETBF, measured by the proportion of TACC not caught in the fishery, has varied across the key species since the shift to quota-based management in 2011. This is to be expected in a multi-species fishery, noting that latency is driven by a range of factors including catch-per-unit-effort, input costs, fish price and market access.

## 1.2 Pole-and-line, purse seine and minor line

The pole-and-line fishery expanded rapidly in the 1950s with the introduction of live-bait-and-pole techniques for southern bluefin tuna and sporadic catches of skipjack and yellowfin tuna. Pole-and-line fishing decreased in the late-1990s with little to no fishing by Australian fleets in the WCPFC Convention Area since then. The introduction of purse seining in the 1970s also increased catches. After peaking at 7,000 t in the early 1990s, purse-seine effort and catches of skipjack have decreased dramatically, with zero to very low effort and catches in recent years, although there is industry interest in re-invigorating the fishery. Minor line effort has been decreasing in the fishery over time, with a peak number of vessels in 2001 (52). There were six dedicated minor line vessels that operated in the ETBF in 2021.


### **1.3 Recreational fishing**

Recreational and charter anglers have taken tuna and billfish off eastern Australia since the early 1900s. During the 1970s, recreational vessels capable of operating offshore became more readily available and angling for tuna and billfish grew in popularity. The continental shelf extends less than 8 nautical miles (nm) offshore in some places along the southeast coast of Australia, allowing anglers to fish for tuna from shore at several locations. The Game Fishing Association of Australia (GFAA) was formed in 1938 and has a membership of several thousand anglers, most based on the east coast of Australia. Many gamefishers tag and release much of their catch, especially marlins.

## 2 Flag state reporting

### 2.1 Domestic longlining catch and effort

Unless otherwise stated, all catch and effort levels in this report are derived from those reported in Australian Fisheries Management Authority (AFMA) logbooks. Thirty-five vessels in the Australian fleet reported longlining in the WCPFC Convention Area during 2021, down from a peak of 180 in 1997 (Figure 1). Total longline effort decreased from 8.2 million hooks in 2020 to 7.44 million hooks in 2021 (Table 1). Overall, effort has declined from the peak effort of 12.40 million hooks deployed in 2003. This decline is mainly the result of the strength of the Australian dollar, increased operating costs, the surrender of permits under the structural adjustment component of the Australian Government Securing Our Fishing Future package, the introduction of hook limits in 2009 and the introduction of individual transferrable quota management in 2011. The number of vessels in 2021 fishing for striped marlin south of 15°S was 31. Thirty-five vessels fished for albacore and 31 fished for swordfish south of 20°S.



**Figure 1. Historical annual vessel numbers for the Australian fleet, by gear (longline, purse seine, pole-and-line and other commercial methods [minor line including trolling, rod-and-reel and handline]) for the WCPFC Convention Area.**


**Table 1. Annual catch and effort estimates (whole weight) for the Australian fleet, by gear and primary species, for the WCPFC Convention Area, 2017–21.**

Fishing method	Year	Effort <sup>a</sup>	Primary species (t)					
			Albacore	Bigeye	Skipjack	Yellowfin	Striped marlin	Swordfish
<b>All gears</b>	2017	-	785.1	374.8	7.4	1490.7	244.6	975.5
	2018	-	694.6	313.8	3.1	1278.2	208.9	819.9
	2019	-	759.9	246.6	4.1	1896.4	211.5	694.8
	2020	-	1106.9	283.0	2.5	1670.3	162.5	536.3
	2021	-	1020.9	349.0	2.2	1377.5	162.8	589
<b>Longline <sup>b</sup></b>	2017	8737	785.0	374.8	7.1	1490.7	244.6	975.5
	2018	7896	694.6	313.8	3.1	1278.2	208.9	819.7
	2019	8567	759.9	246.6	4.0	1896.4	211.5	694.8
	2020	8218	1105.4	283.0	2.5	1670.5	162.7	536.3
	2021	7444	1020.4	349.0	2.2	1377.4	162.8	589.0
<b>Purse seine</b>	2017	0	0	0	0	0	0	0
	2018	0	0	0	0	0	0	0
	2019	0	0	0	0	0	0	0
	2020	0	0	0	0	0	0	0
	2021	0	0	0	0	0	0	0


Note: **a** Longline—number of hooks (000's); purse seine—search hours. **b** Includes small catches from other commercial methods (minor line component including trolling, rod-and-reel, handline and pole-and-line). Note: Australia is currently reviewing catch estimation methods and therefore values reported here may be amended in future reports.

Total longline catches of WCPFC species of interest in the ETBF reported in logbooks decreased from 3,778 t in 2020 to 3514 t in 2021. This is down from a peak of 8,229 t in 2002. Historical catches for the Australian fleet in the WCPFC Convention Area, by primary species, are shown in Figure 2. Longline catches of albacore decreased from 1,105 t in 2020 to 1,020 t in 2021 (846 t caught south of 20°S). Longline catches of bigeye tuna increased from 283 t in 2020 to 349 t in 2021. Longline catches of yellowfin tuna decreased from 1,670 t in 2020 to 1,377 t in 2021. Longline catches of swordfish increased from 536 t in 2020 to 589 t in 2021 (563 t caught south of 20°S). Longline catch of striped marlin was 163 t in both 2020 and 2021 (162 t caught south of 15°S in 2021). Longline catches of skipjack decreased from 2.5 t in 2020 to 2.2 t in 2021. Annual catch distributions of the main target species by the Australian longline fleet for 2016 to 2021 are shown in Figure 3.

a)




b)




**Figure 2. Historical annual longline catch in the WCPFC Convention Area for (a) albacore, bigeye tuna and yellowfin tuna, as well as striped marlin and swordfish, and (b) purse seine catch skipjack tuna.**


a)




b)




c)



d)



e)



**Figure 3 (a–e). Annual distributions of target species catch by the Australian longline fleet active in the WCPFC Convention Area, for 2017 to 2021. Catches have been aggregated to 5-degree blocks to address issues of confidentiality. The diameter of the pie chart in each grid cell is proportional to the total catch of that grid cell. The diameter to catch relationship is provided in the legend. The pie charts also show the proportions of catch of each of the target species. The segments of the pie chart in the legend are for illustrative purposes only. Fishing footprint shows the total extent of waters fished at a spatial resolution of 5-degree square.**

Annual retained catch estimates of major non-target, associated and dependent species, including sharks, by the Australian longline fleet from 2017–21 are presented in Table 2. Estimates of releases derived from logbooks are in Table 3. From July of 2015, the logbooks of the Australian longline fleet were subject to potential verification through e-monitoring coverage. This has resulted in a rise in the levels of discards reported in logbooks since 2016 (Table 3).

Since 1 January 2013, retention of oceanic whitetip sharks by all commercial vessels has been prohibited and no retention was recorded in logbooks in 2021 (Table 2) while 1,629 were reported as discarded in logbooks (Table 3). Of the 22 oceanic whitetips observed caught, 11 were released alive and 11 were released in an undetermined condition.

Since 1 July 2014, retention of silky sharks has been prohibited and no retention was recorded in logbooks in 2021 (Table 2) while 60 were reported as discarded (Table 3). In the 2021 calendar year, no silky sharks were observed caught in the ETBF.

Since 14 September 2014 hammerhead sharks must be reported by species under CITES listings. For the purposes of this report, however, we have continued to group hammerheads into a single group for consistency in data presentation.

Further restrictions on shark catches in the ETBF can be found at [https://afma.govcms.gov.au/sites/default/files/2020\\_etbf\\_management\\_arrangements\\_booklet\\_-\\_final.pdf](https://afma.govcms.gov.au/sites/default/files/2020_etbf_management_arrangements_booklet_-_final.pdf)

Retention of both blue marlin and black marlin by commercial longliners has been prohibited since 1998 and no retention was recorded in logbooks in 2021, while 1,486 black marlin and 1,220 blue marlin were reported as discarded (Table 3). The vast majority of the catch and effort by Australian longliners has been within the AFZ, with little effort on the adjacent high seas (Table 4).

## **2.2 Pole-and-line, purse seine and minor line catch and effort**

In 2021, there were no active purse-seine vessels in the Eastern Skipjack Fishery and no interactions with whale sharks were recorded (CMM 2012-06). Total minor line catches of WCPFC species of interest in the ETBF (including pole and line, trolling, rod-and-reel and handline) was <1 t in 2021. The only catch of target species was albacore (0.47 t) and yellowfin tuna (0.16 t). The number of vessels reporting using minor line in the ETBF has steadily decreased from a peak of 52 vessels in 2001 to seven in 2021 (using troll, rod-and-reel and handline). Minor line effort peaked in 2007 with 975 lines.

**Table 2. Annual retained catch estimates (tonnes, t) of major non-target, associated and dependent species, including sharks, by the Australian fleet, by gear (longliners and other methods combined), in the WCPFC Convention Area, for 2017–21.**

Group	Species	Longline (t)					Other methods (t)				
		2017	2018	2019	2020	2021	2017	2018	2019	2020	2021
Scalefish	Escolar	14.6	23.8	26.5	7.7	0.2	0.0	0.0	0.0	0.0	0.0
	Lancetfish	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0
	Mahi mahi	155	128.1	119.3	81.3	52.9	0.0	0.0	0.0	0.0	0.0
	Moonfish	7.1	14	12.3	25.5	33.5	0.0	0.0	0.0	0.0	0.0
	Ocean sunfish	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	Oilfish	0.0	0.1	0.6	2.2	0.5	0.0	0.0	0.0	0.0	0.0
	Ray's bream	19.5	22.4	19.3	15.5	7.9	0.0	0.0	0.0	0.1	0.0
	Rudderfish	43.4	43.9	19.4	8.4	10.6	0.0	0.0	0.0	0.0	0.0
	Sailfish	2.2	1.6	2.4	1.3	1.7	0.0	0.0	0.0	0.0	0.0
	Shortbill spearfish	12.5	11.8	16.9	10.4	4.4	0.0	0.0	0.0	0.0	0.0
	Wahoo	18.4	14.9	14.7	13.3	12.6	0.0	0.0	0.0	0.0	0.0
<b>Subtotal</b>	<b>272.7</b>	<b>260.6</b>	<b>231.4</b>	<b>165.9</b>	<b>124.3</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.1</b>	<b>0.0</b>	
Sharks	Blacktip shark	0.1	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0
	Blue shark	0.2	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Bronze whaler	0.3	0.4	0.5	0.2	0.0	0.0	0.0	0.0	0.0	0.0
	Dusky shark	2.3	0.3	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	Hammerhead	0.2	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	Longfin mako	0.0	0.2	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	Oceanic whitetip	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Porbeagle	0.8	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	Shortfin mako	31.4	24.9	23.7	16	12.1	0.0	0.0	0.0	0.0	0.2
	Silky shark	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	Thresher shark	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	Tiger shark	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	Whale shark	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>Subtotal</b>	<b>35.4</b>	<b>26</b>	<b>24.8</b>	<b>16.8</b>	<b>12.1</b>	<b>0.0</b>	<b>0.2</b>	<b>0.0</b>	<b>0.0</b>	<b>0.2</b>	
<b>TOTAL</b>	<b>308.1</b>	<b>286.6</b>	<b>256.2</b>	<b>182.7</b>	<b>136.4</b>	<b>0.0</b>	<b>0.2</b>	<b>0.0</b>	<b>0.1</b>	<b>0.2</b>	

**Table 3. Annual longline discard estimates (numbers) of major non-target, associated and dependent species, including sharks, by the Australian fleet in the WCPFC Convention Area, for 2017–21.**


<b>Group</b>	<b>Species</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>	<b>2021</b>
<b>Scalefish</b>	Black marlin	1171	1333	1506	1310	1486
	Blue marlin	1201	1038	984	681	1220
	Escolar	1342	1192	1636	2041	564
	Lancetfish	24,452	17,295	21,421	19,346	21,303
	Mahi mahi	933	864	703	404	568
	Moonfish	9	7	5	82	27
	Ocean sunfish	2402	2834	1379	2149	5770
	Oilfish	47	41	54	258	745
	Ray's bream	340	201	115	181	66
	Rudderfish	2690	1767	3845	4349	3902
	Sailfish	79	24	74	38	73
	Shortbill spearfish	168	115	171	131	100
	Wahoo	73	55	68	47	90
	<b>Subtotal</b>	<b>34,907</b>	<b>26,766</b>	<b>31,961</b>	<b>31,017</b>	<b>35,914</b>
<b>Sharks</b>	Blacktip sharks	1	0	6	0	0
	Blue shark	16,860	13,820	13,394	14,871	20,677
	Bronze whaler	4143	3477	4568	2331	11,378
	Dusky shark	1847	1648	2339	1470	333
	Hammerhead	487	476	433	287	374
	Longfin mako	18	6	1	3	0
	Oceanic whitetip	1359	806	1071	1086	1629
	Porbeagle	124	4	1	1	0
	Shortfin mako	1449	1311	1219	805	1061
	Silky shark	395	130	54	264	60
	Thresher shark	651	664	634	970	856
	Tiger shark	726	749	682	1621	1673
	Whale shark	0	0	0	0	0
	<b>Subtotal</b>	<b>28,060</b>	<b>23,091</b>	<b>24,402</b>	<b>23,709</b>	<b>38,041</b>
	<b>TOTAL</b>	<b>62,967</b>	<b>49,857</b>	<b>56,363</b>	<b>54,726</b>	<b>73,955</b>

**Table 4. Effort by Australian longliners and catch by primary species, within the AFZ and on the high seas, 2017–21. The percentage of catch or effort within the AFZ or on the high seas is provided in parentheses.**

	Year	Effort ('000 hooks)	Primary species catch (t)					
			Albacore	Bigeye	Skipjack	Yellowfin	Striped marlin	Swordfish
<b>Within AFZ</b>	2017	8663	784.3	373.5	7.1	1485.3	240.9	960.7
		(99.2%)	(99.9%)	(99.6%)	(100.0%)	(99.6%)	(1.5%)	(98.5%)
	2018	7783	692.6	311.2	3.1	1273.9	202.1	804.6
		(98.6%)	(99.7%)	(99.2%)	(100.0%)	(99.7%)	(96.7%)	(98.2%)
	2019	8539	757.6	245.9	4.0	1893.9	210.6	690.7
	(99.7%)	(99.7%)	(99.7%)	(100.0%)	(99.9%)	(99.6%)	(99.4%)	
	2020	8134	1100.6	281.1	2.5	1660.9	161.2	525.0
		(99.1%)	(99.5%)	(99.3%)	(99.6%)	(99.4%)	(99.2%)	(97.9%)
	2021	7424	1019.4	348	2.2	1376.8	162.8	588.6
		(99.8%)	(99.9%)	(100%)	(100%)	(99.9%)	(100%)	(99.9%)
<b>High seas</b>	2017	73	0.6	1.3	0	5.4	3.8	14.8
		(0.8%)	(0.1%)	(0.4%)	(0.0%)	(0.4%)	(1.5%)	(1.5%)
	2018	113	2.0	2.6	0	4.3	6.8	15.1
		(1.4%)	(0.3%)	(0.8%)	(0.0%)	(0.3%)	(3.3%)	(1.8%)
	2019	27	2.3	0.7	0	2.5	0.9	4.2
	(0.3%)	(0.3%)	(0.3%)	(0.0%)	(0.1%)	(0.4%)	(0.6%)	
	2020	72	5.9	2	0.01	9.4	1.3	11.2
		(0.9%)	(0.5%)	(0.7%)	(0.4%)	(0.6%)	(0.8%)	(2.1%)
	2021	17	1	0	0	0.6	0	0.1
		(0.2%)	(0.1%)	(0%)	(0%)	(<0.1%)	(0%)	(<0.1%)

## 2.3 Fishing patterns

Fishing patterns vary with target species, location and season. The management area of the ETBF extends from Cape York, at the northern tip of Queensland, to the border between Victoria and South Australia, including waters around Tasmania (Figure 4). In the WCPFC Convention Area, skipjack tuna are fished from southern New South Wales to north-eastern Tasmania.



**Figure 4. Longline effort distribution in the Eastern Tuna and Billfish Fishery (2021). Fishing footprint shows the total extent of waters fished at a spatial resolution of one degree square.**

## 2.4 Fleet operations

Domestic longlining vessels are mostly 15–25 m long and use monofilament gear (Table 5). Vessels usually conduct one longline operation per day, or night, depending on the target species. A typical longline set will comprise about 1400+ hooks. Fishers commonly operate around 150 days per year. Most trips are between 2 and 15 days and typically range from 40–300 nm from port with effort mostly concentrated on the edge of the continental shelf (seaward of 200 m depth) but extending much further eastward off Mooloolaba (Figure 4). The catch is gilled and gutted (depending on species) and stored on ice, in ice slurry or in refrigerated brine.

Historically, most purse-seiners were 20–25 m long, but several were 40–45 m. Most poling vessels were 15–20 m long. Most minor line catches are reported by vessels (e.g. longline vessels) on their way to and from fishing grounds.

**Table 5. Number of Australian vessels, by gear and size category, active in the WCPFC Convention Area, targeting WCPFC species, for 2017 to 2021. Gross registered tonnes (GRT) is the unit for vessel size.**

Year	Longline				Subtotal	Purse seine	Pole-and-line	Troll	Total
	Vessel size (GRT)	≤50	51–200	201–500		≤500	0–50	Unknown	
2017		16	23	0	39	0	0	2	41
2018		16	23	1	40	0	0	0	40
2019		10	26	1	37	0	0	0	37
2020		8	27	1	36	0	0	2	38
2021		9	25	1	35	0	0	2	41 <sup>a</sup>

**a** The total of forty-one vessels that fished in the WCPFC Convention Area in 2021 include three rod and reel vessels and one handline vessel.

## 2.5 Species of special interest

Australia implements a mandatory reporting scheme for fisheries interactions with protected species, which includes species of special interest. Interactions with these species are recorded by fishers in their logbooks and are reported to AFMA. These interactions are then forwarded to the Department of Agriculture, Water and the Environment on a quarterly basis. A summary of these interactions, from logbooks, is presented in Table 6. Life status of the animal involved in the interaction is also recorded. In 2021, interactions were recorded with 16 green turtles (14 alive and 2 dead), 20 leatherback turtles (19 alive and 1 injured), 12 loggerhead turtles (11 alive and 1 dead), 2 hawksbill turtles (both alive), 5 Olive Ridley turtles (4 alive and 1 dead), 1 flatback turtle (dead) and 33 unidentified turtles (28 alive and 5 dead). Interactions were also recorded with 16 flesh footed shearwaters (1 alive and 15 dead), 1 short tailed shearwater (dead), 17 unspecified albatrosses (9 alive and 8 unknown), 11 unspecified birds (10 dead, 1 unknown), 4 short-finned pilot whales (3 alive and 1 dead), 2 long-finned pilot whales (both alive), 1 bottlenose dolphin (alive), 1 unidentified dolphin (alive), 9 unidentified seals (all alive) and 5 unknown or other species of interest (all dead). From July of 2015, the logbooks of the Australian longline fleet were subject to potential verification through electronic monitoring. This has resulted in an increase in the reporting level of interactions with species of special interest since 2016 (Table 6).

Observed captures are reported in Table 7. In 2021, there were 17 observed captures of sea turtles (10 alive, 1 dead and 6 of unknown life status); and 10 captures of seabirds (9 alive and 1 dead).

### 2.5.1 Sea turtles

The Eastern Tuna and Billfish Fishery requires the use of large circle hooks in Australia's shallow-set pelagic longline fisheries targeting swordfish, consistent with CMM2008-03. Mitigation requirements in the ETBF for sea turtles are reported in Appendix B.

## 2.5.2 Seabirds and marine mammals

Australia has extensive mitigation requirements for seabirds in the ETBF which are shown in Appendix B. More specific seabird interaction information, including the observed capture rate and captures by species and area for 2021, is presented in Table 8 and Table 9. Table 10 provides the proportion of mitigation measures used.

**Table 6. Interactions with species of special interest recorded in logbooks for the Australian longline fleet in the WCPFC Convention Area, 2017–21. Interactions not identified to species level are noted as unspecified (unspec).**

Group	Common name	2017	2018	2019	2020	2021
Seabirds	Black-browed albatross	3	4	2	3	0
	Shy albatross	0	0	1	1	0
	Sooty albatross	0	0	0	1	0
	Wandering albatross	1	6	3	0	0
	Albatrosses (unspec)	33	56	50	9	17
	Cape Petrel	0	0	1	0	0
	Sooty shearwater	1	1	0	0	0
	Short-tailed shearwater	0	1	2	0	1
	Flesh-footed shearwater	2	0	0	0	16
	Shearwaters (unspec)	6	17	37	8	0
	Terns	0	0	0	1	0
	Australian gannet	0	4	0	3	0
	Cormorants	1	0	0	0	0
	Birds (unspec)	0	3	5	3	11
	<b>Subtotal</b>		<b>47</b>	<b>92</b>	<b>101</b>	<b>29</b>
Turtles	Green turtle	96	36	52	31	16
	Hawksbill turtle	2	5	8	3	2
	Leatherback turtle	53	68	53	21	20
	Loggerhead turtle	26	18	12	13	12
	Flatback turtle	0	0	1	0	1
	Pacific (Olive) Ridley turtle	5	7	9	7	5
	Turtles (unspec)	18	22	18	20	33
<b>Subtotal</b>		<b>200</b>	<b>156</b>	<b>153</b>	<b>95</b>	<b>89</b>
Mammals	Melon-headed whale	4	0	0	2	0
	Baleen whales	0	0	0	0	0
	Toothed whales	0	0	0	0	0
	Short-finned pilot whale	6	7	4	1	4
	Long-finned pilot whale	4	1	0	0	2
	False killer whale	1	0	4	3	0
	Humpback whale	1	1	0	0	0
	Whales (unspec)	5	0	1	3	0
	Common dolphin	0	0	3	3	0
	Bottlenose dolphin	1	0	1	1	1
	Dolphin (unspec)	9	4	7	4	1
	Australian fur seal	0	0	0	0	0
	New Zealand fur seal	0	0	2	0	0
	Seals (unspec)	2	5	0	1	9
	Dugong	1	0	0	0	0
<b>Subtotal</b>		<b>34</b>	<b>18</b>	<b>22</b>	<b>18</b>	<b>17</b>

Unknown or other	0	0	0	1	5
<b>TOTAL</b>	<b>281</b>	<b>266</b>	<b>276</b>	<b>143</b>	<b>156</b>

**Table 7. Observed captures of species of special interest for the Australian longline fleet in the WCPFC Convention Area, 2017–21. Interactions not identified to species level are noted as unspecified (unspec).**

<b>Group</b>	<b>Common name</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>	<b>2021</b>
<b>Seabirds</b>	Black-browed albatross	0	0	0	0	0
	Shy albatross	0	0	0	0	0
	Wandering albatross	0	0	1	0	0
	Albatrosses (other)	2	9	3	4	5
	Cape petrel	0	0	0	0	0
	Gannets and boobies	0	0	0	3	0
	Great crested tern	0	0	0	0	0
	Flesh-footed shearwater	0	0	0	0	0
	Wilson’s storm petrel	0	0	0	0	0
	Petrels Prions and Shearwaters	0	4	6	0	1
	Shearwaters	0	0	1	2	3
	Birds (unspec)	0	1	0	2	1
	<b>Subtotal</b>		<b>2</b>	<b>14</b>	<b>11</b>	<b>11</b>
<b>Turtles</b>	Green turtle	2	0	5	0	0
	Hawksbill turtle	0	0	0	0	0
	Leatherback turtle	8	11	2	0	6
	Loggerhead turtle	0	0	0	0	0
	Pacific (Olive) Ridley	0	0	0	0	0
	Turtles (unspec)	5	14	16	9	11
<b>Subtotal</b>		<b>15</b>	<b>25</b>	<b>23</b>	<b>9</b>	<b>17</b>
<b>Mammals</b>	Dolphin (unspec)	1	1	2	0	0
	Long-finned pilot whale	0	0	0	0	0
	Short-finned pilot whale	0	0	1	0	0
	Whales (unspec)	4	2	1	1	0
	Australian fur seal	0	0	0	0	0
<b>Subtotal</b>		<b>5</b>	<b>3</b>	<b>4</b>	<b>1</b>	<b>0</b>
<b>TOTAL</b>		<b>22</b>	<b>42</b>	<b>38</b>	<b>21</b>	<b>27</b>

CMM 2011-03 regarding the intentional setting of purse-seine gear on cetaceans entered into force on 1 January 2013. Such setting practices are prohibited in Australian purse-seine fisheries since the introduction of the *Environment Protection and Biodiversity Conservation Act*. There were no reported interactions with cetaceans in purse-seine fisheries in 2021.

**Table 8. Effort and observed seabird captures by fishing year for the ETBF a) south of 30°S, b) for 25°S – 30°S and c) for 23°N – 25°S. No seabird captures have been observed north of 23°N as there was no effort. For each year, the table gives number of longline vessels, total number of hooks (000's), number of observed hooks (000's), observer coverage (percentage of hooks that were observed), number of observed captures and the capture rate (captures per thousand hooks). Mitigation methods provided in Appendix B.**

a) South of 30°S

Year	Fishing effort (000's hooks)				Observed seabird captures	
	Number of vessels	Number of hooks	Observed hooks	% hooks observed	Capture number	Capture rate
2016	31	2471	278	11.2	2	0.007
2017	32	2183	206	9.4	2	0.010
2018	37	3084	345	11.2	8	0.023
2019	33	2537	306	12.1	8	0.026
2020	30	1721	168	9.8	9	0.005
2021	29	1801	170	9.4	7	0.004

b) 25°S – 30°S

Year	Fishing effort (000's hooks)				Observed seabird captures	
	Number of vessels	Number of hooks	Observed hooks	% hooks observed	Capture number	Capture rate
2016	26	3718	326	8.8	1	0.003
2017	31	3816	391	10.3	0	0.000
2018	27	2917	298	10.2	5	0.017
2019	26	3264	391	12.0	3	0.008
2020	22	3990	409	10.2	2	0.005
2021	20	2600	266	10.2	1	0.000

c) 23°N – 25°S

Year	Fishing effort (000's hooks)				Observed seabird captures	
	Number of vessels	Number of hooks	Observed hooks	% hooks observed	Capture number	Capture rate
2016	20	1633	131	8.0	0	0.000
2017	22	2744	300	10.9	0	0.000
2018	22	1897	212	11.2	1	0.005
2019	18	2769	302	10.9	0	0.000
2020	18	2409	219	9.8	0	0.000
2021	17	2990	286	9.6	2	0.000

**Table 9. Number of observed seabird captures in the ETBF, 2021, by species and area.**

Species	South of 30°S	25°S–30°S	23°N–25°S	North of 23°N	Total Captures
Birds	0	0	1	0	1
Albatrosses	5	0	0	0	5
Gannets and boobies	0	0	0	0	0
Shearwaters	2	1	1	0	4
<b>Total</b>	<b>7</b>	<b>1</b>	<b>2</b>	<b>0</b>	<b>10</b>

**Table 10. Proportion of mitigation types used by the fleet 2017 to 2021.**


Combination of mitigation measures	Proportion of observed effort using mitigation measures				
	2017	2018	2019	2020	2021
TL + WB	0.90	0.85	0.84	0.89	0.87
TL + WB + NS	0.10	0.15	0.16	0.11	0.13
<b>Total</b>	<b>1.00</b>	<b>1.00</b>	<b>1.00</b>	<b>1.00</b>	<b>1.00</b>

Note: Proportions were derived from the AFMA observer database and, from July 2015 onwards, from electronic monitoring data. TL = tori line, NS = Night setting, WB = weighted branch lines.


## 2.6 Trends in size composition of retained catch

The size composition (based on processed weights) of albacore, bigeye tuna, yellowfin tuna and striped marlin is shown in Figure 5. Albacore has maintained a high 8-quarter running mean weight of around 18 kg since 2016 (Figure 5a). The 8-quarter running mean of bigeye tuna declined to around 35 kg by the end of 2019 and has since remained fairly stable (Figure 5b). Yellowfin tuna maintained a high 8-quarter running mean weight of 39-40kg between 2016 to 2018 before decreasing to ~ 37kg in 2019 to 2021 (Figure 5c). The 8-quarter running mean weight of striped marlin hovered around 63 kg from 2018 to 2020 before decreasing to 60 kg in 2021 (Figure 5d). Swordfish showed an increasing trend in its 8-quarter running mean between 2015 to 2018 before declining to around 42 kg by the end of 2021 (Figure 5e).


a)




b)




c)



d)



e)



**Figure 5 (a–e).** Time series of 4-quarter and 8-quarter running means from 1997 to 2021, by quarter, of processed fish weight (in kg) of a) albacore, b) bigeye tuna, c) yellowfin tuna, d) striped marlin and e) swordfish sampled across the entire ETBF based on the data collected from the port sampling program.

# 3 Coastal state reporting

There are currently no foreign fishing vessels licensed to operate in the AFZ.

## 4 Socio-economic factors

In 2015–16, the gross value of production (GVP) of the ETBF reached an 11-year peak in real terms (2020–21 dollars) of \$52.9 million. Between 2016-17 and 2020-21 GVP has remained below the 2015–16 peak because of falling catch volumes, despite generally improved prices for key species during that time period. In 2020–21 GVP increased from the previous financial year by 10% to \$35.6 million largely because of lower yellowfin tuna and broadbill swordfish catch volume and prices.

## 5 Dispatch of catch

The combined value of Australian exports of albacore, bigeye tuna and yellowfin tuna (the three key species of tuna caught in the WCPFC) decreased from the previous financial year by 20% to \$14.2 million in 2020–21. The export value of albacore and bigeye tuna increased, driven by higher export volumes and (to a lesser extent) higher export prices. This increase was offset by a significant (37%) decline in yellowfin tuna export value, reflecting lower export volumes. Swordfish is typically the second most valuable species group landed in the ETBF after yellowfin tuna. In 2020–21 the value of swordfish exports decreased by 57% to \$1.8 million, largely reflecting lower export volumes. The trade disruptions caused by COVID-19 in 2020-21 resulted in some producers seeking to balance risk, including using new or alternative selling platforms and diversifying away from export markets to domestic ones.

## 6 Onshore developments

Nil

## 7 Future prospects for the fishery

Commercial operators view the Australian skipjack fisheries as an important development opportunity because significant catching capacity exists in Port Lincoln, South Australia. Currently, catches are low as a result of variability in the availability of skipjack tuna in the AFZ, variable participation levels, low profit margins and the closure of the Port Lincoln cannery; however, there is room for development in this fishery.

The AFMA Commission agreed on the total allowable commercial catches (TACCs) for the ETBF. These apply to the 2022 season which commenced on 1 January 2022. The TACCs for the five main target species are: albacore (2,500 t); bigeye tuna (1,056 t); swordfish (1,047 t); striped marlin (351 t); and yellowfin tuna (2,400 t).

## 8 Status of data collection systems

### 8.1 Logbook data collection and verification

AFMA introduced a logbook for domestic longliners in 1986. The logbook has been revised on several occasions. The latest (AL06—Australian Pelagic Longline Daily Fishing Log) was introduced in 2007; vessels began submitting AL06 logbooks in November 2007. Return of logbooks by Australian longliners improved when, in 1995, it became a condition of fishing permits and has been close to 100% in recent years. Logbooks have also been introduced for the skipjack tuna purse-seine fisheries; PS01—Australian Purse Seine Daily Fishing Log was distributed in July 2002 with the first skipjack tuna catch recorded in this logbook in December 2003. Weights from catch disposal records are verified; weights recorded on logbooks are an estimate only. From 1 July 2015 logbooks have been verified through e-monitoring. As of 2021, all reporting in the ETBF is done via electronic logbooks.

### 8.2 Observer program

AFMA observers were deployed on domestic longliners from 2001 to 2015. From July 2003 to 2015, observers were deployed more broadly across the fishery with more general duties, such as the collection of data on fishing gear and the size and species composition of catches.

AFMA implemented a trial of e-monitoring to evaluate the effectiveness of this technology for a range of fishery monitoring purposes and to conduct a cost-benefit analysis in 2009–10 (Piasente et al. 2012). E-monitoring became compulsory for all ETBF longline vessels from 1 July 2015. E-monitoring replaced human observers in the ETBF for all in-zone observer requirements, although the government maintains the right to place human observers on board vessels if there is a need to do so. At least 10% of the video footage from all hauls is reviewed to verify the accuracy of logbooks which are required to be completed for 100% of shots. This review rate may be increased in some cases. In 2021, the coverage rate was 9.9% (Table 11). During the reporting period, Australian vessels operated principally in Australian waters with occasional forays onto the high seas. Australia's national observer program is accredited under the WCPFC ROP.

**Table 11. Summary of longline observer coverage (by hooks) for 2021.**

CCM Fleet	Fishery	No. of Hooks			Days fished			Days at sea			No. of trips			See notes
		Total est.	Obs.	Obs %	Total est.	Obs.	Obs %	Total est.	Obs.	Obs %	Total est.	Obs.	Obs %	
Australia	Domestic	7.4 million	743,779	9.9										Nil

Note: Blacked-out cells are not applicable.

### 8.3 Port sampling program

The collection of individual processed fish weights from processors receiving longline caught fish from the ETBF commenced in mid-1997. The program mainly focuses on the five principal target species in the fishery (yellowfin tuna, bigeye tuna, albacore, swordfish and striped marlin), though data on a range of other species have also been collected. During the period from July 1997 to December 2021 over 2.2 million individual fish weights pertaining to the five main target species have been collected. Coverage rates (% of landed fish sampled) for the target

species are generally high, and for the 5-year period between 2017 and 2021 have averaged around 87% for yellowfin tuna, 93% for bigeye tuna, 92% for swordfish and 94% for striped marlin, while for albacore 4% of landed fish have been individually sampled (Table 12). Individual fish weights for another 289,980 fish from 51 non-target species have also been collected. Bulk weights for albacore sampled in batches (covering on average 88% of the catch between 2017 and 2021) and some other species are also collected. Where both the total batched weight and the number of fish have been recorded, these bulk weights cover 1,356,275 fish for 37 species (including 920,742 albacore).

**Table 12. Estimated annual coverage of operational catch and effort (logbooks), observer data (% hooks) and port sampling (% coverage rate for the five main target species of individual fish weights collected from processors receiving longline-caught fish in the ETBF) for the Australian fleet active in the WCPFC Convention Area, 2017–21.**

Gear	Year	Operational catch & effort coverage (%)	Observer coverage (%)	Port sampling coverage (%)					
				YFT	BET	ALB	SWO	STM	SKJ
Longline <sup>a</sup>	2017	100	10.2	88	92	8	87	90	0
	2018	100	10.8	74	91	3	96	91	0
	2019	100	11.7	91	100	4	95	100	0
	2020	100	9.7	88	96	3	94	93	0
	2021	100	9.9	91	91	3	92	94	0
Purse seine	2017	NA	NA	NA	NA	NA	NA	NA	NA
	2018	NA	NA	NA	NA	NA	NA	NA	NA
	2019	NA	NA	NA	NA	NA	NA	NA	NA
	2020	NA	NA	NA	NA	NA	NA	NA	NA
	2021	NA	NA	NA	NA	NA	NA	NA	NA

<sup>a</sup> includes fish taken by minor line

Abbreviations: yellowfin tuna (YFT), bigeye tuna (BET), albacore (ALB), striped marlin (STM), swordfish (SWO) and skipjack (SKJ)

Sources: Ashley Williams (CSIRO) and AFMA observer database

## 8.4 Unloading/transshipment

Catch disposal records are the formal method for monitoring unloads and were implemented in the ETBF in January 2006 (Table 13). Catch disposal records are completed by both the fisher and licensed fish receiver at the point of unload to obtain accurate data on fish numbers and verified weight by species. Skippers tend to under-estimate the weights reported in logbooks for most species, so the catch disposal record data have been reported in domestic official statistics since 2007. Compliance checks are conducted on unloads as part of a risk-based compliance programme. Weight estimates are also derived from the size-monitoring program and are likely to be more accurate than logbook data for that part of the time series.

**Table 13. Annual catch estimates (converted whole weights) for the ETBF for 2017–21 derived from catch disposal records. Estimates are in tonnes.**

Year	Albacore	Yellowfin	Bigeye	Striped marlin	Swordfish	Other	Total
2017	992.2	1714.5	449.6	287.9	1179.9	1042.2	5666.3
2018	889.1	1516.5	367.5	245.6	1026.9	1448.0	5493.6
2019	923.8	2089.0	284.5	250.8	793.1	1133.9	5475.0
2020	1176.8	1856.4	308.2	204.4	610.6	1082.3	5238.7
2021	1094.7	1592.2	390.4	207.1	618.8	1244.7	5148.0

No transshipment activities were undertaken in 2021 (Tables 14 and 15). No Australian vessels were authorised to tranship on the high seas.

**Table 14. Transshipment quantities by weight for Australian longline vessels in 2021 within the WCPFC Convention Area.**

	Location	Transhipped within Convention Area (CA)?	Caught within Convention Area (CA)?	Product Form? *	Fishing gear	Species	Tonnes
Offloaded	Nil	Nil	Nil	Nil	Longline	Nil	0

**Table 15. Number of transshipment for Australian longline vessels in 2021.**

	Transhipped within Convention Area (CA)?	Caught within Convention Area (CA)?	Fishing gear	Number of Transshipments
Offloaded	Nil	Nil	Longline	0

## 8.5 Other

AFMA introduced the compulsory requirement for all Commonwealth endorsed fishing vessels to be fitted with Integrated Computer Vessel Monitoring Systems (ICVMS) in 2007. For 2021, no ETBF vessel received formal warnings for non-operational ICVMS. There were no other ICVMS infringements. Compliance with ICVMS requirements has increased markedly since mid-2008. AFMA uses the ICVMS to assist in planning inspections and operations, to assist the observer program in deploying scientific observers and to actively monitor compliance with closed areas.

A range of data is also collected via individual research projects (see the Research Activities section for more information).

# 9 Research activities

The Australian Government and the fishing industry allocate considerable funds to fishery research and monitoring each year. In addition to the logbook and observer programs, key areas of research in past years and ongoing research include:

## 9.1 Biological research projects

- Reproductive dynamics of swordfish in the domestic longline fishery off eastern Australia (Young & Drake 2002)
- Age and growth of bigeye tuna from the eastern and western AFZ (Farley et al. 2003)
- Age and growth of swordfish from Australian waters (Young & Drake 2004)
- Population biology and habitat preferences of striped marlin in eastern Australia (Keller & Davie 2009)
- Population biology of albacore tuna in the Australian region (Farley et al. 2012)
- Spatial dynamics of swordfish in the south Pacific Ocean (Evans et al. 2012)
- Defining regional connections in southwest Pacific swordfish (Wilcox 2012)
- Determination of swordfish growth and maturity relevant to the southwest Pacific stock (Farley et al. 2016)
- Age, growth and maturity of bigeye tuna in the western and central Pacific Ocean (CSIRO, July 2016–ongoing)
- Yellowfin tuna age and growth in the Western and Central Pacific Ocean (CSIRO, January 2018–December ongoing)
- Feasibility of Close-Kin Mark-Recapture assessment for South Pacific albacore in the WCPO (CSIRO & SPC, 2021–22)
- Bomb radiocarbon age validation for bigeye and yellowfin tunas in the WCP (Allen Andrews & CSIRO, 2021)

## 9.2 Assessment-related research projects

- Dynamics of the interactions of the fishery and swordfish on seamounts off eastern Australia (Campbell & Hobday 2003)
- Migration and habitat preferences of bigeye tuna on the east coast of Australia (Gunn et al. 2005)
- Stock assessment of striped marlin in the south-western Pacific Ocean (Langley et al. 2006)
- Developing harvest strategies for the ETBF (AFMA 2007)
- Developing robust stock-status indicators (Basson & Dowling 2008)

- Updating the stock assessment of swordfish in the south Pacific Ocean (Kolody, Campbell & Davies 2008)
- Determining the depths fished and the effective longline effort targeted at various species in the ETBF (Campbell & Young 2010)
- Integrated evaluation of management strategies for tropical multi-species long-line fisheries (Kolody et al. 2010)
- Eastern Tuna and Billfish Fishery size monitoring program (Williams et al. ongoing)
- Integrated analysis and assessment supporting implementation of the management and harvest strategy framework within the ETBF (Campbell 2011)
- Predicting the impact of hook decrementation on the distribution of fishing effort in the ETBF (Wilcox et al. 2011)
- Analysis of recreational fishing catch and effort data to support the striped marlin stock assessment (Ghosn et al. 2012)
- Standardisation of commercial catch and effort data to support the stock assessment of striped marlin (Campbell 2012)
- Changes in fishing strategies in the ETBF in response to the introduction of quota management (Preece, Cambell & Hillary 2012)
- Development of an approach to harvest strategy management of internationally managed multi-species fisheries (Hillary et al. 2016)
- Developing innovative approaches to improve CPUE standardisation for Australia's multi-species longline fisheries (Campbell et al. 2017)
- Data management, assessment and implementation of harvest strategies for Australia's tropical tuna fisheries (Campbell 2017)
- Determination of the spatial dynamics and movement rates of the principal target species within the Eastern Tuna and Billfish Fishery and connectivity with the broader western and central Pacific Ocean - beyond tagging (CSIRO; draft report submitted)
- Where have all the yellowfin tuna gone? Investigating the disjunct between commercial and recreational catches off NSW (Pepperell Research & Consulting Pty Ltd/CSIRO, July 2017 – in prep)
- Data management, assessment and implementation of harvest strategies for Australia's tropical tuna fisheries (CSIRO, August 2017– ongoing)
- ETBF striped marlin harvest strategy revision and evaluation (CSIRO, November 2018– ongoing)
- Management strategy evaluation of the broadbill swordfish ETBF harvest strategies (Hilary 2020)
- Data management, assessment and implementation of harvest strategies for Australia's tropical tuna fisheries (CSIRO, July 2020–May 2023)

### 9.3 Ecological research projects

- Ecological risk assessment for the effects of fishing (Webb et al. 2007)
- Rapid quantitative assessment (Zhou, Smith & Fuller 2007)
- Determining the ecological impacts of longline fishing in the ETBF (Young et al. 2009)
- Ecological risk assessment for the effort of fishing report for the Eastern Tuna and Billfish Fishery: Longline sub-fishery, data since 2015 (Sporcic et al. 2019)
- Investigation of oceanographic and environmental factors impacting on the ETBF (CSRIO, ongoing)

### 9.4 Bycatch research projects

- A review of byproduct interactions and economics in Australia's tuna and billfish fisheries (Bromhead et al. 2005)
- Marine turtle mitigation in Australia's pelagic longline fishery (Robins et al. 2007)
- The effects of bycatch mitigation measures, such as circle hooks and wire leaders, on target and non-target catches (Ward et al. 2008)
- Effect of line shooters on the sink rates of pelagic longlines and the effect on seabird interactions (Robertson, Candy & Wienecke 2010a)
- Factors affecting the sink rates of baited hooks and the impact on seabird interactions in pelagic longline fisheries (Robertson & van den Hoff 2010; Robertson et al. 2010b; Robertson & Candy 2013)
- Branch line weighting regimes to reduce the incidental catch of seabirds in pelagic longline fisheries (Robertson et al. 2013)
- The effects of propeller turbulence on sink rates of baited hooks (Robertson & Candy 2014)
- Development of an underwater bait setting system (Robertson et al. 2015)
- Improving the effectiveness, efficiency and safety of mitigation tools for protected species interactions in the Eastern Tuna and Billfish Fishery (FRDC 2020-041, 2020, ongoing).
- Improving the management of wildlife interactions in pelagic longline fisheries (FRDC 2021-078, 2021, ongoing).

# Appendix A: Common and Scientific Names

Common names	Scientific names
Albacore	<i>Thunnus alalunga</i>
Albatrosses (other)	<i>Diomedeidae</i> spp.
Australian fur seal	<i>Arctocephalus pusillus doriferus</i>
Australian gannet	<i>Morus serrator</i>
Australian sea lion	<i>Neophoca cinerea</i>
Baleen whale	Mysticeti
Bigeye tuna	<i>Thunnus obesus</i>
Birds	Aves
Black marlin	<i>Makaira indica</i>
Black-browed albatross	<i>Thalassarche melanophrys</i>
Blacktip sharks	<i>Carcharhinus</i> spp.
Blue marlin	<i>Makaira mazara</i>
Blue shark	<i>Prionace glauca</i>
Bottlenose dolphin	<i>Tursiops truncatus</i>
Bronze whaler shark	<i>Carcharhinus brachyurus</i>
Buller's albatross	<i>Thalassarche bulleri</i>
Cape petrel	<i>Daption capense</i>
Common dolphin	<i>Delphinus delphis</i>
Cormorants	<i>Phalacrocoracidae</i>
Dolphin	Delphinidae
Dugong	<i>Dugong dugon</i>
Dusky shark	<i>Carcharhinus obscurus</i>
Escolar	<i>Lepidocybium flavobrunneum</i>
False killer whale	<i>Pseudorca crassidens</i>
Flatback turtle	<i>Natator depressa</i>
Flesh-footed shearwater	<i>Puffinus carneipes</i>
Great crested tern	<i>Sterna bergii</i>
Great skua	<i>Catharacta skua</i>
Great-winged petrel	<i>Pterodroma macroptera</i>
Green turtle	<i>Chelonia mydas</i>
Grey-headed albatross	<i>Thalassarche chrysostoma</i>
Hammerhead shark	<i>Sphyrna</i> spp.
Hawksbill turtle	<i>Eretmochelys imbricata</i>
Humpback whale	<i>Megaptera novaeangliae</i>
Lancetfish	<i>Alepisaurus</i> sp.
Leatherback turtle	<i>Dermochelys coriacea</i>
Loggerhead turtle	<i>Carretta carretta</i>
Long-finned pilot whale	<i>Globicephala melas</i>
Longfin mako	<i>Isurus paucus</i>
Mahi mahi	<i>Coryphaena hippurus</i>
Melon-headed whale	<i>Peponcephala electra</i>
Moonfish (opah)	<i>Lampris guttatus</i>
New Zealand fur seal	<i>Arctocephalus fosteri</i>
Northern bluefin tuna	<i>Thunnus orientalis</i>
Ocean sunfish	<i>Mola mola</i>
Oceanic whitetip shark	<i>Carcharhinus longimanus</i>
Oilfish	<i>Ruvettus pretiosus</i>
Pacific (olive) ridley turtle	<i>Lepidochelys olivacea</i>
Petrels, prions and shearwaters	<i>Procellariidae</i> spp.
Porbeagle	<i>Lamna nasus</i>
Ray's bream	<i>Brama brama</i>

Rudderfish	<i>Centrolophus niger</i>
Sailfish	<i>Istiophorus platypterus</i>
Scalloped hammerhead	<i>Sphyrna lewini</i>
Seals	Phocidae
Shearwaters	<i>Puffinus</i> spp.
Shortbill spearfish	<i>Tetrapturus angustirostris</i>
Shortfin mako	<i>Isurus oxyrinchus</i>
Short-finned pilot whale	<i>Globicephala macrorhynchus</i>
Short-tailed shearwater	<i>Puffinus tenuirostris</i>
Shy albatross	<i>Thalassarche cauta</i>
Silky shark	<i>Carcharhinus falciformis</i>
Skipjack tuna	<i>Katsuwonus pelamis</i>
Smooth hammerhead	<i>Sphyrna zygaena</i>
Sooty shearwater	<i>Puffinus griseus</i>
Southern bluefin tuna	<i>Thunnus maccoyii</i>
Southern royal albatross	<i>Diomedea epomophora</i>
Striped marlin	<i>Kajikia audax</i>
Swordfish	<i>Xiphias gladius</i>
Thresher shark	<i>Alopias vulpinus</i>
Tiger shark	<i>Galeocerdo cuvier</i>
Toothed whale	Odontoceti
Turtles	Testudinata
Wahoo	<i>Acanthocybium solandri</i>
Wandering albatross	<i>Diomedea exulans</i>
Wedge-tailed shearwater	<i>Puffinus pacificus</i>
Westland petrel	<i>Procellaria westlandica</i>
Whale shark	<i>Rhincodon typus</i>
Wilson's storm petrel	<i>Oceanites oceanicus</i>
Yellowfin tuna	<i>Thunnus albacares</i>
Yellow-nosed albatross	<i>Thalassarche chlororhynchos</i>

---

# Appendix B: Mandatory mitigation measures in the ETBF 2022

*(Source: AFMA website:*

[https://www.afma.gov.au/sites/default/files/final\\_2022\\_etbf\\_management\\_arrangements\\_booklet.pdf](https://www.afma.gov.au/sites/default/files/final_2022_etbf_management_arrangements_booklet.pdf)

## Seabirds

At all times you must:

- Carry one or more assembled tori lines on board
- Not discharge offal while setting
- Carry at least three seabird feather kits onboard
- Comply with seabird interaction obligations relating to the Threat Abatement Plan (TAP).

When you are fishing south of 25°S you must:

- Deploy a tori line before commencing a shot when fishing between the hours of nautical dawn and nautical dusk
- A tori line if not required to be deployed when performing fishing operations between the hours of nautical dusk and nautical dawn
- Use only non-frozen bait
- Weight longlines with either a minimum of:
  - 60 g swivels at a distance of no more than 3.5 m from each hook ; or
  - 98 g swivels at a distance of no more than 4 m from each hook; or
  - 40 g weights immediately adjacent to the hook, or at no more than 0.5 m from the hook, with dead, non-frozen baits attached to the hooks or
  - 'hook shielding device' with a cap and weighing at least 38 g may be deployed directly at the hook as an alternative.

Note: If you are fishing south of 40° South, AFMA may require you to implement additional seabird mitigation measures as this is an area in which higher than average number of seabird interactions are possible.

Your tori line must be:

- At least 100 m long

- Set up from a position on the boat that allows it to stay above the water for at least 75 m from the stern
- Have streamer attached at a maximum interval of 3.5 m
  - Streamers should be maintained, ensuring that their lengths are as close to the water as possible.
- Have a towed line, material or object at the end of the line to give sufficient drag to meet the 75 m aerial coverage criteria.

### **Seabird Interaction Obligations**

As of the 2020 fishing season, the requirements relating to breaches of the Seabird Threat Abatement Plan (TAP) have changed. If a vessel exceeds the seabird bycatch rate (0.05 birds per 1,000 hooks) during any two of the last three consecutive TAP seasons, interacted with more than 10 seabirds in the current or previous TAP season, or been found to have unreported Seabird interaction the following seabird mitigation options must immediately be implemented:

- a daylight setting ban (ensure that all longline hooks are deployed only during the hours between nautical dusk and nautical dawn); or
- implement amended line weighting of either:
  - 40g or greater attached within 0.5 metre of the hook; or
  - 60g or greater attached within 1 metre of the hook; or
  - 80 g or greater attached within 2 m of the hook, or
- ACAP approved hook shielding devices on all hooks; or
- a northern shift in the area of operation (to at least 5 degrees north of the most northerly seabird interaction with the nominated vessel, as verified and notified by AFMA).

If, after implementing one of the seabird mitigation options listed above, a vessel subsequently catches one additional albatross (dead) or two other seabirds (dead) one of the following must immediately be implemented:

- an additional mitigation measure from the list of seabird mitigation options above; or
- cease the use of live bait (if using) and only use dead bait; or
- relocate fishing activities to north of latitude 25 degrees south; or
- cease fishing using longlines for the remainder of the TAP season.

## **Sea turtles**

### **Circle hooks**

Large circle hooks must be used for shallow sets with less than 8 hooks per basket.

### **De-hooking device**

At all times you must carry on board a minimum of one de-hooking device, with the following specifications:

- The device must enable the hook to be secured and the barb shielded so that the barb does not re-engage with the fish while the hook is being removed
- The device must be blunt with all edges rounded
- Where more than one size of hook is to be carried, a de-hooking device (or devices) must be carried that can be used with all hooks on the boat; and
- The shaft of the device must be a minimum of 1.5 metres in length.

### **Line cutting device**

At all times you must carry on board a minimum of one line cutting device. The line cutting device must be constructed and used in accordance with the following specifications:

- The device must be constructed to allow the line to be cut as close to the hook as possible
- The blade of the device must be enclosed in a blunt rounded (arc-shaped) cover with the hook exposed on the inside of the arc; and
- The shaft of the device must be a minimum of 1.5 metres in length.

# References

- AFMA 2007, *Eastern Tuna and Billfish Fishery Harvest Strategy Framework*, Australian Fisheries Management Authority, Canberra.
- Basson, M & Dowling, NA 2008, *Development of a robust suite of stock status indicators for the Southern and Western and the Eastern Tuna and Billfish Fisheries*, CSIRO Marine and Atmospheric Sciences, Hobart.
- Bromhead, D, Ackerman, J, Graham, S, Wight, M, Wise, B & Findlay, J 2005, *Byproduct: catch, economics and co-occurrence in Australia's pelagic longline fisheries*, Bureau of Rural Sciences, Canberra.
- Campbell, RA 2011, *Integrated analysis and assessment supporting implementation of the management and harvest strategy framework within the Eastern Tuna and Billfish Fishery - Compilation of Related Project Papers*, Final report to the Australian Fisheries Management Authority, Canberra.
- Campbell, R 2012, *Aggregate and size-based standardised CPUE indicators for longline target species caught in the south-west Pacific*, WCPFC-SC8-2012/SA-IP-13, Scientific Committee, Eighth Regular Session, 7–15 August 2012, Korea.
- Campbell, R 2017, *Data management, assessment and implementation of the harvest strategies for Australia's tropical tuna fisheries - Compilation of Related Project Papers*, Final report for project R2014/0822 to the Australian Fisheries Management Authority, Canberra.
- Campbell, R & Hobday, A 2003, *Swordfish-environment-seamount-fishery interactions off eastern Australia*, Report to AFMA, Canberra.
- Campbell, R & Young, J 2010, *Determination of effective longline effort in the Eastern Tuna and Billfish Fishery*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Rep. No. 2005/005, Hobart.
- Campbell, RA, Zhou, S, Hoyle, SD, Hillary, R, Haddon, M & Hall, S 2017, *Developing innovative approaches to improve CPUE standardisation for Australia's multispecies pelagic longline fisheries*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Rep. No. 2014-021, Canberra.
- Evans, K, Kolody, D, Abascal, F, Holdsworth, J & Maru, P 2012, *Spatial dynamics of swordfish in the south Pacific Ocean inferred from tagging experiments*, working paper WCPFC-SC8-2012, Scientific Committee Eighth Regular Session, 7–15 August 2012, Korea.
- Farley, J, Clear, N, Leroy, B, Davis, T & McPherson, G 2003, *Age and growth of bigeye tuna (*Thunnus obesus*) from the eastern and western AFZ*, CSIRO Marine Research Report to the Fisheries Research and Development Corporation, Project 2000/100, Hobart.
- Farley, JH, Williams, AJ, Davies, CR, Clear, NP, Eveson, JP, Hoyle, S & Nicol, SJ 2012, *Population biology of albacore tuna in the Australian region*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Project 2009/012, Hobart.

Farley, JH, Clear, NP, Kolody, D, Krusic-Golub, K, Eveson, P & Young, J 2016, *Determination of swordfish growth and maturity relevant to the southwest Pacific stock*, Final report to the Australian Fisheries Management Authority, Project R-2014/ 821, Canberra.

Ghosn, D, Collins, D, Baiada, C & Steffe, A 2012, *Catch per unit effort and size composition of striped marlin caught by recreational fisheries in southeast Australian waters*, WCPFC-SC8-2012/SA-IP-7, Scientific Committee, Eighth Regular Session, 7–15 August 2012, Korea.

Gunn, J, Hampton, J & Evans, K 2005, *Migration and habitat preferences of bigeye tuna, *Thunnus obesus*, on the east coast of Australia—a project using archival and conventional tags to determine key uncertainties in the species stock structure, movement dynamics and CPUE trends*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Project 1999/109, Hobart.

Hillary, R, Preece, A, Kolody, D, Evans, K, Davies, C 2016, *Development of an approach to harvest strategy management of internationally managed multi-species fisheries*, Final report for project R2013/203 to the Fisheries Research and Development Corporation, Canberra.

Keller, KR & Davie, P 2009, *Population biology and habitat preferences of striped marlin, *Kajikia audax*, in the southwest Pacific Ocean*, report for New South Wales Fisheries and the Australian Fisheries Management Authority, Canberra.

Kolody, D, Campbell, R & Davies, N 2008, *A MULTIFAN-CL assessment of south-west Pacific Swordfish*, working paper WCPFC-SC4-2008/SA-WP-6, Scientific Committee Fourth Regular Session 11–22 August 2008, Papua New Guinea.

Kolody, DS, Preece, AL, Davies, CR, Hartog, JR & Dowling, NA 2010, *Integrated evaluation of management strategies for tropical multispecies long-line fisheries*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Project 2007/017, Hobart.

Langley, A, Molony, B, Bromhead, D, Yokawa, K & Wise, B 2006, *Stock assessment of striped marlin (*Tetrapturus audax*) in the south west Pacific Ocean*, working paper WCPFC-SC2-2006/SA WP-6, Scientific Committee Second Regular Session 7–18 August 2006, Philippines.

Piasente, M, Stanley, B, Timmiss, T, McElderry, H, Pria, M, & Dyas, M 2012, *Electronic onboard monitoring pilot project for the Eastern Tuna and Billfish Fishery*, FRDC Project 2009/048, Australian Fisheries Management Authority, Canberra.

Preece, A, Campbell, R & Hillary, R 2012, *Investigation of possible changes in fishing strategies under quota management and implication for the ETBF harvest strategy*, CSIRO Marine and Atmospheric Research Report to the Australian Fisheries Management Authority, Canberra.

Robertson, G & van den Hoff, J 2010, *Static water sink rate trials to improve understanding of sink rates estimated at sea*, Third meeting of the Seabird Bycatch Working Group, SBWG-3 Doc 31, Agreement on the Conservation of Albatrosses and Petrels, 8–9 April 2010, Mara del Plata, Argentina.

Robertson, G, Candy, SG & Wienecke, B 2010a 'Effect of line shooter and mainline tension on the sink rates of pelagic longlines and implications for seabird interactions', *Aquatic Conservation: Marine Freshwater Ecosystems*, vol. 20, pp. 419–427.

Robertson, G, Candy, SG, Wienecke, B & Lawton, K 2010b, 'Experimental determinations of factors affecting the sink rates of baited hooks to minimize seabird mortality in pelagic longline fisheries', *Aquatic Conservation: Marine Freshwater Ecosystems*, vol. 20, pp. 632–643.

Robertson, G, Candy, SG & Hall, S 2013, 'New branch line weighting regimes to reduce the risk of seabird mortality in pelagic longline fisheries without affecting fish catch', *Aquatic Conservation: Marine Freshwater Ecosystems*, DOI: 10.1002/aqc.2346.

Robertson, G & Candy, SG 2014, 'Does propeller turbulence affect the sink rate of baited hooks and their availability to seabirds in pelagic longline fisheries?', *Aquatic Conservation: Marine Freshwater Ecosystems* vol. 24, pp. 179-191.

Robertson, G, Ashworth, Phillip, Ashworth, Peter, Carlyle I & Candy SG 2015, 'The development and operational testing of an underwater bait setting system to prevent the mortality of albatrosses and petrels in pelagic longline fisheries', *Open Journal of Marine Science* vol. 5, pp. 1-12.

Robins, CM, Bradshaw, EJ & Kreutz, DC 2007, *Marine turtle mitigation in Australia's pelagic longline fisheries*, Fisheries Research and Development Corporation Final Report 2003/013, Canberra.

Sporcic, M, Hobday, A, Bulman, C, Hartog, J & Fuller, M 2019, *Ecological risk assessment for the effects of fishing report for the Eastern Tuna and Billfish Fishery: longline sub-fishery, data to 2015*, report for the AFMA, Canberra.

Ward, P, Epe, S, Kreutz, D, Lawrence, E, Robins, C & Sands, A 2008, *Implementation of bycatch mitigation measures in Australia's pelagic longline fisheries: the effects of circle hooks on target and non-target catches*, Final Report to the Natural Heritage Trust, Bureau of Rural Sciences, Canberra.

Webb, H, Hobday, A, Dowdney, J, Bulman, C, Sporcic, M, Smith, T, Stobutzki, I, Fuller, M & Furlani, D 2007, *Ecological risk assessment for the effects of fishing: Eastern Tuna & Billfish Fishery: longline sub-fishery*, report for the AFMA, Canberra.

Wilcox, C 2012, *Defining regional connections in southwestern Pacific broadbill swordfish*, FRDC 2007/036, draft final report, Canberra.

Wilcox, C, Dowling, N & Pascoe, S 2011, *Predicting the impact of hook decrementation on the distribution of fishing effort in the ETBF*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Project 2008/028, Hobart.

Young, J & Drake, A 2002, *Reproductive dynamics of broadbill swordfish (*Xiphias gladius*) in the domestic longline fishery off eastern Australia*, FRDC 1999/108 Final Report.

Young, J & Drake, A 2004, *Age and growth of broadbill swordfish from Australian waters*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Project 2001/014, Hobart.

Young, JW, Lansdell, MJ, Hobday, AJ, Dambacher, JM, Griffiths, SP, Cooper, S, Kloser, R, Nichols, PD & Reville, A 2009, *Determining ecological effects of longline fishing in the Eastern Tuna and Billfish Fishery*, CSIRO Marine and Atmospheric Research Report to the Fisheries Research and Development Corporation, Project 2004/063, Hobart.

Zhou, S, Smith, T & Fuller, M 2007, *Rapid quantitative risk assessment for fish species in selected Commonwealth fisheries*, report to the AFMA, Canberra.