



**COMMISSION  
NINETEENTH REGULAR SESSION**

Da Nang, Viet Nam  
27 November – 3 December 2022

---

**Reference Document for the Review of CMM 2015-02 and the Development of  
Harvest Strategies for South Pacific Albacore**

---

**WCPFC19-2022-14**  
11 November 2022

**Secretariat**

**A. INTRODUCTION**

1. The purpose of this paper is to provide a quick reference guide to the latest recommendations of the latest Scientific Committee (SC) and Technical and Compliance Committee (TCC) of relevance to the discussions in support to the review of CMM for South Pacific Albacore (CMM 2015-02), including the development of the WCPFC harvest strategy framework for South Pacific albacore under CMM 2014-06 and the South Pacific Albacore Roadmap Intercession Working Group meeting ([WCPFC19-2022-SPALB RM](#)). This paper also includes the latest stock status, management advice, research recommendations from SC17 (**Attachment 1**).

**B. SCIENTIFIC COMMITTEE RECOMMENDATIONS**

**B.1 Development of the harvest strategy framework for South Pacific albacore tuna**

**1) South Pacific albacore target reference point** (*Paragraphs 80 - 83, SC18 Outcomes Document*)

2. Noting the concerns expressed at WCPFC18 regarding the delayed process to implement an interim TRP adopted in 2018 and the need to achieve a long-term TRP, and the request from WCPFC18 for SC18 to review any additional information on TRPs for South Pacific albacore tuna, SC18 reviewed the information in SC18-MI-WP-04 (*Further analyses to inform discussions on South Pacific albacore objectives and the TRP*).

3. SC18 noted the implications of a potential MP to be developed across the South Pacific, particularly with the areas outside of the WCPFC jurisdiction, and sought advice on how an MP that only applied to the WCPO could be developed. The Scientific Service Provider explained that this could be undertaken in a similar manner as done for skipjack tuna, where fishing in WCPO archipelagic waters is not controlled by the MP. The MP would be designed so it only applied to the WCPO, and not to the EPO.

4. Noting the request for additional catch scenarios to inform management options to clarify management objectives, several CCMs suggested a 10% and 20% reduction in catch from the 2017-2019

baseline for consideration.

5. SC18 recommended forwarding this updated working paper to WCPFC19 for its deliberations on alternative target reference points for south-Pacific albacore tuna.

## **2) South Pacific albacore operating models** (*Paragraphs 84 - 88, SC18 Outcomes Document*)

6. Noting the Commission is scheduled to adopt an MP for South Pacific albacore tuna in 2022, SC18 reviewed the current grid of OMs that has been developed to reflect all important sources of uncertainty and plausible states of nature for South Pacific albacore as outlined in SC18-MI-WP-05 (*Progress update and technical challenges for the South Pacific albacore MSE framework*).

7. SC18 noted the two alternative sets of OMs listed in Table 1 of SC18-MI-WP-05 – one based on the 2018 assessment (WCPO area only) and one based on the 2021 assessment (including the EPO) – but also noted that it was not able to definitively agree on the reference set of OMs for South Pacific albacore tuna because it was necessary for the Commission to decide whether or not to consider the impacts of fishing within the EPO in their decision making. Nevertheless, SC18 agreed to specify an OM grid for both options so there is a clear way forward for this work pending the Commission’s decision.

8. SC18 noted the axes of uncertainty currently outlined in each set of OMs and recommended that additional axes be considered for inclusion in each (if practical). For the 2018 grid a movement axis should be considered, while for the 2021 grid the addition of an axis exploring CPUE uncertainty should be considered. For both grids, axes examining effort creep and hyperstability should be explored.

9. One CCM also noted that both options exhibit some retrospective bias and suggested that adjustment of terminal estimates to account for retrospective bias in projections might be included as another axis of uncertainty (i.e., with or without bias adjustment).

10. SC18 sought advice from WCPFC19 on whether the impacts of fishing within the EPO need to be included in a set of OMs for south Pacific albacore tuna, and recommended that both the Science-Management-Dialogue and the Commission note the further additions recommended to the alternative sets of OMs.

## **3) South Pacific albacore management procedures** (*Paragraphs 89 - 91, SC18 Outcomes Document*)

11. Noting the Commission is scheduled to adopt an MP for South Pacific albacore tuna in 2022, SC18 reviewed the progress on developing and testing MPs for South Pacific albacore tuna as outlined in SC18-MI-WP-05 (*Progress update and technical challenges for the South Pacific albacore MSE framework*).

12. SC18 noted the progress on the development of MPs using model-based approaches (SPiCT<sup>1</sup>) for South Pacific albacore tuna and recommended that candidate HCRs for this species be adapted from those already considered for skipjack tuna.

13. SC18 recommended that both the Science-Management Dialogue and WCPFC19 take note of the progress to date on the development of an MSE framework for South Pacific albacore tuna and that further work is required prior to adoption of an MP.

---

<sup>1</sup> Stochastic Production model in Continuous Time

## **B.2 Review of the South Pacific Albacore Roadmap-IWG activities**

14. The Commission at WCPFC18 provided the following recommendations on Roadmap for effective conservation and management of South P albacore (*Paragraphs 183-184, WCPFC18 Summary Report*):

*183. The Commission noted the report of progress from the SP Albacore Roadmap Working Group (WCPFC18-2021-South Pacific albacore-RM-IWG).*

*184. The Commission noted the priorities for the SP albacore Roadmap IWG for 2022, in particular the development of a Harvest Strategy for SPA or SPA fisheries, the setting of a TAC and allocation, and encouraged CCMs to continue to work cooperatively in the IWG.*

15. The Commission reviewed the CMM for South Pacific Albacore (CMM 2015-02) and provided the following recommendations (*195 - 199, WCPFC18 Summary Report*)

*195 The Commission noted that the review of CMM 2015-02 Conservation and Management Measure for South Pacific Albacore is ongoing as part of the work of the SP Albacore Roadmap IWG. The Commission encouraged the SPA Roadmap IWG to speed up its process and progress its recommendations.*

*196 The Commission noted concerns regarding the delayed process to implement an interim TRP adopted in 2018, and the need to take action to rebuild the stock to support the economic viability of fleets, and achieve a long-term TRP.*

*197 Given the urgency of these concerns, the Commission agreed that until a new CMM for South Pacific Albacore is adopted, each CCM is encouraged to take steps to implement interim controls on South Pacific albacore catch or effort across the convention area south of the equator.*

- i. CCMs are encouraged to limit commercial fishing of South Pacific albacore within EEZs to domestically applied catch or effort limits in recent years.*
- ii. Each CCM is encouraged to ensure that its flagged vessels for this species shall not exceed the high seas catch or effort (such as number of vessels) of South Pacific albacore by its flagged vessels in recent years.*

*198 These interim arrangements do not confer the allocation of rights to any CCM and are without prejudice to any future decisions of the Commission.*

*199 These interim arrangements shall not prejudice the legitimate rights and obligations under international law for small island developing State and Participating Territory CCMs in the Convention Area for whom South Pacific albacore is an important component of the domestic tuna fishery in waters under their national jurisdiction, and who may wish to pursue a responsible level of development of their fisheries for South Pacific albacore in their EEZs or adjacent high seas.*

16. The IWG convened one on-line meeting, chaired by Ms. Neomai Ravitu (Fiji), on 15 July 2022, and SPC provided presentations on iTRP and management objectives for South Pacific albacore, development of management strategy evaluation, and fishing patterns for total allowable catch and allocations. The IWG also considered its work plan to direct future activities. The Chair's Summary of the

IWG meeting is posted at <https://meetings.wcpfc.int/node/16581>.

17. The general outcomes of the IWG meeting include:

- 1) iTRP and management objectives - There were members (PNA) who raised the issue of having more options on the reference period. While the 2017-2019 is ok for this paper, it is important to look at others, suggesting the 2015-2018 period. Another member asked that this needs more time to be absorbed, in the meantime to continue using the current measure with the use of limiting the number of vessels as it stands. SPC advised that there is a need for managers to identify candidate reference period(s) that reflect the types of performance that they desire from the Southern Longline fishery.
- 2) Harvest Strategy (management procedure) - Members raised questions with regards to how an MP can be operated bycatch/effort in a single management control rule, but generally recognized the need to incorporate all fisheries into MP/HCR development. SPC highlighted that the HS process sets the overall harvest level and that this then needs to go through a separate process for developing and designing an allocation framework. It was also noted that those options presented were relatively straightforward and that more complex options could be investigated based on further discussions with members about what they wish to achieve for their fisheries. One member raised that work should focus on the estimation model, candidate the harvest control rule(s), and management strategy evaluation. However, they raised concerns with other high-priority upcoming tasks in 2023 including the review of the TT CMM, taking precedence over SPA work progression.
- 3) TAC - Members raised concerns around the use of recent catches 2017-2019 as the baseline as these average catches represent the highest catches in the time series and a member suggested the proposal to IATTC for EPO SPA management should be taken into account going forward. A member suggested using vessel numbers rather than catch levels to reflect the working in the existing CMM and asked SPC to provide this data, which SPC noted may be found in WCPFC-TCC17-2021-IP07.
- 4) Work Plan - Members suggested that the Work Plan be revised to focus on the development of the MP work and the other issues we have discussed today. It is too ambitious to have a draft CMM for the WCFPC19 to consider. Further notes that the IWG has the responsibility to provide recommendations on the different components which in our view still has a lot of work to be undertaken. Another member suggested that the IWG work plan be aligned with the SPA Harvest Strategy and see how the IWG can progress/support this work.

## **C. TECHNICAL AND COMPLIANCE COMMITTEE RECOMMENDATIONS**

**C.1 Ongoing difficulties related to interpretation of the term “actively fishing for” (and similar terms such as “directed at” and “targeting”) in certain CMMs which continue to present challenges and makes it difficult for TCC to complete the assessments of some obligations during the CMR (paragraphs 140 and 141 of the TCC18 draft Summary Report)**

18. TCC18 noted the longstanding difference in the interpretation of “actively fishing for” and recommended that the Commission resolve to ensure fairness and consistency in the application of the CMM.

19. TCC18 noted the proposal from the EU ([TCC18-2022-DP03](#)) to assist in resolving the interpretation of “actively fishing for” in CMM 2015-02 and recommended that this and other proposals

should be considered further at WCPFC19.

**C.2 CMM 2015-02 South Pacific albacore** (*paragraph 203 of the TCC18 draft Summary Report*)

20. TCC18 noted the update from the Chair of South Pacific Albacore Roadmap Intersessional Working Group (Fiji).

**The Commission for the Conservation and Management of  
Highly Migratory Fish Stocks in the Western and Central Pacific Ocean**

**Scientific Committee  
Seventeenth Regular Session**

Electronic Meeting  
11–19 August 2021

---

**WCPO SOUTH PACIFIC ALBACORE TUNA STOCK ASSESSMENT**  
(Paragraphs 110 – 138, SC17 Summary Report)

---

**PROVISION OF SCIENTIFIC INFORMATION**

1. SC17 accepted as SC17-SA-WP-02 (*Stock assessment of South Pacific albacore*), which described the 2021 stock assessment of South Pacific albacore. An additional three years of data were available since the previous assessment in 2018 that included data to 2016. The new assessment extends through to the end of 2019 and provides the best available scientific information for the purpose of stock assessment determination.

**a) Stock status and trends**

2. The median values of relative recent (2016-2019) spawning biomass depletion ( $SB_{\text{recent}}/SB_{F=0}$ ) and relative recent (2015-2018) fishing mortality ( $F_{\text{recent}}/F_{\text{MSY}}$ ) over the uncertainty grid of 72 models (Table SPA-01) were used to define South Pacific albacore stock status. The values of the upper 90<sup>th</sup> and lower 10<sup>th</sup> percentiles of the empirical distributions of relative spawning biomass and relative fishing mortality from the uncertainty grid were used to characterize the probable range of stock status.

3. A description of the updated structural sensitivity grid used to characterize uncertainty in the assessment is illustrated in Table SPA-01. Tables SPA-02, SPA-03, and SPA-04 show reference points for South Pacific-wide, WCPFC-CA (Convention Area) and IATTC-CA, respectively, including the median values of relative ‘recent’ (2016-2019) and ‘latest’ (2019) spawning biomass depletion ( $SB_{\text{recent}}/SB_{F=0}$ ) and relative recent (2015-2018) fishing mortality ( $F_{\text{recent}}/F_{\text{MSY}}$ ) over the uncertainty grid of 72 models used to define stock status. These values are based on the uncertainty grid with the downweighted SEAPODYM (M2) movement hypothesis. The values of the upper 90<sup>th</sup> and lower 10<sup>th</sup> percentiles of the empirical distributions of relative spawning biomass and relative fishing mortality from the uncertainty grid were used to characterize the probable range of stock status.

4. The spatial structure used in the 2021 stock assessment is shown in Figure SPA-01. Time series of total annual catch by fishing gear over the full assessment period and by region are shown in Figure SPA-02. Estimated annual average recruitment, spawning potential, and total biomass by model region for the diagnostic case model are shown in Figure SPA-03. Estimated trends in spawning potential by region for the diagnostic case are shown in Figure SPA-04, and juvenile and adult fishing mortality rates from the diagnostic model are shown in Figure SPA-05. Time series of estimated spawning potential for the 72 models are shown in Figure SPA-06. Time-dynamic percentiles of depletion ( $SB_t/SB_{t,F=0}$ ) for the 72 models are shown in Figure SPA-07. Majuro and Kobe plots summarizing the results for each of the

72 models in the weighted structural uncertainty grid are shown in Figures SPA-08 and SPA-09 for the ‘recent’ and ‘latest’ periods, respectively.

5. The most influential axis of uncertainty with respect to estimated stock status was movement, where assuming SEAPODYM-derived movement resulted in more pessimistic outcomes.

6. SC17 noted that the median value of relative recent (2016-2019) spawning biomass depletion for South Pacific albacore ( $SB_{2016-2019}/SB_{F=0}$ ) was 0.52 with a 10<sup>th</sup> to 90<sup>th</sup> percentile interval of 0.41 to 0.57.

7. SC17 further noted that there was 0% probability (0 out of 72 models) that the recent (2016-2019) spawning biomass had breached the adopted limit reference point (LRP).

8. SC17 noted that there has been a long-term increase in fishing mortality for adult South Pacific albacore, with a notable steep increase in fishing mortality since 2000.

9. SC17 noted that the median of relative recent fishing mortality for South Pacific albacore ( $F_{2015-2018}/F_{MSY}$ ) was 0.24 with a 10<sup>th</sup> to 90<sup>th</sup> percentile interval of 0.15 to 0.37.

10. SC17 further noted that there was 0% probability (0 out of 72 models) that the recent (2015-2018) fishing mortality was above  $F_{MSY}$ .

11. SC17 noted the results of stochastic projections (based on the weighted grid, SC17-SA-WP-02a, Figures 1 and 2) from the 2021 assessment, which indicated the potential stock consequences of fishing at “status quo” conditions (2017–2019 or 2020 average catch or, separately, fishing effort) using the uncertainty framework approach endorsed by SC17. These results are provided for both South Pacific-wide and for the WCPFC Convention area only. All projections show a steep and rapid decline in biomass towards the LRP in the year 2021 followed by an increase in biomass thereafter.

**Table SPA-01.** Description of the structural uncertainty grid used to characterize uncertainty in the management quantities derived from this assessment. Note that the M2-SEAPODYM hypothesis was downweighted by 50% by the SC17.

Axis	1	2	3
Steepness (S)	0.65	<b>0.80</b>	0.95
Movement (M)	<b>M1-Estimated, age-dependent</b>	M2-SEAPODYM	
Size data weight (D)	Low (50)	<b>Medium (25)</b>	High (10)
Recruitment distribution (R)	<b>R1-SEAPODYM</b>	R2-Regions 3 and 4	
Growth/M (G/M)	<b>Fixed otolith, Nat-M1</b>	Estimated from length frequency, Nat-M2	

**Table SPA-02.** South Pacific-wide (all regions) reference point estimates from the assessment based on the weighted grid.


	Mean	Median	Min	10%	90%	Max
$C_{latest}$	87,184	86,827	83,519	85,092	87,633	130,936
$F_{MSY}$	0.06	0.06	0.05	0.05	0.07	0.08
$f_{mult}$	4.37	4.25	2.11	2.69	6.62	7.84
$F_{recent} / F_{MSY}$	0.25	0.24	0.13	0.15	0.37	0.47
$MSY$	115,661	120,020	68,200	75,584	158,600	166,240
$SB_0$	623,542	660,200	361,800	392,590	845,100	929,300
$SB_{F=0}$	675,861	678,345	524,886	537,740	824,855	873,278
$SB_{latest} / SB_0$	0.41	0.41	0.34	0.37	0.46	0.48
$SB_{latest} / SB_{F=0}$	0.37	0.40	0.25	0.27	0.45	0.46
$SB_{latest} / SB_{MSY}$	2.50	2.33	1.45	1.69	3.921	4.28
$SB_{MSY}$	109,710	104,100	48,040	61,497	157,500	190,000
$SB_{MSY} / SB_0$	0.18	0.18	0.11	0.11	0.22	0.23
$SB_{MSY} / SB_{F=0}$	0.16	0.16	0.09	0.11	0.22	0.23
$SB_{recent} / SB_{F=0}$	0.50	0.52	0.37	0.41	0.57	0.59
$SB_{recent} / SB_{MSY}$	3.34	3.22	2.07	2.24	5.18	5.33
$Y F_{recent}$	81,998	85,020	58,440	63,656	94,720	101,400

**Table SPA-03.** WCPFC-CA reference point estimates from the assessment based on the weighted grid.

	Mean	Median	Min	10%	90%	Max
$C_{latest}$	78,788	78,455	75,673	76,959	79,126	118,706
$SB_{F=0}$	459,648	463,424	415,746	431,617	491,092	501,602
$SB_{latest} / SB_{F=0}$	0.37	0.39	0.26	0.28	0.43	0.45
$SB_{recent} / SB_{F=0}$	0.51	0.52	0.39	0.42	0.58	0.61


**Table SPA-04.** IATTC-CA reference point estimates from the assessment based on the weighted grid.

	Mean	Median	Min	10%	90%	Max
$C_{latest}$	8,396	8,242	7,845	8,074	8,760	12,229
$SB_{F=0}$	216,213	233,755	92,190	98,063	356,491	379,718
$SB_{latest} / SB_{F=0}$	0.38	0.42	0.22	0.25	0.46	0.48
$SB_{recent} / SB_{F=0}$	0.47	0.52	0.28	0.32	0.56	0.57




**Figure SPA-01.** The geographical area covered by the stock assessment and the boundaries of the four model regions used for South Pacific-wide 2021 albacore assessment. The overlap region between the WCPFC and IATTC convention areas is the area between 130° - 150° west demarcated by the dashed line. The catch from the ‘overlap’ area is included within the WCPFC-CA for this assessment.







**Figure SPA-02.** a) Spatial pattern of albacore catch by gear type over the last decade, and b) historical catches of albacore across the model region from 1952-2019 by gear type.




**Figure SPA-03.** Estimated annual average a) spawning potential, b) recruitment, and c) total biomass by model region for the diagnostic case model, showing the relative levels among regions.




**Figure SPA-04.** Estimated temporal spawning potential by model region, grouped by region (WCPFC-CA, EPO) and South Pacific as a whole for the diagnostic case model. The dotted lines are included to indicate the  $SB/SB_{F=0}$  interim target reference point (iTRP)=0.56 and the LRP=0.2 for the WCPFC-CA albacore fishery. Regions 1-3 represent the WCPFC-CA (including the “overlap”), Region 4 is the IATTC-CA.




**Figure SPA-05.** Estimated annual average juvenile and adult fishing mortality for the diagnostic case model.




**Figure SPA-06.** Estimated spawning potential across all models in the structural uncertainty grid over the period 1960-2019. The dashed line represents the median. The darker band shows the 10<sup>th</sup>-90<sup>th</sup> percentile, and the lighter band shows the 25<sup>th</sup>-75<sup>th</sup> percentile of the model estimates. Regions 1-3 represent the WCPFC-CA (including the “overlap”), Region 4 is the IATTC-CA. The bars at right in each plot are the median values (points) and percentiles for recent (left) and latest (right) spawning potential.



**Figure SPA-07.** Estimated spawning depletion across all models in the structural uncertainty grid over the period 1960-2019. The dashed line represents the median. The darker band shows the 10<sup>th</sup>-90<sup>th</sup> percentile, and the lighter band shows the 25<sup>th</sup>-75<sup>th</sup> percentile of the model estimates. Regions 1-3 represent the WCPFC-CA (including the “overlap”), Region 4 is the IATTC-CA. The dashed horizontal lines indicate the depletion LRP (0.2) and the WCPFC-CA TRP for  $SB/SB_{F=0}$  (0.56). The bars at right in each plot are the median values (points) and percentiles for  $SB_{recent}/SB_{F=0}$  (left) and  $SB_{latest}/SB_{F=0}$  (right).



**Figure SPA-08.** Majuro (bottom) and Kobe (top) plots summarizing the Pacific-wide results for each of the models in the structural uncertainty grid for the ‘recent’ (2016-2019) period. The blue point is the median value based on the weighted grid models, with the more heavily weight models indicated by the larger black dots.



**Figure SPA-09.** Majuro (bottom) and Kobe (top) plots summarizing the Pacific-wide results for each of the models in the structural uncertainty grid for the ‘latest’ (2019) period. The blue point is the median value based on the weighted grid models, with the more heavily weighted models indicated by the larger black dots.

***b. Management advice and implications***

12. Annual catch estimates for albacore in the South Pacific peaked at 93,835 mt (all gears) in 2017 (SC17-SA-IP-04). Catch by longliners represented 93% of the catch weight in 2020 at 64,963 mt and represented a 21% decrease from 2019 despite a shift of effort from the tropical to the southern longline fishery in 2020. By comparison, the 2020 total albacore catch within the southern part of the WCPFC-CA was 61,778 mt and the longline catch was 57,006 mt.

13. The 2021 South Pacific albacore stock assessment provided results consistent with the 2018 assessment. The addition of the EPO region into the current entire South Pacific assessment did not notably alter the main assessment outcomes, and similar trajectories and terminal depletion were estimated in both RFMO regions.

14. The spawning stock biomass has become more depleted across the model period (1960-2019), with a notable increase in depletion in the most recent years. Based on the set of models in the SC endorsed structural uncertainty grid, the South Pacific albacore assessment indicates the stock is not overfished, and there was zero estimated risk of the stock being below the Limit Reference Point of  $20\%SB_{F=0}$ . However, the decline in the latest estimated  $SB_{latest}/SB_{F=0}$  (year 2019; median 0.40; 10<sup>th</sup> and 90<sup>th</sup> percentiles 0.27 - 0.45) is notably more pessimistic than those of  $SB_{recent}/SB_{F=0}$  (years 2016-2019; median 0.52; 10<sup>th</sup> and 90<sup>th</sup> percentiles 0.41 - 0.57) indicating that there has been a substantial decline in stock status estimated over the last three years. The general trends are consistent for estimates across all regions of the South Pacific stock, and for the WCPFC-CA only.

15. For the WCPFC-CA region, the 'recent' and 'latest' SB estimates are on average both below the interim TRP of 0.56. Further, 86% of models (62 out of 72 models) in the structural uncertainty grid endorsed by SC17 estimated that  $SB_{recent}/SB_{F=0}$  was below the interim TRP. In relation to management objectives for the WCPFC-CA longline fishery, this assessment estimated that the median 'latest' (2019) and 'recent' (2016-2019) longline vulnerable biomass for the WCPFC-CA are 56% and 76% of the 2013+8% target level that defined the interim TRP.

16. SC17 noted CPUE declines in many domestic longline fisheries in the southern portion of the WCPFC-CA.

17. SC17 noted that depletion is greatest in regions north of 25°S, specifically in assessment Regions 1 and 2 where most domestic Pacific Island Countries and Territories (PICTs) fleets operate, including Small Island Developing States (SIDS) and Participating Territories that may have no high seas access. These are areas mostly unaffected by current management measure for South Pacific albacore (CMM 2015-02), which prescribe effort controls and reporting provisions south of 20°S.

18. SC17 expressed great concern with the projected status of South Pacific albacore if recent catch or effort levels are maintained (SC17-SA-WP-02a REV2). Projections indicated that South Pacific albacore stock has a greater than 20% risk of falling below the LRP in 2021 under both catch and effort scenarios. These projections indicate an extended period where biomass is below the current interim TRP and in most cases the TRP is not achieved within the 30-year projection period.

19. Recalling its previous advice from SC11, SC12, and SC13, SC17 recommended that longline catch be reduced to avoid further and extended declines in the vulnerable biomass so that economically viable catch rates can be maintained, especially for longline catch of adult albacore.

20. SC17 recommended a recalibration of the interim TRP for review at WCPFC18 in accordance with the process agreed at WCPFC15 (WCPFC15 Summary Report, para 207). Further, SC17



recommended projections be undertaken to estimate the constant catch levels that would achieve that TRP on average over the long-term. SC17 recommended that these analyses be provided to WCPFC18 to guide its consideration of reductions in longline fishing mortality that will be required to return the vulnerable biomass to the 2013 +8% level as agreed.

*c. Future research recommendations*

21. SC17 noted with concern that the standardized CPUE indices do not show linear contrast with catches over the past 20 years when the catch has increased by 2 to 3-fold and also that the fit to the indices show a residual pattern over time. SC17 supported the assessment scientist's suggestion to consider split indices in future assessments, which might allow for the incorporation of more informative catchability and density covariates during the contemporary period, which is more important for estimates of recent status.

22. SC17 noted a possible nonlinear relationship between catch and effort or a time-varying relationship with changing fishing power and catchability. The next assessment could investigate such nonlinear relationships and explore alternative effort metrics.

23. SC17 noted with concern that the standardized CPUE model with hooks between floats (HBF) did not converge. The time-series is almost 70 years with substantial shifts to deploy more HBF though time. These gear changes have probably altered South Pacific albacore catchability and require additional research. HBF is one characteristic of longline gear that could affect catchability; operational longline data are largely absent of detailed vessel and gear characteristics that could be valuable in a standardization model. Reliably collecting additional gear characteristics will better inform these models on variability in catchability among vessels and fleets and over time and these data enhancements could be achieved by revisiting the minimum logsheet data standards, increasing observer coverage, or expanding electronic monitoring applications. Without this additional information the large uncertainties associated with the use of standardised-CPUE in assessments will remain unresolved and continue to impact on future assessments.

24. SC17 noted the need to both recalibrate the interim TRP according to the procedure agreed at WCPFC15 (WCPFC15 Summary Report, para 207) and estimate the constant catch levels that would achieve that TRP on average over the long-term. Specifically, based upon the SC-agreed 2021 South Pacific albacore stock assessment:

- a) re-calibrate the WCPFC interim TRP (the median depletion in the WCPFC-CA,  $SB/SB_{F=0}$ ) that would on average achieve the agreed objective of an 8 % increase in vulnerable biomass (CPUE proxy) for the southern longline fishery as compared to 2013 levels.
- b) undertake projections to estimate the constant catch levels that would achieve the recalibrated TRP, on average, over the long-term.
- c) within that projection-based analysis, WCPFC-CA longline and troll fisheries should be modelled based upon catch, and fishing levels within the EPO should be adjusted in the same way as the WCPO for one scenario and fixed at recent catch levels for another scenario. Future recruitment should be sampled from the long-term recruitment pattern.

25. A number of key research needs were identified in undertaking the assessment that should be investigated either internally or through directed research.

26. As with the previous South Pacific albacore assessment, the fishery dependent CPUE-based indices of abundance lacked contrast to inform population responses to increased fishing pressure. This continues to be a significant concern for the reliability of estimates of population size. The CPUE analysis has been a major focus of preparatory work for this and previous assessments, and despite the attempts of

various scientists, application of new approaches including attempts at splitting time series and testing various covariates, the CPUE continues to lack contrast. It is recommended that alternative fishery independent estimates of population size be explored, especially the genetic method of Close-Kin Mark-Recapture (CKMR).

27. The implications of uncertainty in movement were clearly evident in this year's assessment, with this being the most influential uncertainty for management advice. In the absence of strong empirical data to inform decisions on alternative movement hypotheses and based on the quality of fits to the data, the SC decided to downweight one of the two movement hypothesis for provision of management advice. This is an unsatisfactory situation and there is a clear need to improve understanding of connectivity among albacore populations across the South Pacific, and, in particular, the fishery regions in the WCPFC and IATTC convention areas. This is particularly critical if South Pacific-wide assessments are to continue. The CKMR method as a by-product can also address this uncertainty.

28. Despite applying the new growth data to this assessment, the modal structure in the New Zealand troll fishery size composition was still not fit adequately. Further work on growth modelling is required. It should also be noted that otolith-based growth data being used is mostly derived for otolith samples collected in 2009 -2010. Further, to update the growth information for albacore, samples from the IATTC-CA are needed. Again, samples required to address this issue could be collected as part of a CKMR project that would also include a component to develop (tissue-based) epigenetic ageing methods and sex determination. This would be a major advance in including more contemporary growth information in tuna assessments.

29. Follow-up studies to assess the reliability of size composition data for providing information on recruitment and population trends, and if necessary, develop better stratification methods to improve the representativeness of size composition data should be considered.

30. Finally, the current model is highly parameterized, and reducing model parameters and complexity should be considered to improve model fits and diagnostics. One key advancement would be the application of the "catch conditioned" approach that will be available in MULTIFAN-CL for the next assessment.