

SCIENTIFIC COMMITTEE SEVENTH REGULAR SESSION

9-17 August 2011 Pohnpei, Federated States of Micronesia

ANNUAL REPORT TO THE COMMISSION PART 1: INFORMATION ON FISHERIES, RESEARCH, AND STATISTICS

WCPFC-SC7-AR/CCM-02

CANADA

2011 Annual Report to the Western and Central Pacific Fisheries Commission

Canada

PART I. INFORMATION ON FISHERIES, RESEARCH, AND STATISTICS For 2010

John Holmes
Fisheries and Oceans Canada
Science Branch
Nanaimo, British Columbia, Canada

August 2011

| Scientific data was provided to the Commission in accordance with the decision relating to the provision of scientific data to the Commission by 30 April 2011 | YES |
|--|-----|
| | |

Summary

Catch, effort and catch per unit of effort (CPUE) data for the Canadian albacore (*Thunnus alalunga*) fishery in the WCPFC Convention Area for 2010 are summarized in this document. The Canadian tuna fishery is a troll fishery that uses jigs and targets albacore exclusively. The Canadian fishery was inactive within the WCPFC convention area in 2010, with no effort or catch reported in statistical zones within the convention area in either the north or south Pacific Oceans. Annual albacore tuna catches by the Canadian troll fleet in the convention area between 2002 and 2008 have ranged from 83 t in 2005 to 453 t in 2003 and effort has ranged from 56 v-d in 2007 to 408 v-d in 2002. Both catch and effort by the Canadian fleet in the WCPFC convention area have declined since 2002. Two Canadian scientists participated in and Chaired two ISC-Albacore Working Group workshops in 2010.

2011 Annual Report to the Western and Central Pacific Fisheries Commission

Canada

PART I. INFORMATION ON FISHERIES, RESEARCH, AND STATISTICS (For 2010)

Canadian fishermen have been fishing for albacore tuna (*Thunnus alalunga*) since the mid-1930s. The Canadian fishery started in the coastal waters off British Columbia and is a troll fishery using jigs to target albacore tuna in the surface waters of the Pacific Ocean. Although the Canadian fleet will follow albacore tuna concentrations into offshore waters, in recent years the majority of effort and catch has occurred in the coastal waters of Canada and the United States and this trend continued in 2010. Access by Canadian vessels to waters in the U.S. Exclusive Economic Zone (EEZ) is governed by a bilateral Canada-United States albacore tuna treaty, which enables Canadian and U.S. fishers to catch north Pacific albacore in each other's EEZ, and land albacore tuna at designated ports in Canada and the United States.

Canada is committed to providing detailed catch and effort statistics, logbook data, and fishing vessel information, as is required under the Highly Migratory Species Convention. Management regulations for Canadian vessels fishing albacore tuna in 2010 are documented in the *Pacific Region Integrated Fisheries Management Plan: Tuna - April 1, 2010 to March 31, 2011,* which is available electronically at: which is available at http://www.dfo-mpo.gc.ca/Library/340295.pdf. These regulations specify that Canadian fishers must obtain a licence to fish for albacore tuna and that they must maintain accurate records of daily harvest operations in the *Canadian Pacific Albacore Tuna Logbook*. Logbooks are purchased from the Canadian Highly Migratory Species Foundation (http://www.canadianalbacoretuna.com/) and fishers are required to submit their logbooks within 7 days of their final landing or mid-November, whichever is first. The Canadian tuna fishery in the coastal and high seas waters of the Pacific Ocean was open from 01 April 1 2010 to 31 March 2011, but all reported catch and effort by the fleet occurred between 15 June and 31 October 2010.

This report presents summaries of catch, effort and catch per unit of effort (CPUE) data for the Canadian albacore tuna troll fishery in the WCPFC Convention Area in 2010.

1.1 Annual Fisheries Information

Data on albacore tuna catch and effort are compiled from hailing records, logbooks, and sales slips from processing plants and stored in the *Canadian Albacore Tuna Catch and Effort Relational Database* (Stocker et al. 2007). This database contains all fishery-related scientific data from 1995 to the present and provides the best estimate of total annual catch and effort by vessel and geographic area. All fishing vessels are required to hail out when they intend to start fishing and hail in when fishing ceases. Hail data from vessels fishing in Canadian waters are obtained from Marine Communications and Traffic Services, Canadian Coast Guard, and hail

data for vessels fishing in U.S. waters are obtained from Ship.com. The hail data are used to estimate total vessels fishing (Stocker et al. 2007). Canadian vessels must also carry logbooks while fishing for highly migratory species in any waters of the Pacific Ocean. Daily catch and effort data at the highest temporal and spatial scales are obtained from completed copies of the logbooks submitted at the end of the fishing season. A full description of the type of information recorded in the logbooks is provided by Stocker et al. (2007). Sales slips records of landings provide the most accurate estimates of albacore landings (weight), although they underestimate total annual landings because they do not fully account for international sales, domestic public sales or take-home totals (Stocker et al. 2007). Logbooks, sales slips and at-sea trans-shipment slips, completed at the time fish are landed and sold, must be returned to Fisheries and Oceans Canada (DFO) for entry into the albacore catch database (Argue et al. 1999; Stocker et al. 2007). Canada does not currently have a domestic program to collect biological data (lengths, weights, sex) from catch landed in Canadian ports.

The data in this report were obtained from Version 11.03.29 of the *Canadian Albacore Tuna Catch and Effort Relational Database* using codebase Version 11.03.10 to query and compile the data. The 2010 data presented in this report are considered preliminary while the data from 2002 to 2009 are definitive. Similar summaries of data from 2002 to 2008 are provided by Stocker and Shaw (2003, 2004, 2005), Stocker (2006, 2007), and Holmes (2008, 2009, 2010).

1.1.1 Annual Catch in the WCPFC Convention Area

The preliminary Canadian catch of north Pacific albacore tuna in 2010 was 6,497 t (Table 1) in the eastern Pacific Ocean (Table 2). Canadian catches of north Pacific albacore within the WCPFC convention area have ranged between 11 and 1,007 t historically, averaging 431 t for the 1995-2005 period and south Pacific albacore catch has varied between 0 and 313 t, averaging 135 t, from 1996 to 2007 (Figure 1; Table 1). Canadian vessels ceased entering the WCPFC convention area in the north Pacific Ocean after the 2005 fishing season and became inactive in the south Pacific Ocean following the 2007 fishing season. All catches of albacore tuna in the north Pacific reported by the Canadian troll fleet since 2006 (Table 1) occurred east of 150 °W longitude, within the convention area of the Inter-American Tropical Tuna Commission (IATTC). Catch figures in the north Pacific for 2008 and 2009 have been updated (Table 1) relative to previous reports as a result of more complete logbook since the last report was provided to SC (Holmes 2010).

1.1.2 Annual Effort in the WCPFC Convention Area

The number of Canadian vessels operating in the convention area has declined from a peak of 35 in 1997 to zero (0) vessels since 2008 (Table 1) and similarly, fishing effort (measured as the number of vessel fishing days (v-d)) has declined from 1,188 v-d in 1997 to 59 v-d in 2007 and 0 thereafter. Historically, the majority of catch and effort within the convention area occurred in the north Pacific west of 150 °W longitude. Effort figures in the north Pacific for 2008 and 2009 have been updated (Table 1) relative to previous reports as a result of more complete logbook submission since the last report was provided to SC (Holmes 2010).

1.1.3 Annual CPUE in the WCPFC Convention Area

Nominal (reported) catch-per-unit-effort (CPUE) for north Pacific albacore has ranged from a low of 193 kg/v-d in 2005 to a high of 1,653 kg/v-d in 1996 and for south Pacific albacore it has

ranged from 488 kg/v-d in 1996 to 1,504 kg/v-d in 1998. Average annual CPUEs are 1,057 kg/v-d for north Pacific albacore and 893 kg/v-d for south Pacific albacore (Table 1).

1.1.4 Interactions with other Species in the WCPFC Convention Area

There were no reported interactions or bycatch of sharks, seabirds, or sea turtles by the Canadian fishery in the WCPFC convention area in 2010.

1.1.5 Swordfish

Canadian-flagged vessels or Canadian vessels under charter, lease or similar arrangements operating as part of the domestic Canadian fishery did not fish for or catch swordfish (*Xiphias gladius*) south of 20 °S for the 2000-2010 period.

1.1.6 Transhipment

Transhipment activities within the WCPFC convention area were not reported by Canadian vessels in 2010.

1.2 Research and Statistics

1.2.1 Stock assessment studies

Canada currently chairs the ISC Albacore Working Group (ISC-ALBWG) and participated in two workshops in 2010. Results of these workshops, including scientific advice, are contained in Annexes 6 and 9 of the "Report of the Tenth Meeting of the International Committee for Tuna and Tuna-like Species in the North Pacific Ocean", which is available electronically at: http://isc.ac.affrc.go.jp/reports/isc/isc10_reports.html.

1.2.2 Size Sampling of the Catch

Canada implemented an on-board sampling program in 2009. Harvesters were asked to measure and record the lengths (rounding down to the lowest whole number) of the first 10 fish landed on a daily basis, or as often as possible. Thirty-four vessels participated in 2010 and turned in 9,772 fork length measurements of north Pacific albacore, for a sampling rate of 1.04% of the fish caught.

References

- Holmes, J.A. 2010. Annual report to the Commission. Part 1: Information on fisheries, research, and statistics for 2009. Canada. Canadian annual report to the Western and Central Pacific Fisheries Commission, Sixth Regular Session of the Scientific Committee, 10-19 August 2010, Nuku'alofa, Tonga. WCPFC-SC6-AR/CCM-02: 9 p.
- Holmes, J.A. 2009. Annual report to the Commission. Part 1: Information on fisheries, research, and statistics for 2008. Canada. Canadian annual report to the Western and Central Pacific Fisheries Commission, Fifth Regular Session of the Scientific Committee, Port Vila, Vanuatu, 10-21 Aug 2009. WCPFC-SC5-AR/CCM-02: 11 p.
- Holmes, J.A. 2008. Canada. Part I. Annual Report Part 1. Information on fisheries, research, and statistics for 2007. Canadian annual report to the Western and Central Pacific Fisheries Commission, Scientific Committee Fourth Regular Session, Port Moresby, Papua New Guinea, 11-22 Aug 2008. WCPFC-SC4-AR PART 1/WP-03: 10 p.
- Stocker, M. 2007. Canada Part I. Information on fisheries, research, and statistics for 2006. Canadian annual report to the Western and Central Pacific Fisheries Commission, Honolulu, HI, U.S.A., 13-24 Aug 2007. AR WP-3: 5 p.
- Stocker, M. 2006. Canadian albacore tuna fisheries in the north and south Pacific Ocean in 2005. Canadian National Fishery Report for the 2nd Regular Session of the Scientific Committee of the Western and Central Pacific Fisheries Commission, Manila, Philippines, August 7-18, 2006. FR WP-2: 6 p.
- Stocker, M. and W. Shaw. 2005. Canadian albacore tuna fisheries in the north and south Pacific Ocean in 2004. Canadian National Fishery Report for the 1st meeting of the Scientific Committee of the Western and Central Pacific Fisheries Commission, Noumea, New Caledonia, August 8-19, 2005. FR WP-4: 7 p.
- Stocker, M. and W. Shaw. 2004. An update for Canadian tuna fisheries in the north and south Pacific Ocean for 2003. Report prepared for the 17th meeting of the Standing Committee on Tuna and Billfish, Majuro, Marshall Islands, August 9-18, 2004. NFR-3: 9 p.
- Stocker, M. and W. Shaw. 2003. An update for Canadian tuna fisheries in the north and south Pacific Ocean for 2002. Report prepared for the 16th meeting of the Standing Committee on Tuna and Billfish, Mooloolaba, Australia, July 9-16, 2003. NFR-3: 8 p.
- Stocker, M., Stiff, H., Shaw, W., and Argue, A.W. 2007. The Canadian albacore tuna catch and effort relational database. Canadian Technical Report of Fisheries and Aquatic Sciences 2701: vi+76 p. http://www.dfo-mpo.gc.ca/Library/327827.pdf

Table 1. Fisheries statistics for the Canadian troll fleet targeting albacore in the WCPFC convention area, 1995 to 2010. 0 means no reported data. CPUE values are not standardized.

| | North Pacific ^A | | | North Pa | North Pacific – inside WCPFC CA ^{B,C} | | | South Pacific ^C | | | | |
|-------------------|----------------------------|-----------------|--------------------|------------------|--|-----------------|--------------------|----------------------------|--------------|-----------------|--------------------|------------------|
| Year | Catch (t) | Effort (v-d) | Num. of Vessels | CPUE (kg/v-d) | Catch (t) | Effort (v-d) | Num. of Vessels | CPUE (kg/v-d) | Catch (t) | Effort (v-d) | Num. of Vessels | CPUE (kg/v-d) |
| 1995 | 1,763 | 5,930 | 284 | 297 | 23 | 17 | 3 | 1,353 | | | | |
| 1996 | 3,316 | 8,151 | 292 | 407 | 811 | 523 | 25 | 1,551 | 82 | 168 | 3 | 488 |
| 1997 | 2,168 | 4,324 | 197 | 501 | 1,007 | 1,017 | 32 | 990 | 149 | 171 | 3 | 871 |
| 1998 | 4,177 | 6,018 | 214 | 694 | 752 | 455 | 27 | 1,653 | 167 | 111 | 3 | 1,504 |
| 1999 | 2,734 | 6,969 | 233 | 392 | 151 | 327 | 14 | 462 | 254 | 197 | 5 | 1,289 |
| 2000 | 4,531 | 8,769 | 238 | 517 | 586 | 608 | 12 | 964 | 313 | 348 | 5 | 899 |
| 2001 | 5,248 | 10,021 | 244 | 524 | 569 | 383 | 7 | 1,486 | 208 | 168 | 4 | 1,238 |
| 2002 | 5,379 | 8,323 | 228 | 646 | 259 | 250 | 7 | 1,036 | 144 | 158 | 4 | 911 |
| 2003 | 6,861 | 8,429 | 192 | 814 | 453 | 389 | 15 | 1,165 | 0 | 4 | 1 | 0 |
| 2004 | 7,856 | 9,943 | 220 | 790 | 123 | 159 | 5 | 774 | 63 | 67 | 1 | 940 |
| 2005 | 4,845 | 8,565 | 213 | 566 | 11 | 57 | 1 | 193 | 72 | 111 | 2 | 649 |
| 2006 | 5,832 | 6,243 | 174 | 934 | 0 | 0 | 0 | | 135 | 105 | 2 | 1,286 |
| 2007 | 6,075 | 7,113 | 198 | 854 | 0 | 0 | 0 | | 38 | 59 | 1 | 644 |
| 2008 | 5,446 | 5,907 | 134 | 922 | 0 | 0 | 0 | | 0 | 0 | 0 | |
| 2009 | 5,643 | 6,589 | 135 | 856 | 0 | 0 | 0 | | 0 | 0 | 0 | |
| 2010 ^D | 6,497 | 7,532 | 157 | 863 | 0 | 0 | 0 | | 0 | 0 | 0 | |

A – Total catch and effort in the north Pacific, including catch and effort within the WCPFC convention area B – North Pacific albacore catch and effort west of 150 °W longitude (inside the WCPFC convention area).

C – Total catch and effort in the WCPFC convention area are the sum of figures in these columns

D – Preliminary estimates from database version 11.03.29.

Table 2. Aggregated Canadian albacore tuna catch and effort data in the Pacific Ocean for 2010. Statistical areas are defined in Figure 2.

| Area | Catch (t) | Effort (vessel-days) |
|--------------------------------------|-----------|----------------------|
| 1. North Pacific Ocean | 6,497 | 7,532 |
| 2. South Pacific Ocean | 0 | 0 |
| 3. WCPFC Statistical Area N | 0 | 0 |
| 4. WCPFC Statistical Area S | 0 | 0 |
| 5. WCPFC Statistical Area E of 150°W | 0 | 0 |
| 6. IATTC E of 130°W | 0 | 0 |
| | | |






Figure 1. Canadian albacore tuna catch (\bullet) and nominal catch-per-unit-effort, CPUE (\circ) , in the WCPFC convention area north of the equator (A) and south of the equator (B). Note the differences in times scales and vertical scales. Lines are two-year running averages of catch (solid) and CPUE (dashed).




Figure 2. Statistical areas used for reporting of albacore tuna aggregated catch and effort data in the WCPFC convention area. Numbers refer to area catch and effort data shown in Table 2.