

SCIENTIFIC COMMITTEE THIRTEENTH REGULAR SESSION

Rarotonga, Cook Islands 9 – 17 August 2017

Report of the Seventeenth Meeting of the International Scientific Committee for Tuna and Tuna-like Species in the North Pacific Ocean

WCPFC-SC13-2017/ GN-IP-02

ISC¹

¹ International Scientific Committee for Tuna and Tuna-like Species in the North Pacific Ocean



REPORT OF THE SEVENTEENTH MEETING OF THE INTERNATIONAL SCIENTIFIC COMMITTEE FOR TUNA AND TUNA-LIKE SPECIES IN THE NORTH PACIFIC OCEAN

PLENARY SESSION

12-17 July 2017 Vancouver, British Columbia Canada

TABLE OF CONTENTS

1	INTR	ODUCTION AND OPENING OF THE MEETING	12	
	1.1	Introduction	12	
	1.2	OPENING OF THE MEETING	12	
2	ADOI	TION OF AGENDA	13	
3	DELE 13	GATION REPORTS ON FISHERY MONITORING, DATA COLLECTION AN	D RESEAR	
	3.1	Canada	13	
	3.2	CHINESE-TAIPEI	14	
	3.3	Japan	15	
	3.4	REPUBLIC OF KOREA	16	
	3.5	MEXICO	17	
	3.6	U.S.A.	17	
4	REPO	ORT OF THE CHAIRMAN	18	
5	INTE	RACTIONS WITH REGIONAL ORGANIZATIONS	20	
	5.1	WCPFC	20	
	5.2	PICES	20	
	5.2.1	ISC-PICES Collaboration	20	
	5.2.2	2018 PICES Symposium Co-Sponsor.	21	
6	REPORT OF SPECIES WORKING GROUPS AND REVIEW OF ASSIGNMENTS			
	6.1	Albacore	21	
	6.2	PACIFIC BLUEFIN TUNA	23	
	6.2.1	PBF Stakeholders' Meeting	25	
	6.3	BILLFISH	26	
	6.4	Shark	29	
7	STOC	K STATUS AND CONSERVATION INFORMATION	30	
	7.1	Albacore	30	
	7.2	Blue shark	39	
	7.3	SHORTFIN MAKO SHARK	44	
	7.4	North Pacific Swordfish	44	
	7.5	WESTERN AND CENTRAL NORTH PACIFIC STRIPED MARLIN	46	
	7.6	Pacific Blue Marlin	47	
	7.7	Pacific Bluefin Tuna	48	
8	REVIEW OF STOCK STATUS OF SECONDARY STOCKS IN THE WESTERN PACIFIC52			
	8.1	EPO	52	
	8.2	WCPO	52	
9	REVI	EW OF STATISTICS	53	
	9.1	STATWG REPORT	53	
	9.2	TOTAL CATCH TABLES	55	
10	REPO	ORT OF THE SEMINAR	56	
11	REVI	EW OF MEETING SCHEDULE	57	
	11.1	TIME AND PLACE OF ISC18	57	

	11.2	TIME AND PLACE OF WORKING GROUP INTERCESSIONAL MEETINGS	57
12	ADM	INISTRATIVE MATTERS	59
	12.1	FORMALIZATION OF ISC	59
	12.2	PEER REVIEW OF FUNCTION AND PROCESS FOR STOCK ASSESSMENTS	59
	12.3	Work Group Election Results	60
	12.4	ELECTION OF THE CHAIR AND VICE CHAIR	60
	12.5	OTHER BUSINESS	62
		1 Template for Stock Status and Conservation Advice	
	12.5.	2 Candidate Reference Points	62
13	ADOI	PTION OF REPORT	63
14	CLOS	SE OF MEETING	63
15	CATO	CH TABLES	65

LIST OF TABLES

Table 7-1. Estimates of maximum sustainable yield (MSY), female spawning biomass (SSB) quantities, and fishing intensity (F) based reference point ratios for north Pacific albacore tuna for the base case assessment and important sensitivity analyses. SSB₀ and SSB_{MSY} are the unfished biomass of mature female fish and at MSY, respectively. The Fs in this table are not based on instantaneous fishing mortality. Instead, the Fs are indicators of fishing intensity based on SPR and calculated as 1-SPR so that the Fs reflects changes in fishing mortality. SPR is the equilibrium SSB per recruit that would result from the current year's pattern and intensity of fishing mortality. Current fishing intensity is based on the average fishing intensity during 2012-Table 7-2. Estimates of key management quantities for the North Pacific blue shark SS stock Table 7-3. Projected trajectory of spawning biomass (in metric tons) for alternative harvest Table 15-1. North Pacific albacore catches (in metric tons) by fisheries, 1952-2012. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional.......65 Table 15-2. Pacific bluefin tuna catches (in metric tons) by fisheries, 1952-2012. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional......69 Table 15-3. Annual catch of swordfish (Xiphias gladius) in metric tons for fisheries monitored by ISC for assessments of North Pacific Ocean stocks, 1951-2010. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not Table 15-4. Annual catch of striped marlin (Kajikia audax) in metric tons for fisheries monitored by ISC for assessments of North Pacific Ocean stocks, 1951-2011 "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not Table 15-5. Retained catches (metric tons, whole weight) of ISC Members of blue marlin (Makaira nigricans) by fishery in the North Pacific Ocean, north of the equator. "0": Fishing

LIST OF FIGURES

Figure 6-1. An example of flowchart of development and implementation of an emergency rule Figure 7-1. Maximum likelihood estimates of (A) total age-1+ biomass (open circles) (B), female spawning biomass (SSB) (solid blue line), and (C) age-0 recruitment (open circles) of north Pacific albacore tuna (Thunnus alalunga). Dashed lines (B) and vertical bars (C) indicate 95% confidence intervals of the female SSB and recruitment estimates respectively. Red line indicates the 20%SSB_{current, F=0} limit reference point, which is based on dynamic SSB₀. Closed black circle and error bars in (B) are the maximum likelihood estimate and 95% confidence intervals of unfished female spawning biomass, SSB₀. Since the assessment model represents time on a quarterly basis, there are four estimates of total biomass (A) for each year, but only one annual estimate of female SSB (B) and recruitment (C). Figure 7-2 Estimated sex-specific instantaneous fishing mortality-at-age (F-at-age) for the 2017 Figure 7-3. Historical catch-at-age of north Pacific albacore (*Thunnus alalunga*) estimated by the Figure 7-4. Fishery impact analysis on north Pacific albacore (*Thunnus alalunga*) showing female spawning biomass (SSB) (red) estimated by the 2017 base case model as a percentage of dynamic unfished female SSB (SSB₀). Colored areas show the relative proportion of fishing impact attributed to longline (USA, Japan, Chinese-Taipei, Korea and others) (green) and surface (USA, Canada, and Japan) (blue) fisheries (primarily troll and pole-and-line gear, but including all other gears except longline). Figure 7-5. (A) Kobe plot showing the status of the north Pacific albacore (*Thunnus alalunga*) stock relative to the 20% SSB current, F=0 biomass-based limit reference point (WCPFC), and equivalent fishing intensity (F20%; calculated as 1-SPR20%) over the base case modeling period (1993-2015). Note that no F-based reference point is established for north Pacific albacore by either the IATTC or WCPFC. F20% was chosen for this plot as a convenient analogue to the limit reference point but it should not be inferred that it represents advice on the choice of an Fbased reference point on this stock. Black triangle indicates the start year (1993) and black circle with 95% confidence intervals indicates the terminal year (2015). (B) Kobe plot showing stock status and 95% confidence intervals in the terminal year (2015) of the base case model (black; closed circle) and important sensitivity runs with M = 0.3 y-1 for both sexes (pen square), and CV = 0.06 for L_{inf} in the growth model (white; open triangle). Fs in this figure are not based on instantaneous fishing mortality. Instead, the Fs are indicators of fishing intensity based on SPR

and calculated as 1-SPR so that the Fs reflects changes in fishing mortality. SPR is the
equilibrium SSB per recruit that would result from the current year's pattern and intensity of
fishing mortality
Figure 7-6. (A) Historical and future trajectory of north Pacific albacore (<i>Thunnus alalunga</i>)
female spawning biomass (SSB) under a constant fishing intensity (F ₂₀₁₂₋₂₀₁₄) harvest scenario.
Future recruitment was based on the expected recruitment variability and autocorrelation. Black
line and blue area indicates maximum likelihood estimates and 95% confidence intervals (CI),
respectively, of historical female SSB, which includes parameter uncertainty. Red line and red
area indicates mean value and 95% CI of projected female SSB, which only includes future
recruitment variability and SSB uncertainty in the terminal year. (B) Expected annual catch
under a constant fishing intensity ($F_{2012-2014}$) harvest scenario (2016-2025). The red line is the
current average catch (2010-2014 = 82,432 t)
Figure 7-7. (A) Historical and future trajectory of north Pacific albacore (<i>Thunnus alalunga</i>)
female spawning biomass (SSB) under a constant catch (average 2010-2014 = 82,432 t) harvest
scenario. Future recruitment was based on the expected recruitment variability and
autocorrelation. Dashed line indictaes the average limit reference point threshold for 2012-2014.
Black line and blue area indicates maximum likelihood estimates and 95% confidence intervals
(CI), respectively, of historical female SSB, which includes parameter uncertainty. Red line and
red area indicates mean value and 95% CI of projected female SSB, which only includes future
recruitment variability and SSB uncertainty in the terminal year. (B) Projected fishing intensity
relative to the current fishing inetnsity (2012-2014) (red line) under a constant catch scenario
(average 2010-2014)
Figure 7-8. Results of the SS stock assessment reference case model for the North Pacific BSH
assessment: (A) estimated age-0 recruits (circles) and 95% confidence intervals (vertical bars);
(B) estimated female spawning biomass and 95% confidence intervals (blue shaded area); (C)
estimated fishing mortality (sum of F's across all fishing fleets). Red solid lines indicate the
estimates of SB _{MSY} and F _{MSY} in (B) and (C), respectively41
Figure 7-9. Kobe plots of the trends in estimates of relative fishing mortality and biomass of
North Pacific blue shark between 1971 and 2015 for the reference case of (A) the SS stock
assessment model, and (B) the BSSPM stock assessment model
Figure 7-10. Comparison of future projected North Pacific blue shark spawning biomass under
different F harvest policies (status quo, +20%, -20%, and F_{MSY}) using the SS reference case
model. Status quo fishing mortality was based on the average from 2012-201443

LIST OF ANNEXES

Annex 1	List of Participants
Annex 2	ISC Meeting Provisional Agenda
Annex 3	List of Meeting Documents
Annex 4	Report of the Albacore Working Group Workshop (November 2016)
Annex 5	Report of the Shark Working Group Intersessional Workshop (November 2016)
Annex 6	Report of the Statistics Working Group Intersessional Workshop (January 2017)
Annex 7	Report of the Pacific Bluefin Tuna Working Group Intersessional Workshop (February-March 2017)
Annex 8	Report of the Shark Working Group Workshop (March 2017)
Annex 9	Report of the Albacore Working Group Workshop (April 2017)
Annex 10	Report of the Billfish Working Group Workshop (June 2017)
Annex 11	Draft Summary of Additional Projections: International PBF Stakeholders Meeting (April 2017)
Annex 12	Stock Assessment of Albacore Tuna in the North Pacific Ocean in 2017
Annex 13	Stock Assessment and Future Projections of Blue Shark in the North Pacific Ocean Through 2015
Annex 14	Summary Report of the Pacific Bluefin Tuna International Stakeholders Meeting (April 2017)
Annex 15	Report of the Seminar: HMS Tagging and Lessons Learned
Annex 16	Report of the Statistics Working Group Workshop (July 2017)

ACRONYMS AND ABBERVIATIONS

Names and FAO Codes of ISC Species of Interest in the North Pacific Ocean

FAO Code	Common English Name TUNAS	Scientific Name
ALB	Albacore	Thunnus alalunga
BET	Bigeye tuna	Thunnus obesus
PBF	Pacific bluefin tuna	Thunnus orientalis
SKJ	Skipjack tuna	Katsuwonus pelamis
YFT	Yellowfin tuna	Thunnus albacares
	BILLFISHES	
BIL	Other billfish	Family <i>Istiophoridae</i>
BLM	Black marlin	Makaira indica
BUM	Blue marlin	Makaira nigricans
MLS	Striped marlin	Kajikia audax
SFA	Sailfish	Istiophorus platypterus
SSP	Shortbill spearfish	Tetrapturus angustirostris
SWO	Swordfish	Xiphias gladius
	SHARKS	
ALV	Common thresher shark	Alopias vulpinus
BSH	Blue shark	Prionace glauca
BTH	Bigeye thresher shark	Alopias superciliosus
FAL	Silky shark	Carcharhinus falciformis
LMA	Longfin mako	Isurus paucus
LMD	Salmon shark	Lamna ditropis
OCS	Oceanic whitetip shark	Carcharhinus longimanus
PSK	Crocodile shark	Pseudocarcharias kamonharai
PTH	Pelagic thresher shark	Alopias pelagicus
SMA	Shortfin mako shark	Isurus oxyrinchus
SPN	Hammerhead spp.	Sphyrna spp.

ISC Working Groups

Acronym	Name	Chair
ALBWG	Albacore Working Group	John Holmes (Canada)
BILLWG	Billfish Working Group	Jon Brodziak (U.S.A.)
PBFWG	Pacific Bluefin Working Group	Hideki Nakano (Japan)
SHARKWG	Shark Working Group	Suzanne Kohin (U.S.A.)
STATWG	Statistics Working Group	Ren-Fen Wu (Chinese
		Taipei)

Other Abbreviations and Acronyms Used in the Report

CDS Catch documentation scheme
CIE Center for Independent Experts
CKMR Close-kin mark-recapture

CMM Conservation and Management Measure

CPFV Charter passenger fishing vessel

CPUE Catch-per-unit-of-effort

CSIRO Commonwealth Scientific and Industrial Research Organization

DWLL Distant-water longline
DWPS Distant-water purse seine
EEZ Exclusive economic zone
EPO Eastern Pacific Ocean
F Fishing mortality rate
FAD Fish aggregation device

FAO Fisheries and Agriculture Organization of the United Nations

FL Fork length

HCRHarvest control ruleHMSHighly migratory species H_{MSY} Harvest rate at MSY

IATTC Inter-American Tropical Tuna Commission

ISC International Scientific Committee for Tuna and Tuna-Like Species in the

North Pacific Ocean

ISSF International Seafood Sustainability Foundation LFSR Low fecundity spawner recruitment relationship

LTLL Large-scale tuna longline LRP Limit reference point

MSE Management strategy evaluation MSY Maximum sustainable yield NC Northern Committee (WCPFC)

NRIFSF National Research Institute of Far Seas Fisheries (Japan)
OFDC Overseas Fisheries Development Council (Chinese Taipei)

PICES North Pacific Marine Science Organization
PIFSC Pacific Islands Fisheries Science Center (U.S.A.)

SAC Scientific Advisory Committee (IATTC)

SC Scientific Committee (WCPFC)

SG-SCISC Study Group on Scientific Cooperation of ISC and PICES

SPC-OFP Oceanic Fisheries Programme, Secretariat of the Pacific Community

SPR Spawning potential ratio, spawner per recruit

SSB Spawning stock biomass

SSB_{F=0} Spawning stock biomass at a hypothetical unfished level

SSB_{CURRENT} Current spawning stock biomass

SSB_{MSY} Spawning stock biomass at maximum sustainable yield

STLL Small-scale tuna longline

t, mt Metric tons, tonnes

WCNPO Western Central and North Pacific Ocean

WCPFC Western and Central Pacific Fisheries Commission

WPO Western Pacific Ocean

WWF World Wildlife Fund for Nature - Japan

GRT Gross registered tons

REPORT OF THE SEVENTEENTH MEETING OF THE INTERNATIONAL SCIENTIFIC COMMITTEE FOR TUNA AND TUNA-LIKE SPECIES IN THE NORTH PACIFIC OCEAN

PLENARY SESSION

12-17 July 2017 Vancouver, British Columbia Canada

Highlights of the ISC17 Plenary Meeting

The 17th ISC Plenary, held in Vancouver, British Columbia Canada from 12-17 July 2017 was attended by Members from Canada, Chinese Taipei, Japan, Korea, Mexico and the United States, as well as the Western and Central Pacific Fisheries Management Commission and the North Pacific Marine Science Organization. The Plenary reviewed results, conclusions, new data, and updated analyses of the Billfish, Albacore, Shark and Pacific Bluefin Tuna working groups. The Plenary endorsed the findings that North Pacific albacore tuna (NPALB) is not experiencing overfishing and is not overfished, and considers the NPALB stock assessment to be best available scientific information. The Plenary also endorsed the findings that North Pacific blue shark (BSH) is not experiencing overfishing and is not overfished, and considers the North Pacific BSH stock assessment to be best available scientific information. It re-iterated stock status and conservation information proffered at ISC16 for North Pacific swordfish, Pacific blue marlin, North Pacific striped marlin, and North Pacific shortfin make shark. It reiterated stock status condition proffered at ISC16 for Pacific bluefin tuna, and provided revised conservation information based on extensive harvest scenario testing requested by WCPFC and IATTC. A science seminar on HMS tagging and lessons learned was held and the Plenary agreed to assess the feasibility of developing an multinational North Pacific bluefin tuna tagging program. Observers from Pew Charitable Trusts, World Wildlife Fund for Nature – Japan, Monterey Bay Aquarium, Wild Oceans, Tohoku University, Waseda University, and American Fisherman's Research Foundation/American Albacore Fishing Association attended. The ISC workplan for 2017-18 includes completing Pacific Bluefin tuna, North Pacific shortfin make shark, and North Pacific swordfish assessments, improving catch and CPUE time series and advancing biological information for shark species, conducing the third MSE workshop, developing protocols for an ISC stock assessment review process, and enhancing database and website management. John Holmes (Canada) was elected as the new ISC Chair and Eric Chang (Chinese Taipei) as Vice Chair. The next Plenary will be held in the Korea in July 2018.

1 INTRODUCTION AND OPENING OF THE MEETING

1.1 Introduction

The ISC was established in 1995 through an intergovernmental agreement between Japan and the United States (U.S.A.). Since its establishment and first meeting in 1996, the ISC has undergone a number of changes to its charter and name (from the Interim Scientific Committee to the International Scientific Committee) and has adopted a number of guidelines for its operations. The two main goals of the ISC are (1) to enhance scientific research and cooperation for conservation and rational utilization of the species of tuna and tuna-like fishes that inhabit the North Pacific Ocean during a part or all of their life cycle; and (2) to establish the scientific groundwork for the conservation and rational utilization of these species in this region. The ISC is made up of voting Members from coastal states and fishing entities of the region as well as coastal states and fishing entities with vessels fishing for highly migratory species in the region, and non-voting Members from relevant intergovernmental fishery and marine science organizations, recognized by all voting Members.

The ISC provides scientific advice on the stocks and fisheries of tuna and tuna-like species in the North North Pacific Ocean to the Member governments and regional fisheries management organizations. Fishery organizations. Fishery data tabulated by ISC Members and peer-reviewed by the species and statistics statistics Working Groups (WGs) form the basis for research conducted by the ISC. Although some data for some data for the most recent years are incomplete and provisional, the total catch of highly migratory migratory species (HMS) by ISC Members estimated from available information is in excess of 500,000 500,000 metric tons (t) annually and dominated by the tropical tuna species. In 2016, the catch of priority priority species monitored by ISC Member countries was 53,543 t of North Pacific albacore tuna (NPALB, (NPALB, Thunnus alalunga), 13,167 t of Pacific bluefin tuna (PBF, T. orientalis), 8,867 t of North Pacific North Pacific swordfish (SWO, Xiphias gladius), 2,034 t of North Pacific striped marlin (MLS, Kajikia audax), Kajikia audax), 7,157 t of Pacific blue marlin (BUM, Makaira nigicans), 346 t of shortfin mako shark (SMA, shark (SMA, Isurus oxyrinchus) and 22,642 t of North Pacific blue shark (BSH, Prionace glauca). The total glauca). The total estimated catch of these seven species is 107,746 t, or approximately 91% of the 2015 total the 2015 total estimated catch of 118,994 t. Annual catches of priority stocks throughout their ranges are ranges are shown in Table 15-1 through

_

¹ FAO three-letter species codes are used throughout this report interchangeably with common names.

Table 15-7.

1.2 Opening of the Meeting

The Seventeenth Plenary session of the ISC (ISC17) was convened in Vancouver, British Columbia, Canada, at 0900 on 12 July 2017 by the ISC Chairman, G. DiNardo. A roll call confirmed the presence of delegates from Canada, Chinese Taipei, Japan, Republic of Korea, Mexico, and U.S.A. (ISC/17/ANNEX/1). A representative from the Western and Central Pacific Fisheries Commission (WCPFC) was also present. Pew Charitable Trusts, World Wildlife Fund for Nature - Japan (WWF), Monterey Bay Aquarium, Wild Oceans, Tohoku University, Waseda University, and American Fishermen's Research Foundation/American Albacore Fishing Association were present as observers.

ISC Member China, as well as the non-voting Members the Secretariat of the Pacific Community (SPC), Food and Agriculture Organization of the United Nations (FAO), and the Inter-American Tropical Tuna Commission (IATTC), while extended an invitation, did not attend the Plenary.

J. Holmes introduced Dr. Carmel Lowe, Regional Director Science, Pacific Region, Fisheries and Oceans Canada, who gave the welcome address for the meeting. Dr. Lowe welcomed the Members and emphasized the importance of albacore to Canada and Canada's history of fishing on the stock in the EPO, which has sustained industry and coastal communities. In addition to albacore, ISC provides sound science to manage other North Pacific tuna and tuna-like species and Canada is committed to science-based decision making. She noted that 2017 is a special year for Canada, because it marks the 150th anniversary of its sovereign independence. She encouraged meeting participants to enjoy the many cultural events associated with anniversary. She concluded by emphasizing the continued importance of the ISC's work, especially on depleted stocks such as Pacific bluefin tuna. Canada supports the ISC's ongoing work, as evidenced by its hosting of this meeting and the upcoming North Pacific albacore management strategy evaluation workshop.

2 ADOPTION OF AGENDA

The proposed agenda for the session (ISC/17/ANNEX/02) was considered and adopted. It was noted that observers would be given the opportunity at the end of each day to offer comments and seek clarification on topics discussed. C. Dahl was assigned lead rapporteur duties. A list of meeting documents is contained in ISC/17/ANNEX/03.

3 DELEGATION REPORTS ON FISHERY MONITORING, DATA COLLECTION AND RESEARCH

3.1 Canada

J. Holmes presented a summary of Category I, II, and III data from Canadian fisheries for highly migratory species in 2016 (ISC/17/PLENARY/04). The Canadian fleet of 152 vessels targets juvenile North Pacific ALB and operates primarily in the coastal waters of Canada and the United States, with little effort or catch outside of these areas in 2016. Preliminary estimates of effort and catch for 2016 are 5,359 vessel-days (v-d) and 2,842 t, respectively. While effort

increased about 2% relative to 2015, catch declined 35% relative to 2015. Catch and effort were split primarily between Canadian waters (55% of the catch and 64% of the effort) and U.S. waters (44% of the catch and 35% of the effort). The Canadian fleet experienced the highest catch rates in June, after which catch rates were below average for the remainder of the fishing season. This early peak in catch rates is unusual for this fishery, but the below-average catch rates are consistent with lower catches – North Pacific albacore (NPALB) were difficult to find in coastal waters in 2016. The size composition data sampled from the catch (N=14,189) in 2016 exhibit a primary mode at 68-71 cm FL, with a smaller secondary mode between 79 and 83 cm FL. These modes correspond to two-year old and three-year old fish, respectively.

Discussion

There does not appear to be an explanation for the shift in the timing of the catch (more catch early in the season) in 2016 compared to previous years. Both potential environmental effects and changes in the spatial distribution of fishing effort have been ruled out. The apparent decline in catch-per-unit-of-effort (CPUE) (suggested by the simultaneous decline in catch and increase in fishery participation) cannot be explained by new entrants with limited fishing experience.

It was noted that for the past six or more years the fishing season established under the United States-Canada Albacore Treaty now ends earlier (September 15) than under past fishing access regimes. Canadian vessels fish for a few weeks in Canadian waters after access to U.S. waters closes, and fish generally migrate out of the Canadian EEZ by mid-October.

The number of undersize, released fish (colloquially referred to as "peanuts") declined substantially in 2016 from a spike in 2015. The 2015 catch of undersize fish suggested the potential for a spike in recruitment in later years, which apparently has not materialized yet. It was noted that there is a spatial component to the size distribution of catch; historically when Canadian vessels fish farther south (e.g., off of California) they have encountered more undersize fish. However, in 2015 small fish were also caught and released in the Canadian EEZ, consistent with a northern shift in the stock.

3.2 Chinese-Taipei

W. Wang presented the Chinese-Taipei National Report (ISC/17/PLENARY/05). There are two principal tuna fisheries of Chinese-Taipei operating in the North Pacific Ocean, the tuna longline fishery and the distant water purse seine fishery. Other offshore and coastal fisheries include the harpoon, setnet and gillnet fisheries, and account for a small proportion of overall tuna and tuna-like species catch. The catches of longline and purse seine fisheries account for 99% of the total tuna and tuna-like species catches in the North Pacific Ocean by Chinese-Taipei. Longline fisheries are composed of the large-scale tuna longline (LTLL, vessels larger than 100 gross registered tons, GRT) and small-scale tuna longline (STLL, vessels less than 100 GRT) fleets. The total catch of tunas and billfish (including SWO, MLS, BUM, BLA, and SFA) for the combined longline fishery (LTLL and STLL) in the North Pacific Ocean was 23,771 t for 2016 with 79 active LTLL vessels and 1,303 STLL vessels. For the purse seine fishery, the total catch was 185,692 t caught by 34 authorized vessels in the Pacific Ocean in 2016.

For the LTLL fishery, the sources of Category I data (annual catch by gear) include weekly catch reports and commercial data from individual fishing vessels. Category II and III data are all compiled from logbook data. Fishermen are required to measure the length of the first 30 fish caught in each longline set. For the STLL fishery, the sources of Category I data include landings and auction records of local fish markets, reports of market sales, and monthly catch reports from individual fishing vessels. For the purse seine fishery, Category I, II, and III data are obtained from logbooks.

In March 2010, a catch documentation scheme (CDS) was established and implemented for vessels fishing for PBF. In addition to prior authorization, fishing vessels targeting PBF are required to notify the relevant authority and attach a tag to every PBF caught by them. Moreover, port samplers are dispatched to measure length and weight of each PBF. In addition, Chinese-Taipei collected 1,408 PBF tissue samples in 2016 under the close-kin mark recapture project.

An observer program for the LTLL fleet was implemented in the Pacific Ocean in 2002. The program has gradually expanded in recent years and hence the number of observers has increased. The program was further expanded to the STLL fleet in 2012. A total of thirty observers were deployed on longline vessels in 2016, including 10 observers on LTLL vessels and 20 observers on STLL vessels.

Discussion

The change in overall fishing effort in longline fisheries was discussed. While the number of registered STLL vessels has remained stable, the number of active vessels has declined in recent years due to various economic and operational factors, resulting in a decreased catch.

3.3 Japan

H. Okamoto presented the Japan National Report (ISC/17/PLENARY/06). Japanese tuna fisheries consist of the three major fisheries (i.e., longline, purse seine, pole-and-line) and other miscellaneous fisheries like troll, driftnet, and setnet fisheries. The number of vessels has shown a decreasing trend in the longline and pole-and-line fisheries, and has been stable in the purse seine fishery in this decade. The total catch of tunas (excluding SKJ) caught by Japanese fisheries in the North Pacific Ocean was 111,115 t in 2015 and 94,884 t in 2016. The total catch of tunas (including SKJ) caught by Japanese fisheries in the North Pacific Ocean was 300,652 metric t in 2015 and 247,425 t in 2016. Preliminary catch of PBF and ALB in 2016 was 8,302 t and 35,582 t, respectively. ALB caught by pole-and-line ranged in size from 60 to 90 cm FL in 2016, and fewer individuals smaller than 60 cm were caught compared to 2015. The total catch of SWO and MLS in 2016 was 4,885 t and 1,320 t, respectively. In addition to this information, three research topics initiated in the Pacific Ocean in 2016 were noted: a PBF egg/larvae research cruise, recruitment monitoring on juvenile PBF, and a tagging research cruise on shark.

Discussion

It was pointed out that PBF catch in purse seine fisheries appears to have increased since 2013, possibly exceeding quotas established pursuant to the WCPFC Conservation and Management

Measure (CMM). It was clarified that in Japan some quotas set at the prefectural level have been exceeded and the total catch with respect to the catch limit will be reported to WCPFC.

The catch of small PBF can be quite variable from year to year due to the availability of fish in the target year classes (age 0 and 1) and Japan's recruitment monitoring program has documented this variability in recruitment. The next stock assessment for PBF will undoubtedly provide more information on recruitment and fishing mortality.

Catch of ALB declined in 2016; a change in fishing area has been ruled out as a reason but it was noted that the northward movement of pole-and-line fishing grounds occurred later than in previous years.

3.4 Republic of Korea

D. Kim presented the Korea National Report (ISC/17/PLENARY/07). The Korean distant water fishery uses two types of fishing gears, purse seine and longline, that engage in fishing for tuna and tuna-like species in the North Pacific. The total number of active longline vessels showed a stable pattern, ranging from 122 vessels in 2010 to 125 in 2013; however, the number decreased to 96 in 2016. The total number of active purse seine vessels was constant during 2010-2014 but decreased slightly in 2015 and 2016 to 25 vessels. The total catch of tuna and tuna-like species caught by Korean distant water fisheries in the North Pacific Ocean was 84,586 t in 2016. The total longline fishery catch was 10,593 t, which is 53.4 % of the historical highest catch in 2004. Catch by the purse seine fishery was 73,993 t, which corresponds to 73.5% of the historical highest catch in 2003. As for the catch composition of the longline fishery, dominant species were BET, which is over 65% of the total catch, and YFT and BUM comprising 15% and 10%, respectively. In the purse seine fishery, SKJ, YFT, and BET comprised 85%, 13%, and 1%, of the catch respectively. The offshore large purse seine fishery caught 1,029 t of PBF in the Korean EEZ in 2016. This catch was mainly distributed in the South Sea around Jeju Island throughout the year with the highest catches in March. Large PBF (≥ 30kg) were 46% of the total PBF catch and most of the large PBF were caught during only two days in March. For close-kin analysis 1,045 PBF tissue samples were collected in 2016 and 348 samples were collected in 2017.

Discussion

Korea provided additional information about its fishery monitoring program. It has an electronic monitoring system with reporting in real time through satellite transmission. Fishermen have up to 72 hours to correct the reported information on catch and effort, if necessary. Potential inaccuracies in logbook reporting were questioned due to fishermen being less able to identify non-target bycatch species. Educating fishermen on species identification is warranted. However, logbook formats do allow reporting of all data required by RFMOs including length frequency data. It was noted that coastal fishery participants have remarked on the utility of the data they report through the electronic monitoring program. This may provide an incentive for accurate self-reporting. The coastal fishery does not have at-sea fishery observers but shoreside observers are used to document landings.

It was noted that the purse seine fishery exhibited a marked eastward shift in the distribution of fishing effort in 2015. This effort shift is likely attributable to the strong El Niño occurring in that year.

The increase in PBF landings by weight in 2016 may be attributable in part to an increase in the average weight of the fish caught. More fish \geq 30 kg were caught in 2016 compared to 2015. Catch of larger fish may have contributed to Korea exceeding its PBF quota in 2016 but Korea will adjust quotas downward in future years in compensation so that catch over the five-year period 2017-2021 is consistent with the WCPFC CMM.

3.5 Mexico

M. Dreyfus presented the Mexico National Report (ISC/17/PLENARY/08). Billfishes except SWO have been reserved for the sport fishery since 1980s. Most of the sport fishing activities are located in areas near the southern tip of the Baja California Peninsula. It is a catch and release activity where most of the catch is dolphin fish and about 20% are billfishes, dominated by MLS.

Pacific shark fisheries have traditionally reported catch in two groups (big and small shark individuals). A logbook and requirement, under which fishermen provide species composition of the catch, was introduced in 2007. As a conservation measure, in 2012 Mexico implemented a three-month closure covering the period of reproduction of many of the species found in Mexican waters. Since 2007 there also has been an observer program, which covers both longline and artisanal fleets at different coverage rates. Mexican scientists have been cooperating with the ISC SHARKWG; providing time series of catch estimates for SMA and BSH, as well as size composition of the catch. They are also collaborating with the U.S.A. and Japan in an SMA age and growth estimation study.

In relation to tunas, Mexico has a fleet that primarily targets YFT and complements its catch with SKJ. All vessels above 400 cubic meters capacity have a scientific observer on board from the IATTC or the national program. Some of those vessels also participate in the PBF fishery each year. The catch is transferred to net pen grow out farms in northwest Baja California Peninsula. A quota for the PBF commercial fishery has been in place since 2012. In 2016 2,700 t was retained after the release of 192 t to comply with the voluntary measure Mexico adopted for that year. Since 2013 Mexico also has been providing size composition of the PBF catch for PBF assessments, and is currently processing 2015-2017 data for that same purpose. Related to the PBF close-kin analysis project, tissue sampling began this year using animals captured last year. This activity is expected to continue for several years with the help of the farming industry.

Discussion

Mexico noted that when collecting samples for the PBF close-kin analysis project lengths of individual animals are not always collected. However, in collaboration with the PBF net pen operations the size distribution of each harvest is accurately documented. A general indication of the size of fish sampled for close-kin analysis can be derived from this information.

Variability in the catch of MLS and SFA was discussed. Mexico asserted that recent inter-annual variability is not outside the historical range.

3.6 U.S.A.

M. Seki presented the United States National Report to the Plenary (ISC/17/PLENARY/09). The Pacific Ocean produces about 71% of the global tuna catch. And in the Pacific Ocean, about 79% of the tuna catch is from the Western Pacific. Purse seine gear accounts for 66.8% of the tuna catch, followed by longline (9.6%) and pole-and-line (8.1%). In 2016, there were 571 vessels in the U.S.A. albacore troll fishery in the North Pacific, the fewest since 2008 when there were 525 vessels. The 2016 albacore troll catch was 10,686 t. For the U.S.A. purse-seine fishery, the number of vessels fishing in the North Pacific Ocean decreased to 42 in 2016, compared to 46 in 2015. Total U.S.A. purse-seine catch in the North Pacific Ocean was 61,300 t in 2016, an increase from 53,822 t in 2015. The catch consisted of 86% skipjack and 11% yellowfin tuna. The 2013-2016 catches are considered preliminary since the species composition of juvenile YFT and BET have not been accounted for and has not been adjusted. U.S.A. longline fishing activity for BET and SWO has remained stable since 2008. In 2016, 141 longline vessels landed a total of 12,088 t. Of this, 5,848 t were BET, and 1049 t were SWO. Research updates were provided on the impact of oceanic fronts on ALB foraging efficiency, possible signs of increased PBF in the Southern California Bight in spring 2017, PBF foraging ecology, PBF close-kin sample collection and a workshop planned for 2017, BUM age estimation, and shark post-release survival in longline fisheries. With respect to bycatch, research on the effect of hook attachment rings on catch and bycatch, and passive acoustic studies on false killer whale encounters and depredation with longline gear was also presented. Additionally, studies concerning the impact of climate change on North Pacific carrying capacity, development of a recruitment index for BET to forecast catch, ongoing efforts to improve modeling for stock assessments, and economic performance of pelagic longline fishing in Hawaii were highlighted.

Discussion

It was explained that the age validation study of a very large BUM specimen using bomb radiocarbon dating noted in the U.S.A. report was a one-time activity and enabled by the availability of the specimen. Future work is dependent on acquiring specimens.

In response to a question, the Chair detailed progress on the ISC PBF close-kin analysis project. ISC Members agreed to collect tissue samples starting in 2015 but questions have arisen about processing the samples. The U.S.A. is planning to hold a workshop for ISC Members to review and standardize processing methods. The Chair noted that Japan has already made considerable progress in sample processing and genetic analysis; their expertise should be a starting point for determining standard processing techniques at the workshop. He also noted that the cost of sample processing and genetic analysis varies widely among countries; in general it is much cheaper in the U.S.A., so it may make sense for countries to send their samples to firms in the U.S.A. for processing. Both Japan and the U.S.A. will work collaboratively to develop the necessary simulation models to test procedures for estimating the stock abundance from close-kin analysis results.

4 REPORT OF THE CHAIRMAN

G. DiNardo, ISC Chair presented the following report to the Plenary.

The ISC had another busy year since the ISC Plenary met in Sapporo, Japan in July 2016. The year was spent completing a stock assessment update for blue shark, a benchmark assessment for North Pacific albacore tuna, advancing research collaborations with PICES, and collection of biological samples to facilitate close-kin research, as well as preparing for assessments on Pacific bluefin tuna and shortfin make shark in 2018, evaluating requested Pacific Bluefin tuna harvest strategies, and developing a process to formalize the structure/existence of the ISC. While numerous accomplishments and successes advanced the scientific integrity of ISC, we cannot afford to waiver from our scientific mission. The failure of ISC to provide best available scientific information has far-reaching implications. While international fisheries management can be political at times, politics has no place in science. As an independent scientific organization ISC must stay the course, which includes appointing leaders with proper scientific credentials.

Progress was made by improving best practices and scientific reporting procedures, compiling a catalogue and inventory of the ISC database, and advancing development of the website and data enterprise system. Seven intersessional workshops and two webinars were held to facilitate collaboration among Member scientists in implementing ISC work plans and coordinating research on the stocks. In addition, the ISC convened a Pacific Bluefin Tuna International Stakeholders Meeting in Tokyo Japan, initiated planning for the third international workshop on Management Strategy Evaluations tentatively scheduled for October 2017 in Vancouver, BC, Canada, and in November 2016 co-convened a workshop at the 2016 Annual PICES Meeting in San Diego, CA as part of the joint ISC-PICES Working Group-34 to assess the impacts of climate variability on highly migratory species. We continue to address recommendations stemming from the 2013 peer review of the ISC function. Jon Brodziak (U.S.A.) and Ren-Fen Wu (Chinese-Taipei) were reelected as Chairs of the ISC Billfish and Statistics Working Groups, respectively. Hidetada Kiyofuji (Japan) was elected as Chair of the ISC Albacore Working Group, Mikihiko Kai (Japan) as Vice Chair of the Shark Working Group and Hirotaka Ijima (Japan) as Vice Chair of the Billfish Working Group.

Managing ISC activities continued to be a challenge during the past year. As before, the challenge is an inherent consequence of the ISC framework adopted by the Members. That is, ISC relies on in-kind contributions from its Members rather than monetary contributions to support a "Secretariat" to oversee day-to-day operations of the organization. Given this framework, the Office of the Chairman takes on the role of a Secretariat, but not a full-service one at that, owing to uncertain support from the Chairman's funding source. Likewise, the working groups depend on in-kind contributions from Members who elect to participate in specific working groups. This support is uneven among the Members, and can delay progress of a working group in completing assignments. To date, the support for administration of ISC activities has been provided by the U.S. for day-to-day operations of the Office of the Chairman, and by Japan for operating the ISC website and database. Member countries with scientists serving as chairpersons of the working groups have contributed to supporting administrative services of the working groups. All of the support is appreciated and acknowledged. Efforts to formalize the ISC through a Memorandum of Understanding (MOU) are moving forward and should provide the necessary framework to address many of these concerns including support for a fulltime secretariat.

Over the past seven years, the scientific stature, commitment, and competency of ISC has increased significantly, owing to the unwavering dedication and integrity of its scientists. At the same time, the breadth/scope of the organization's research, scientific partnerships, and visibility have also expanded and flourished due to the persistence of its scientists. For that I am deeply grateful.

As this is my last year as ISC Chair, I close this report by thanking all my colleagues who have worked on ISC tasks and who have provided the support to ISC and the Office of the Chair in advancing the objectives and purpose of the organization. The service of Chi-lu Sun, Vice-Chairman, for support and insightful advice is acknowledged, as well as the services of Freddie Logan, Tarah Sullivan, and Stephanie Flores. A special thanks and appreciation is owed to the Chairs of the working groups, namely Ren- Fen Wu, Jon Brodziak, John Holmes, Hideki Nakano, and Suzanne Kohin, who provided unselfish leadership in guiding the work of the Working Groups. In addition, the leadership role of Hideki Nakano with respect to the Data Administrator, Izumi Yamasaki, and Webmaster, Kirara Nishikawa, is appreciated. Finally, I acknowledge the professional assistance and dedicated service of Sarah Shoffler to the ISC in completing tasks assigned to the Chairman. In that capacity, she served as point of contact for the Office of the Chairman, led in writing, editing and assembling technical information required for agenda items of meetings and for responding to inquiries, and served as advisor to me on aspects of ISC. Thanks to all of you for contributing to another successful year for ISC and for the support and service provided.

5 INTERACTIONS WITH REGIONAL ORGANIZATIONS

5.1 WCPFC

A. Beeching presented an SC12 summary of the two ISC stock assessments, PBF and BUM. SC12 showed concern regarding the poor PBF stock status with spawning stock biomass at 2.6% of the estimated unfished spawning stock biomass ($SB_{F=0}$) and low recruitment. He also detailed the issues around the need for an extraordinary meeting of the NC to satisfy some of the PBF issues raised during WCPFC13. The outputs of that meeting were accepted, and they agreed to adopt CMM 2016-04 to establish a multi-annual rebuilding plan for PBF. Finally, the ISC Plenary was briefed on the operations of the WCPFC tissue bank.

Discussion

The Chair noted that in 2015 he sent a letter to WCPFC requesting that all requests of the ISC be put in writing and submitted to the ISC formally by letter. Requests emanating from WCPFC13 were not communicated to ISC in this way. The Chair will remind WCPFC of this process.

5.2 PICES

5.2.1 ISC-PICES Collaboration

G. DiNardo reported on the activities of the joint ISC-PICES Working Group (WG-34) aimed at assessing the impacts of climate variability on highly migratory species that is being facilitated under the auspices of PICES (ISC/17/PLENARY/10). Understanding the impacts of climate

variability on pelagic fish dynamics and spatial structure, and incorporating these processes into stock assessments to support effective fisheries management decision-making, represents the next generation of stock assessment models. The joint collaboration spans 2015-2018, with an opportunity to extend the joint collaboration for an additional year.

It was reported that WG-34 convened its first workshop (W-4) and business meeting at the 2016 PICES Annual Meeting in San Diego, CA, U.S.A. A second business meeting was convened at ISC17 to review progress and membership, as well as potential venues for an international workshop to present research findings and lessons learned stemming from WG-34. A second workshop (W-3) and third possible business meeting will be convened at the 2017 PICES Annual Meeting in Vladivostok, Russia.

Discussion

None.

5.2.2 2018 PICES Symposium Co-Sponsor

G. DiNardo introduced the invitation from PICES for ISC to co-sponsor an international symposium on "Understanding changes in transitional areas in the Pacific" (ISC/17/PLENARY/10). Given the importance of transitional areas (TA) to highly migratory species and fisheries that exploit them, and the importance of this area to the mission of ISC, inviting ISC to participate as a co-sponsor was logical.

The goal of the 2018 symposium is to update and expand our understanding of Pacific TAs. The symposium will focus on questions surrounding natural and anthropogenic climate variability and change in Pacific TAs, and discuss evidence of climate-driven changes in the position and physical structure of TAs and their biological communities. The symposium will also address socioeconomic questions related to the challenges of managing the highly migratory and transboundary resources in TAs. The symposium organizers will strive to publish the proceedings in a peer-reviewed journal.

Discussion

It was noted that the symposium on TAs covers more types of oceanographic features than just frontal zones (such as meso-scale eddies), and many high seas fisheries operate in these areas. Scientists from U.S.A., Japan, and Mexico are assisting in organizing the symposium.

The ISC Chair polled Members as to whether ISC should be a co-sponsor of the symposium noting that ISC sponsorship would consist of symposium participation (oral presentations) and perhaps planning. All Members present agreed that ISC should co-sponsor the symposium, and to encourage local scientists to present relevant research at the symposium. G. DiNardo has agreed to represent ISC on the symposium scientific steering committee and as a co-convener of session 1.

6 REPORT OF SPECIES WORKING GROUPS AND REVIEW OF ASSIGNMENTS

6.1 Albacore

J. Holmes reported on the activities of the ALBWG over the past year (ISC/17/ ANNEX/04/09). The WG held a data preparation workshop 8-14 November 2016 in Nanaimo, Canada (ISC/17/ ANNEX/04), and the stock assessment workshop 11-19 April 2017 in La Jolla, U.S.A. (ISC/17/ ANNEX/09). The WG also conducted a webinar in January 2017 to discuss unresolved data issues prior to the data submission deadline for the stock assessment. The data issues were resolved and the stock assessment was successfully completed. The 2017 assessment includes some important changes in model structure and assumptions relative to previous assessments (see ISC/17/ ANNEX/09 and Section 6a) that the WG believes are important advancements in the north Pacific albacore modeling effort:

- 1. New coherent definitions of fleets, especially the JPN LL and PL fleets, based on area and season that achieve constant length-selectivity; and
- 2. A new standardization procedure for the JPN LL index which shows that the model is able to measure changes in the albacore population due to fishing. The new standardization procedure is a zero-inflated negative binomial mixed effects model, using 5°x5° area and vessel as random effects (Ochi and Kiyofuji 2017: ISC/17/ALBWG.01).

An election was held for a new WG Chair at a half-day meeting in advance of ISC17. Hidetada Kiyofuji (Japan) was unanimously elected Chair of the ALBWG for a three-year term (2017-2020). Steve Teo (U.S.A.) was elected Vice-Chair of the ALBWG for a three-year term (2017-2020).

The ALBWG proposed	a schedule	for 2017/18	as follows:
--------------------	------------	-------------	-------------

Meeting	Dates	Location	Goals
SC13	August 9-17, 2017	Rarotonga, Cook Islands	Steve Teo to present stock assessment results and information
NC13	August 28- September 2, 2017	Busan, Korea	J. Holmes to present stock assessment results and information and discuss MSE progress
MSE Managers Workshop	Proposed dates October 17-19, 2017	Vancouver, Canada	Input on acceptable risk, harvest control rules, operating model
Workshop	March/April 2018	TBD	WG Technical input on MSE modelling issues
1SC18	July 1, 2018	TBD	1-day meeting – catch tables, national contacts, other administrative issues

The ALBWG noted the following ongoing issues affecting its work:

1. Clarification of ALB catches reported by China and Vanuatu to the IATTC and WCPFC (see ISC15 Plenary Report for a fuller description)

2. Clarification of engagement with managers/stakeholders from the IATTC and WCPFC on management strategy evaluation (MSE).

Discussion

Members discussed the proposed dates for and the purpose of the third MSE workshop. Desiree Tommasi, a new hire with CIMEC, University of CA, Santa Cruz, and stationed at the NOAA Southwest Fisheries Science Center, will be coordinating development of the MSE and as such, will plan, with support from the ALBWG, specific topics to be covered at the workshop. Members raised concerns about the proposed dates for the workshop (17-19 October 2017) and asked for the opportunity for more input through the ALBWG to settle on a mutually agreeable date. The new WG Chair was tasked with polling WG members for alternative meeting dates and delineating the purpose of the workshop to clearly define when managers and scientist in their countries could attend the workshop. At the same time, members stressed that the workshop announcement should be circulated as soon as possible so as to maximize the opportunity for managers and other stakeholders to participate.

The ISC Chair was tasked with resolving issues regarding albacore catches reported by China and Vanuatu.

6.2 Pacific Bluefin Tuna

H. Fukuda reported on the key activities of the PBFWG over the past year (ISC/17/ANNEX/07/11). The PBFWG conducts bench mark stock assessments every four years and an update assessment between benchmarks. Because a benchmark stock assessment was completed in 2016, a stock assessment was not conducted this year. An update assessment is scheduled for 2018 and a benchmark assessment in 2020.

Between the two consecutive assessments, the PBFWG updated abundance indices to monitor the most recent trends of spawning stock and recruitment. The updated standardized CPUEs from the Japanese and Taiwanese longline fleets, which represent the relative abundance of large spawners, showed a consistent increase and catch-at-length data from both fleets indicated new modes of smaller fish in the catch. Historically the spawning population was comprised of a single cohort of large fish ranging from 155 to 236 cm. In recent years, the spawning population is comprised of multiple cohorts ranging from 100 to 236 cm.

The updated standardized catch-per-unit-of-effort (CPUE) from the Japanese troll fleet, which represents the relative abundance of age-0 fish recruitment, showed a slight increase in 2015 from that of 2014, but is at a low level. The presenter also introduced the real-time recruitment monitoring program conducted by Japan, and based on the 2016 data reported that the recruitment level in 2016 is likely to be higher than that of 2015 and 2014. The PBFWG considered this positive information for stock recovery, even though the best estimate of recruitment is only available from the assessment.

The results of projections requested by the joint meeting of the WCPFC-NC and IATTC were presented and it was reported that the findings have already been forwarded to the WCPFC-NC and IATTC (ISC/17/ANNEX/11). Stochastic projections of PBF were conducted for 19

harvesting scenarios, which were evaluated by six candidate rebuilding targets under three recruitment assumptions. Performance measures were examined for all scenarios, such as probabilities of achieving candidate rebuilding targets and expected annual yield by flag or area. Trajectories of spawning stock biomass and total yield during 2015-2035 were shown, allowing comparisons between scenarios, as well as a table of performance measures, providing material for further discussion towards the next CMMs. The results were presented to the ISC PBF stakeholder meeting in April 2017. (See Section 6.2.1.)

Finally, the presenter summarized the PBFWG's discussion of the request from WCPFC 13 to define a "drastic" drop in recruitment, which the PBFWG found difficult to do. As a path forward, the WG provided a framework to evaluate how recruitment in the most recent one or two years would affect the probability of the stock rebuilding to the rebuilding target within the specified time period (Figure 6-1).




Figure 6-1. An example of flowchart of development and implementation of an emergency rule based on recruitment index. (Source: ISC/17/ANNEX/07, p. 15)

The development of a framework would require a certain amount of time, because it would entail properly evaluating the ability of the new Japanese recruitment/troll index to predict recruitment, developing a procedure to convert the index to recruitment (relative or absolute), and revising the projection model so it would be capable of the calculation suggested here.

The PBFWG proposed schedule for 2017/18 as follows:

Meeting	Dates	Location
Stock Assessment Update	March 1, 2018	SWFSC, La Jolla, USA
PBFWG Meeting prior to ISC18 (1 day)	July 1, 2018	Korea

Discussion

It was noted that the request from the WCPFPC for an emergency rule is not necessary given the frequency of assessments. Plenary accepted the PBFWG's proposed framework for implementation of an emergency rule process when a drastic drop in recruitment is detected as shown in Figure 6-1. It was noted that regular use of the framework (i.e., associated with each assessment) provides a powerful feedback mechanism for managers to assess whether current management measures will achieve the target.

It was emphasized that assessment-based projections will need to take recruitment variability into account.

While the PBFWG was unable to come up with criteria for identifying a "drastic drop" in recruitment, it was suggested that they look at the patterns of recruitment variability of other teleost stocks as a means of defining the meaning of a "drastic drop."

6.2.1 PBF Stakeholders' Meeting

G. DiNardo provided an overview of the PBF International Stakeholders Meeting (ISC/17/ANNEX/14). The International Scientific Committee for Tuna and Tuna-like Species in the North Pacific Ocean (ISC) hosted the First Pacific Bluefin Tuna International Stakeholders Meeting at the Mita Conference Hall (Mita Kaigisho) in Tokyo, Japan from 25-27 April 2017. The Objective of the meeting was to discuss the expected performance of 37 harvest scenarios relative to six candidate rebuilding targets and 9 performance metrics, under a range of assumptions regarding future recruitment conditions (low, average and combined low to average), to facilitate discussion on the selection of the next interim Pacific bluefin tuna (PBF) rebuilding target scheduled for decision at the August 2017 WCPFC-Northern Committee Meeting. Dr. Gerard DiNardo, Fisheries Resources Division Director at NOAA Fisheries, Southwest Fisheries Science Center and Chairman of the ISC, along with Mr. Masanori Miyahara, Japan Fisheries Research and Education Agency President and Chairman of the WCPFC-Northern Committee (NC), co-chaired the event.

As this was a public meeting, stakeholders were urged to candidly express their perceptions regarding the status of PBF, as well as perspectives on future rebuilding targets and potential rebuilding strategies to achieve the targets. Stakeholders were also reminded that at the 2016 WCPFC Commission Meeting the NC was requested to take due account of two specific suggestions: rebuilding PBF SSB to 20% SSB0 by 2034 and development of emergency rules concerning drastic drops in recruitment and associated risks. While no final decisions were expected at this meeting, the discussions will inform decisions on an interim rebuilding strategy and target at the 2nd Joint IATTC-NC Meeting scheduled for August 2017.

Approximately 150 stakeholders participated in the event, including resource managers, scientists, industry (fisherman, farmers, retailers, and processors), representatives from Pacific Ocean tuna RFMOs, environmental organizations, and other stakeholders interested in PBF. Dr. H. Fukuda (NRIFSF) made a presentation entitled Pacific bluefin tuna (PBF) stock assessment and related information. Dr. Gerard DiNardo presented the additional PBF projections requested

by IATTC and WCPFC and completed by Akita et al. (2017), that were reviewed during the February 2017 PBFWG Workshop, as well as a summary of the findings.

While all participants were committed to rebuilding PBF as soon as feasibly possible, the need to restructure fishing activities was also recognized. However, despite a full day of presentation materials on the critically low status of the PBF stock, many stakeholders either believed that the stock was already rebuilding, or that changing environmental conditions was the reason for low stocks. Large-scale industry stakeholders urged the ISC and the Government of Japan not to implement harsh measures, as fishery economic conditions in Japan are dire. The projection results constitute new information relative to PBF conservation advice and these are the conclusions as reported:

- Different recruitment scenarios forecast entirely different levels of SSB in the future;
- Under average recruitment conditions, all harvest scenarios achieve the initial rebuilding target of SSB_{MED1952-2014} by 2024;
- Under all recruitment conditions with zero removals (no fishing), SSB trajectories achieved all rebuilding targets by approximately 2020 and the initial rebuilding target, SSB_{MED1952-2014}, within 2-3 years;
- Achieving 20% SSB0 during the projection period was not possible in most of the low recruitment scenarios;
- The probability of SSB falling below the historical lowest level at any time during the projection period is low (< 2%) in all projections;
- Scenarios that do not have catch limits for large fish in the EPO and WPO, or has a higher catch limit for large fish in WPO, do not achieve the initial rebuilding target, SSB_{MED1952-2014}, by 2024 under low recruitment conditions; and
- Reducing the catch of small fish results in positive impacts on SSB trajectories, even with increases in the catch of large fish in the WPO.

Discussion

The ISC Chair thanked members of the PBFWG for completing the WCPFC-NC and IATTC harvest scenario projections and participating in the PBF Stakeholders' Meeting.

6.3 Billfish

J. Brodziak provided the BILLWG Report (**ISC/17/ANNEX/10**). The BILLWG conducts all of its work in accord with the scientific method which includes empirical testing, open debate, documentation and reproducibility, reporting uncertainty, and peer review. The BILLWG focuses on an ongoing consistent application of this method to produce the best available scientific information for the assessment of billfish stocks in the Pacific Ocean.

The BILLWG held one intersessional meeting during the ISC17 work cycle for providing assessment information and conservation advice to the ISC17 Plenary.

BILLWG June 2017 Workshop

- 1. BILLWG Assessment Research Workshop was held in Keelung, Taiwan, at National Taiwan Ocean University during 1-7 June 2017 with participants from Japan, Taiwan, and U.S.A.
- 2. The primary goals of this workshop were (1) to review and update the standardization of Japanese longline CPUE for swordfish and (2) to review and finalize North Pacific swordfish population structure for the 2018 assessment.
- 3. Secondary goals of this workshop were to:
 - a. Review and update billfish biological and fishery information for the ISC BILLWG webpage;
 - b. Analyze and describe the spatiotemporal distributions of juvenile and adult swordfish; and
 - c. Identify and prioritize uncertainties in billfish stock assessments for management strategy evaluation.

All of the ISC16 BILLWG goals and associated work assignments were addressed at the June 2017 workshop. Six working papers were presented and reviewed by the BILLWG to this end.

In addition, one research presentation on swordfish genetic structure was considered. This was "Bayesian analyses of Pacific swordfish (*Xiphias gladius*) genetic differentiation using multilocus single nucleotide polymorphism data" by Ching-Ping Lu based on Lu, et al. (2016. *Journal of Experimental Marine Biology and Ecology*. 482:1-17).

The BILLWG noted that the key findings of Lu's new research on swordfish genetics were:

- Overall the new research suggested that there were two major swordfish genetic clusters in the Pacific Ocean and these were so-called <u>tropical</u> and <u>temperate genetic patterns</u>.
- The genetic model in Lu, et al. (2016) shows that swordfish exhibit complex genetic population structure in the Pacific.

In addition, the BILLWG noted that there is new information on individual swordfish movements across the southern boundary of the EPO stock area at 20°S latitude, which was provided by the tagging data in the study of Abascal, et al. (2010. Horizontal and vertical movements of swordfish in the Southeast Pacific. *ICES Journal of Marine Science*, 67: 466–474). This paper showed a consistent migratory pattern with fish moving northwest across the stock boundary at 20° S latitude by autumn and presumably returning south by early spring.

Based on the new information on swordfish genetic and movement patterns along with the existing status quo treatment of North Pacific stock structure, the BILLWG agreed to proceed with the 2018 benchmark assessment and account for some stock structure uncertainty by assessing:

- 1. The WCNPO stock (boundary same as used in 2014) as a benchmark assessment, attempting to use Stock Synthesis or similar structured model, and
- 2. The single North Pacific swordfish management unit north of the Equator as a supplementary assessment scenario.

Given this decision on how to treat the 2018 swordfish stock assessment, the BILLWG provided the following work plan for 2017-2018.

Resolution of BILLWG Work Plan for 2017-2018

Secondary tasks for the ISC18 work plan were:

- Ongoing development of management strategy evaluation algorithms for ISC BILLWG stocks
- Continue to evaluate stock structure uncertainty for North Pacific billfish stocks
- Complete the update of BILLWG webpage information
- Continue work on billfish population dynamics and fisheries in the North Pacific

The primary task for the ISC18 work plan is to conduct a benchmark assessment for North Pacific swordfish accounting for some stock structure uncertainty, as described above.

The BILLWG proposed schedule for 2017/18 as follows:

Meeting	Dates	Location
2017 Swordfish Data Preparation and Modeling Webinar/Teleconference Meetings: September (SS3 version) DATA PLAN	November 1, 2017	Webinar
2017 Swordfish Data Preparation and Modeling Webinar/Teleconference Meetings: September (SS3 version) DATA PROGRESS	December 1, 2017	Webinar
Swordfish Data Preparation Meeting	January 17-23, 2018	Honolulu, HI, USA
Swordfish Assessment Modeling Meeting	TBD	Spring 2018

The BILLWG also identified assessment scientists to work on the 2018 benchmark North Pacific swordfish stock assessment. These were Dr. Hirotaka Ijima, Japan; Dr. Michelle Sculley, U.S.A.; and Dr. Yi-Jay Chang, Taiwan.

The BILLWG also identified the need to have a data manager for the BILLWG, noting that Darryl Tagami successfully served for 2010-2016 and that there is no current data manager for the BILLWG.

Last the BILLWG resolved the need for a new Chair and Vice Chair by holding elections with

- Jon Brodziak was reelected as Chair for one year (2017-2018)
- Hirotaka Ijima was elected as Vice Chair for three years (2017-2020)

Discussion

The first use of a structured model for the SWO assessment was discussed. It was difficult to achieve SS2 model convergence in a previous attempt for the EPO stock, in part because of uncertainty about the growth curve. The WCNPO stock did not have the same convergence issues in the previous attempt. The BILLWG Chair also reviewed oceanographic and other information used to determine the current stock boundary. Recent genetic work is reasonably consistent with this stock boundary.

The ISC Chair recommended that the date and location for the proposed 2018 BILLWG assessment modeling meeting be determined soon. He also emphasized the need for the BILLWG to appoint a data manager. This appointment is the WG Chair's responsibility.

The ISC Chair noted that as MSE research for billfish stocks advances it will be important to shift to a format that allows stakeholder engagement. The BILLWG Chair emphasized that current work is focused on resolving preliminary scientific uncertainty questions as a first step. As progress is made stakeholders will be invited to participate. It was noted that any MSE is unlikely to be conducted before 2020.

6.4 Shark

S. Kohin, SHARKWG Chair, provided a summary of SHARKWG activities over the past year (ISC/17/ANNEX/05/08/13). The focus of the SHARKWG was on blue shark (BSH) with the goal of completing a benchmark North Pacific BSH stock assessment by ISC 17 Plenary. Full meetings of the SHARKWG since ISC16 were held in Busan, Republic of Korea and La Jolla, CA, U.S.A. The SHARKWG also held two webinars between meetings to review data updates and plan for the BSH assessment. Other SHARKWG activities included a meeting of a subgroup of assessment modelers in advance of the full assessment workshop, participation in the STATWG Steering Committee meeting, a webinar involving the subgroup of shark age and growth specialists, and a short meeting in advance of ISC17 to conduct administrative business and plan for the coming year. Canada, China, Chinese Taipei, Japan, Korea, Mexico, U.S.A., IATTC and the WCPFC all actively participated in at least one SHARKWG meeting or webinar. Although BSH was the focus of the Working Group over the past year, some information on SMA fishery data and biology was also discussed as the Working Group prepares for the SMA full assessment in 2018.

Highlights of the meetings and webinars were briefly presented; ISC/17/ANNEX/5/8 contain the reports of the full Working Group meetings. The SHARKWG Chair expressed appreciation to Korea for hosting the BSH data preparatory meeting and to U.S.A. for hosting the assessment workshop. Through the hard work of WG members at the meetings and during the intercessional webinars, the SHARKWG completed a very thorough and rigorous, fully-integrated BSH benchmark assessment that is based on the most complete fishery and biological data compiled for a pelagic shark species to date.

The SHARKWG proposed a work plan for the coming year to complete the first full assessment of North Pacific SMA before ISC18. The SHARKWG recognizes the difficulty in estimating shark catch and discards and the challenges presented by spatial segregation of pelagic sharks by

size and sex. In addition, life history parameters for pelagic sharks are still rather uncertain. Work leading up to ISC18 will focus on improving catch and CPUE time series for SMA and advancing research on SMA growth. SHARKWG age and growth specialists are working to corroborate age readings across labs and will convene a workshop in October 2017 to conduct meta-analyses of SMA vertebrae band pair readings and produce consensus growth curves for the SMA assessment.

Continued improvements in the monitoring of BSH catches, through carefully designed observer programs and species-specific logbooks, including recording the size and sex of sharks retained and discarded for all fisheries, as well as continued research into the biology and ecology of BSH in the North Pacific are recommended.

Suzanne Kohin has completed her final term as SHARKWG Chair. Mikihiko Kai of Japan was elected Vice Chair of the SHARKWG and will act as interim Chair until the SHARKWG meets in November 2017 when it will hold an election for Chair. It was recommended that the ISC Chair conduct the WG chair election at the November meeting.

The SHARKWG proposed schedule for 2017/18 as follows:

Meeting	Dates	Location
Mako ageing webinar	September/ October 2017	Webinar
Mako ageing workshop	October 19 - 24, 2017	Shizuoka, Japan
Mako data prep webinar	Early November 2017	Webinar
Mako data prep meeting	November 27 - December 1, 2017	Shizuoka, Japan
Mako data prep webinar	January/February 2018	Webinar
Mako assessment	March/April 2018	La Jolla, CA
meeting	March/April 2018	USA

Discussion

The availability of catch estimate time series data was reviewed; these data are necessary to conduct a fully structured SMA assessment. Most catch is attributable to Chinese Taipei, Japan, Mexico, and the U.S.A. and these Members have committed to provide the necessary data.

A new Chair needs to be elected since the current Chair's term is ending. There is also a need to identify the team lead for the SMA assessment and a data manager. In general, it is better if WGs address the election of officers at its regular meetings when more members are in attendance, rather than during meetings that precede the Plenary. This gives members more time to consult on who can fulfill those roles.

It was noted that both spatial differences and sexual dimorphism in growth have been observed for BSH and SMA. Sexual dimorphism in growth will be addressed in the upcoming SMA growth and age workshop.

The SHARKWG Chair reached out to SPC about collaborating on the 2017 BSH benchmark stock assessment but they were not able to contribute.

There have been difficulties in obtaining needed data from the WCPFC and IATTC. For example, for the BSH assessment, public data from the RFMOs were used to update China fisheries time series but some data were excluded under data confidentiality rules. One of the first tasks of the new ISC Chair will be to engage with these organizations to resolve data provision problems. Related to data issues, the Chair emphasized the importance of the species WGs engaging with the STATWG when assistance is needed on external data requests.

7 STOCK STATUS AND CONSERVATION INFORMATION

7.1 Albacore

J. Holmes, ALBWG Chair, summarized results of the benchmark ALB stock assessment (ISC/17/ANNEX/12). There were three major changes to the 2017 base case model relative to the 2014 assessment. First, the model time frame was shortened to 1993-2015 (1966-2012 in the previous assessment) to eliminate the influence of poorly fit size composition data in the 1975-1992 period, and a conflict between these size composition data and the primary adult albacore index. Second, a new procedure was used to standardize the Japanese longline abundance index (1996 – 2015) used to indicate trends in adult albacore abundance and the results represent a substantial improvement relative to 2014 and earlier assessments. This new index had good contrast and was informative on both population trend and scale, and was the only index fitted in the 2017 base case model. Third, sex-specific M-at-age vectors were developed from a meta-analysis, with a sex-combined M that scaled with size for ages 0-2, and annual sex-specific natural mortality (M) fixed at 0.39 and 0.48 for age-3+ males and females, respectively, because support for the previous assumption (M fixed at 0.3 y-1 for both sexes at all ages) is poor in the scientific literature.

All available fishery data for north Pacific albacore for the 1993 to 2015 period were used in the stock assessment and it was assumed that there is instantaneous mixing of albacore on a quarterly basis. Catches during the modeling period (1993-2015) reached a peak of 119,300 t in 1999 and then declined in the early 2000s, followed by a recovery in later years with catches fluctuating between 68,900 and 93,100 t in recent years (2010-2015). Surface gears (troll, pole-and-line), which primarily harvest juvenile albacore, have accounted for approximately two-thirds of the reported albacore catch and longline gear, which primarily harvest adult albacore, the remaining third.

Catch and size composition data from ISC countries (Canada, China, Chinese Taipei, Japan, Korea, and U.S.A.) and catch data from some IATTC and WCPFC member countries, were obtained for the assessment and assigned to 29 fisheries defined for this assessment (based on flag, gear, area, and season). Thirteen indices of abundance (standardized catch-per-unit-effort) were provided by Japan, U.S.A., Canada, and Chinese Taipei but only the index for the Japan LL in Area 2 was fitted in the base case model. A sensitivity run fitting to the Japan pole-and-line (PL) index was also conducted. Size composition data were available from 18 fisheries.

The North Pacific albacore tuna stock was assessed using a length-based, age-, and sexstructured Stock Synthesis (SS Version 3.24AB) model over the 1993-2015 period. Sex-specific growth curves from the 2014 assessment were used because of evidence of sexually dimorphic growth, with adult males attaining a larger size-at-age than females after maturity. Sex-specific M-at-age vectors were developed from a meta-analysis, with a sex-combined M that scaled with size for ages 0-2, and annual sex-specific M fixed at 0.48 and 0.39 for age-3+ females and males, respectively. The steepness of the Beverton-Holt stock-recruitment relationship was assumed to be 0.9, based on two prior analyses. The assessment model was fitted to the Japan LL index (1996-2015; Area 2) and all representative size composition data in a likelihood-based statistical framework. All fleets were assumed to have dome-shaped length selectivity, and age-based selectivity for ages 1-5 was also estimated for surface fleets (troll and pole-and-line) to address age-based changes in juvenile albacore availability and movement. Selectivity was assumed to vary over time for fleets with important changes in fishing operations. Maximum likelihood estimates of model parameters, derived outputs, and their variances were used to characterize stock status. Several sensitivity analyses were conducted to evaluate changes in model performance or the range of uncertainty resulting from changes in model parameters, including natural mortality, stock-recruitment steepness, starting year, selectivity estimation, variability of size-at-age and weighting of size composition data.

An age-structured production model diagnostic analysis showed that the estimated catch-at-age and fixed productivity parameters (growth, mortality and stock-recruitment relationship without annual recruitment deviates) were able to explain trends in the Japan longline index. Based on these findings, the ALBWG concluded that the base case model was able to estimate the stock production function and the effect of fishing on the abundance of the NPALB stock. The link between catch-at-age and the Japan longline index adds confidence to the data used, and represents a major improvement in the 2017 assessment of the north Pacific albacore stock. Due to the moderate exploitation levels relative to the productivity, the production function was weakly informative about NPALB stock size, resulting in asymmetric uncertainty in the absolute scale of the stock, with more uncertainty in the upper limit of the spawning stock biomass than the lower limit of spawning stock biomass. It is important to note that the primary aim of estimating the female spawning biomass (SSB) in this assessment was to determine if the estimated SSB was lower than the limit reference point (i.e., determine whether the stock is in an overfished condition). Since the lower bound is better defined, it adds confidence to the ALBWG's evaluation of stock condition relative to the limit reference point.

Discussion

With regard to the benchmark stock assessment, ongoing uncertainty about growth parameters was discussed. Growth parameters appear to vary both spatiotemporally as well as by sex. The ALBWG tried looking at regional growth models in the past but they did not perform well in the sensitivity runs. There is a need for Pacific-wide data collection to address this uncertainty. The use of a sex-specific genetic marker for albacore is in development and may be available in the next year or so. There will then be a need to design and implement a sampling program to gather samples and do the analysis.

There was confusion over the use of the term "fishing intensity" and how this differs from the more familiar term fishing mortality. Generally, it was agreed that the term should be clearly

defined when used in presenting stock status information. While including the supporting tables and figures from the stock assessment report (ISC/17/ANNEX/11) in the Plenary Report is useful, presenting a concise, clear statement to fishery managers is paramount.

The difference between the projections of stock status under a constant fishing intensity scenario versus a constant catch scenario was discussed at length. It was noted that constant fishing intensity is probably a closer approximation of the current effort-based management measure adopted by both Pacific RFMOs. However, it may be necessary to provide a straightforward explanation of why stock biomass is projected to decline more under a constant catch scenario based on recent catch levels. Recent recruitment has supported higher catches under recent fishing intensity levels.

The Plenary endorsed the NPALB stock assessment and considers it to be the best available scientific information.

Stock Status and Conservation Information

Stock Status

Estimated total stock biomass (males and female at age-1+) declines at the beginning of the time series until 2000, after which biomass becomes relatively stable (Figure 7-1A). Estimated female SSB exhibits a similar population trend, with an initial decline until 2003 followed by fluctuations without a clear trend through 2015 (Figure 7-1B). The estimated SPR (spawners per recruit relative to the unfished population) in 2015 is 0.53, which corresponds to a moderate fishing intensity (i.e., 1-SPR = 0.47). Instantaneous fishing mortality at age (F-at-age) is similar in both sexes through age-5, peaking at age-4 and declining to a low at age-6, after which males experience higher F-at-age than females up to age 13 (Figure 7-2). Juvenile albacore aged 2 to 4 years comprised, on average, 70% of the annual catch between 1993 and 2015 (Figure 7-3) as reflected by the larger impact of the surface fisheries (primarily troll, pole-and-line) which remove juvenile fish, relative to longline fisheries, which primarily remove adult fish (Figure 7-4).

The WCPFC adopted a limit reference point (LRP) for the spawning stock biomass of 20% of the spawning stock biomass when F=0 (20% SSB_{current, F=0}) (https://www.wcpfc.int/harvest-strategy). The 20% SSB_{current, F=0} LRP is based on dynamic biomass and fluctuates depending on changes in recruitment. However, there are no adopted F-based limit reference points.

Stock status is depicted in relation to the limit reference point (LRP; $20\%SSB_{current, F=0}$) for the stock and the equivalent fishing intensity (F_{20%}; calculated as 1-SPR_{20%}) (Figure 7-5A). Fishing intensity (F, calculated as 1-SPR) is a measure of fishing mortality expressed as the decline in the proportion of the spawning biomass produced by each recruit relative to the unfished state.

The Kobe plot shows that the estimated female SSB has never fallen below the LRP since 1993, albeit with large uncertainty in the terminal year (2015) estimates (Figure 7-5A). Even when alternative hypotheses about key model uncertainties such as natural mortality and growth were evaluated, the point estimate of female SSB in 2015 (SSB₂₀₁₅) did not fall below the LRP, although the risk increases with these more extreme assumptions (Figure 7-5B). The SSB₂₀₁₅ was

estimated to be 80,618 t and was 2.47 times greater than the LRP threshold of 32,614 t (Table 7-1). Current fishing intensity, $F_{2012-2014}$ (calculated as 1- $SPR_{2012-2014}$), was lower than potential F-based reference points identified for the north Pacific albacore stock, except $F_{50\%}$ (calculated as 1- $SPR_{50\%}$) (Table 7-1).

Based on these findings, the following information on the status of the north Pacific albacore stock is provided:

- 1. The stock is likely not overfished relative to the limit reference point adopted by the Western and Central Pacific Fisheries Commission (20%SSB current F=0), and
- 2. No F-based reference points have been adopted to evaluate overfishing. Stock status was evaluated against seven potential reference points. Current fishing intensity $(F_{2012-2014})$ is below six of the seven potential reference points (see ratios in Table 7-1), except $F_{50\%}$.

Conservation Information

Two harvest scenarios were projected to evaluate impacts on future female SSB: F at the 2012-2014 rate over 10 years ($F_{2012-2014}$) and constant catch² (average of 2010-2014 = 82,432 t) over 10 years. Median female SSB is expected to decline to 63,483 t (95% CI: 36,046 - 90,921 t) by 2025, with a 0.2 and <0.01 % probability of being below the LRP by 2020 and 2025, respectively, if fishing intensity remains at the 2012-2014 level³ (Figure 7-6). In contrast, employing the constant catch harvest scenario is expected to reduce female SSB to 47,591 t (95% CI: 5,223 - 89,958 t) by 2025 and increases the probability that female SSB will be below the LRP to about 3.5 and 30 % in 2020 and 2025, respectively (Figure 7-7). In addition, as biomass declines during the projection period the fishing intensity approximately doubles by 2025. The probabilities of declining below the LRP in both harvest scenarios are likely higher in the future because projection results did not capture the full envelope of uncertainty. The ALBWG notes that the lack of sex-specific size data, uncertainty in growth and natural mortality, and the simplified treatment of the spatial structure of north Pacific albacore population dynamics are important sources of uncertainty in the assessment.

Based on these findings, the following conservation information is provided:

1. If a constant fishing intensity $(F_{2012-2014})$ is applied to the stock, then median female spawning biomass is expected to undergo a moderate decline, with a < 0.01% probability of falling below the limit reference point established by the WCPFC by 2025. However, expected catches in this scenario will be below the recent average catch level for this stock.²

² It should be noted that the constant catch scenario is inconsistent with current management approaches for NPALB adopted by the IATTC and the WCPFC.

³ Median future catch for the constant F scenario is expected to be below the average catch level for 2010-2014 (82,432 t – red line in Figure 7-6). This result is likely due to low estimated recruitment in 2011, which is expected to reduce female SSB beginning in 2015, the first year of the projection period.

If a constant average catch ($C_{2010-2014} = 82,432$ t) is removed from the stock in the future, then the decline in median female spawning biomass will be greater than in the constant F intensity scenario and the probability that SSB falls below the LRP will be greater by 2025 (30%). Additionally, the estimated fishing intensity will double relative to the current level ($F_{2012-2014}$) by 2025 as spawning biomass declines.




Figure 7-1. Maximum likelihood estimates of (A) total age-1+ biomass (open circles) (B), female spawning biomass (SSB) (solid blue line), and (C) age-0 recruitment (open circles) of north Pacific albacore tuna (*Thunnus alalunga*). Dashed lines (B) and vertical bars (C) indicate 95% confidence intervals of the female SSB and recruitment estimates respectively. Red line indicates the 20%SSB_{current, F=0} limit reference point, which is based on dynamic SSB₀. Closed black circle and error bars in (B) are the maximum likelihood estimate and 95% confidence intervals of unfished female spawning biomass, SSB₀. Since the assessment model represents time on a quarterly basis, there are four estimates of total biomass (A) for each year, but only one annual estimate of female SSB (B) and recruitment (C).




Figure 7-2 Estimated sex-specific instantaneous fishing mortality-at-age (F-at-age) for the 2017 base case model, averaged across 2012-2014.




Figure 7-3. Historical catch-at-age of north Pacific albacore (*Thunnus alalunga*) estimated by the 2017 base case model.





Figure 7-4. Fishery impact analysis on north Pacific albacore (*Thunnus alalunga*) showing female spawning biomass (SSB) (red) estimated by the 2017 base case model as a percentage of dynamic unfished female SSB (SSB₀). Colored areas show the relative proportion of fishing impact attributed to longline (USA, Japan, Chinese-Taipei, Korea and others) (green) and surface (USA, Canada, and Japan) (blue) fisheries (primarily troll and pole-and-line gear, but including all other gears except longline).

Table 7-1. Estimates of maximum sustainable yield (MSY), female spawning biomass (SSB) quantities, and fishing intensity (F) based reference point ratios for north Pacific albacore tuna for the base case assessment and important sensitivity analyses. SSB_0 and SSB_{MSY} are the unfished biomass of mature female fish and at MSY, respectively. The Fs in this table are not based on instantaneous fishing mortality. Instead, the Fs are indicators of fishing intensity based on SPR and calculated as 1-SPR so that the Fs reflects changes in fishing mortality. SPR is the equilibrium SSB per recruit that would result from the current year's pattern and intensity of fishing mortality. Current fishing intensity is based on the average fishing intensity during 2012-2014 ($F_{2012-2014}$).

Quantity	Base Case	$M = 0.3 \text{ y}^{-1}$	$\begin{aligned} &Growth \\ &CV = 0.06 \text{ for } L_{inf} \end{aligned}$
MSY (t) A	132,072	92,027	118,836
$SSB_{MSY}(t)^{B}$	24,770	42,098	22,351
$SSB_0(t)^B$	171,869	270,879	156,336
SSB_{2015} (t) ^B	80,618	68,169	63,719
$SSB_{2015}/20\%SSB_{current, F=0}$ B	2.47	1.31	2.15
$F_{2012-2014}$	0.51	0.74	0.57
$F_{2012-2014}/F_{MSY}$	0.61	0.89	0.68
$F_{2012-2014}/F_{0.1}$	0.58	0.90	0.65
$F_{2012-2014}/F_{10\%}$	0.56	0.81	0.63
$F_{2012-2014}/F_{20\%}$	0.63	0.91	0.71
$F_{2012-2014}/F_{30\%}$	0.72	1.04	0.81
$F_{2012-2014}/F_{40\%}$	0.85	1.21	0.96
$F_{2012-2014}/F_{50\%}$	1.01	1.47	1.16

A – MSY includes male and female juvenile and adult fish

B – Spawning stock biomass (SSB) in this assessment refers to mature female biomass only.






Figure 7-5. (A) Kobe plot showing the status of the north Pacific albacore (*Thunnus alalunga*) stock relative to the 20%SSBcurrent, F=0 biomass-based limit reference point (WCPFC), and equivalent fishing intensity (F20%; calculated as 1-SPR20%) over the base case modeling period (1993-2015). Note that no F-based reference point is established for north Pacific albacore by either the IATTC or WCPFC. F20% was chosen for this plot as a convenient analogue to the limit reference point but it should not be inferred that it represents advice on the choice of an F-based reference point on this stock. Black triangle indicates the start year (1993) and black circle with 95% confidence intervals indicates the terminal year (2015). (B) Kobe plot showing stock status and 95% confidence intervals in the terminal year (2015) of the base case model (black; closed circle) and important sensitivity runs with M=0.3 y-1 for both sexes (pen square), and CV=0.06 for L_{inf} in the growth model (white; open triangle). Fs in this figure are not based on instantaneous fishing mortality. Instead, the Fs are indicators of fishing intensity based on SPR and calculated as 1-SPR so that the Fs reflects changes in fishing mortality. SPR is the equilibrium SSB per recruit that would result from the current year's pattern and intensity of fishing mortality.




Figure 7-6. (A) Historical and future trajectory of north Pacific albacore (*Thunnus alalunga*) female spawning biomass (SSB) under a constant fishing intensity ($F_{2012-2014}$) harvest scenario. Future recruitment was based on the expected recruitment variability and autocorrelation. Black line and blue area indicates maximum likelihood estimates and 95% confidence intervals (CI), respectively, of historical female SSB, which includes parameter uncertainty. Red line and red area indicates mean value and 95% CI of projected female SSB, which only includes future recruitment variability and SSB uncertainty in the terminal year. (B) Expected annual catch under a constant fishing intensity ($F_{2012-2014}$) harvest scenario (2016-2025). The red line is the current average catch (2010-2014 = 82,432 t).




Figure 7-7. (A) Historical and future trajectory of north Pacific albacore (*Thunnus alalunga*) female spawning biomass (SSB) under a constant catch (average 2010-2014 = 82,432 t) harvest scenario. Future recruitment was based on the expected recruitment variability and autocorrelation. Dashed line indictaes the average limit reference point threshold for 2012-2014. Black line and blue area indicates maximum likelihood estimates and 95% confidence intervals (CI), respectively, of historical female SSB, which includes parameter uncertainty. Red line and red area indicates mean value and 95% CI of projected female SSB, which only includes future recruitment variability and SSB uncertainty in the terminal year. (B) Projected fishing intensity relative to the current fishing inetnsity (2012-2014) (red line) under a constant catch scenario (average 2010-2014).

7.2 Blue shark

S. Kohin, Chair of the SHARKWG Chair, summarized the results of the 2017 benchmark North Pacific BSH assessment (ISC/17/ANNEX/13). The stock was assessed using a fully-integrated sex-specific Stock Synthesis (SS version 3.24f) model using data from 1971-2015. The SHARKWG also conducted a second assessment using a Bayesian state-space production model (BSSPM) in order to provide continuity with the 2014 ISC BSH assessment.

Data for the assessment were compiled for 18 fishing fleets and included catch time series from 1971 to 2015, six abundance indices, and sex-specific size data. A two-sex base case model that included all the size and catch data, abundance indices from the Japanese Kinkai shallow-set longline fishery, and a low fecundity stock recruitment relationship (LFSR) was developed using an iterative process. Model fits to the data and selection of the base case parameterization was based on a full suite of diagnostics. Additional models were run to explore uncertainty in biological and fishery data, and model performance.

Discussion

Variation in BSH CPUE across fleets was discussed in relation to potential spatial differences. It appears that trends in CPUE are different in the EPO compared to the WCPO. There may be a

relationship between BSH CPUE and temperature; the EPO has been relatively warmer in the last few years and this may partially explain the recent lower CPUE.

The assessment team focused on improving model fit by modifying model structure instead of down-weighting data. Even with this approach some minor down-weighting was done

The Plenary endorsed the BSH stock assessment and considers it to be the best available scientific information.

Stock Status and Conservation Information

Stock Status


The assessment uses a fully integrated, sex-specific Stock Synthesis (SS) model. Model inputs have been greatly improved since the previous assessment. The main differences between the present assessment and the 2014 assessment are: 1) use of SS with a thorough examination of the size composition data, and the relative weighting of CPUE and composition data; 2) improved life history information, such as growth and reproductive biology, and their contribution to productivity assumptions; 3) an improved understanding and parametrization of the low fecundity stock-recruit relationship (LFSR); 4) catch, CPUE, and size time series updated through 2015; 5) a suite of model diagnostics including implementation of an age structured production model implemented in SS. There remain some uncertainties in the time series based on the quality (observer versus logbook) and timespan of catch and relative abundance indices, limited size composition data for several fisheries, potential under-reporting of catch, and life history parameters.


Extensive model explorations showed that the reference run had the best model performance and showed fits most consistent with the data. The CPUE indices used in the reference case were considered most representative of the North Pacific blue shark stock due to their broader spatial temporal coverage in the core distribution of the stock and the statistical soundness of the standardizations. Alternate CPUE series for the latter part of the time series produced different stock trajectories depending upon the index used, but in each case, median SSB during the last three years exceeded SSB_{MSY}. Using alternate assumptions on stock productivity (i.e., form of the stock recruitment relationship) also resulted in variation in the stock trajectories; assuming stock productivity lower than supported by current biological studies, resulted in lowered spawning stock biomass relative to MSY.

Results of the reference case model showed that the spawning stock biomass was near a time-series high in the late 1970s, declined to its lowest level between 1990 to 1995, subsequently increased gradually to reach the time-series high again in 2005, and has since shown small fluctuations with no apparent trend (Figure 7-8 B)close to the time-series high. Recruitment has fluctuated around 37,000,000 age-0 sharks annually with no apparent trend (Figure 7-8 A). Stock status is reported in relation to maximum sustainable yield (MSY) based reference points. Based on these findings, the following information on the status of the North Pacific blue shark stock is provided:

1. Female spawning biomass in 2015 (SSB₂₀₁₅) was 69% higher than at MSY and estimated to be 295,774 t (Table 7-2; Figure 7-8).

- 2. The recent annual fishing mortality ($F_{2012-2014}$) was estimated to be well below F_{MSY} at approximately 38% of F_{MSY} (Table 7-2; Figure 7-8).
- 3. The reference run produced terminal conditions that were predominately in the lower right quadrant of the Kobe plot (not overfished and overfishing not occurring) (Figure 7-9).








Figure 7-8. Results of the SS stock assessment reference case model for the North Pacific BSH assessment: (A) estimated age-0 recruits (circles) and 95% confidence intervals (vertical bars); (B) estimated female spawning biomass and 95% confidence intervals (blue shaded area); (C) estimated fishing mortality (sum of F's across all fishing fleets). Red solid lines indicate the estimates of SB_{MSY} and F_{MSY} in (B) and (C), respectively.

Table 7-2. Estimates of key management quantities for the North Pacific blue shark SS stock assessment reference case model and the range of values for 13 sensitivity runs.

Management Quantity	Reference Case Model	Range for Sensitivity Runs
SSB ₁₉₇₁	301,739 t	174,381 - 980,878 t
SSB_{2015}	295,774 t	140,742 - 1,082,300 t
SSB_{MSY}	175,401 t	100,984 - 482,638 t
F_{1971}	0.15	0.01 - 0.15
$F_{2012-2014}$	0.14	0.06 - 0.15
F_{MSY}	0.36	0.26 - 0.66
$SSB_{2015}\!/SSB_{MSY}$	1.69	1.39 - 2.59
$F_{2012-2014}/F_{MSY}$	0.38	0.15 - 0.50




Figure 7-9. Kobe plots of the trends in estimates of relative fishing mortality and biomass of North Pacific blue shark between 1971 and 2015 for the reference case of (A) the SS stock assessment model, and (B) the BSSPM stock assessment model.

Conservation Information

Target and limit reference points have not yet been established for pelagic sharks by the WCPFC or the IATTC, the organizations responsible for management of pelagic sharks caught in international fisheries for tuna and tuna-like species in the Pacific Ocean.

The 2015 SSB exceeds SSB_{MSY} and $F_{2012-2014}$ is below F_{MSY} . Future projections under different fishing mortality (F) harvest policies (status quo, +20%, -20%, F_{MSY}) show that median BSH spawning biomass in the North Pacific will likely remain above SSB_{MSY} in the foreseeable future

(Table 7-3, Figure 7-10). Other potential reference points were not considered in these evaluations.

Table 7-3. Projected trajectory of spawning biomass (in metric tons) for alternative harvest scenarios.

Year	Average F + 20%	$\mathbf{F}_{\mathbf{MSY}}$	Average F - 20%	Average F (2012-2014)
2015	295,774	295,774	295,774	295,774
2016	306,782	306,782	306,782	306,781
2017	316,169	312,081	318,183	317,173
2018	322,317	312,329	326,829	324,559
2019	324,795	310,499	332,111	328,419
2020	326,757	307,209	336,929	331,782
2021	328,323	303,909	341,210	334,675
2022	329,623	300,842	344,988	337,181
2023	330,719	298,091	348,286	339,349
2024	331,656	295,663	351,138	341,218




Figure 7-10. Comparison of future projected North Pacific blue shark spawning biomass under different F harvest policies (*status quo*, +20%, -20%, and F_{MSY}) using the SS reference case model. *Status quo* fishing mortality was based on the average from 2012-2014.

7.3 Shortfin Mako Shark

S. Kohin noted that a SMA stock assessment was not conducted by the SHARKWG (ISC/17/ANNEX/5/8) in the past year.

Discussion

The Plenary agreed to forward the stock status and conservation information statements adopted at ISC16 for SMA except for the omission of accompanying figures and tables with clarifying modifications

Stock Status and Conservation Information

SMA was last assessed in 2015 and the ISC Plenary endorsed the stock status and conservation information adopted at ISC16 (see pp. 62-63 in the <u>ISC16 Plenary Report</u>), which is reproduced here with slight clarifying modifications.

Stock Status

Shortfin mako is a data poor species. Recognizing that information on important fisheries is missing, the untested validity of indicators for determining stock status, and conflicts in the available data, stock status (overfishing and overfished) could not be determined. Managers should consider the undetermined stock status of SMA in the North Pacific when developing and implementing management measures.

The SHARKWG reviewed a suite of information to determine the stock status of SMA shark in the North Pacific. Of the three indices considered to have the greatest value in providing stock status information (Japan shallow-set longline, Hawaii shallow-set longline and Hawaii deep-set longline), abundance trends in two of the series appear to be stable or increasing, while the abundance trend in the Hawaii shallow-set longline fishery CPUE series appears to be declining.

Conservation Information

It is recommended that data for fisheries operating in the North Pacific for which catch estimates are not yet available be developed for use in the next stock assessment scheduled for 2018 and that available catch and CPUE data be monitored for changes in trends. It is further recommended that data collection programs be implemented or improved to provide species-specific shark catch data for fisheries in the North Pacific Ocean.

7.4 North Pacific Swordfish

J. Brodziak noted that a North Pacific swordfish (SWO) stock assessment was not conducted by the BILLWG in the past year.

Discussion

The Plenary agreed to forward the stock status and conservation information statements adopted at ISC16 for SWO unchanged, except for the omission of accompanying figures and tables and slight clarifying modification.

Stock Status and Conservation Information for the Stock

The SWO stock was last assessed in 2014 and the ISC Plenary endorsed the stock status and conservation information adopted at ISC16 (see pp. 56-61 in the <u>ISC16 Plenary Report</u>), which is reproduced here with clarifying modifications.

Western and Central North Pacific Ocean Stock Status

Catches and harvest rates of WCNPO swordfish had a declining trend from 2007-2011, with exploitable biomass fluctuating around 70,000 t. The Kobe plot shows that the WCNPO swordfish stock did not appear to have been overfished or to have experienced overfishing relative to MSY-based reference points throughout most of the assessment time horizon of 1951-2012.

Based on these results, it is unlikely that the WCNPO swordfish population biomass was below B_{MSY} in 2012 (Pr(B2012 < B_{MSY})=14%). Similarly, it is unlikely that the swordfish population was being fished in excess of harvest rate at MSY (H_{MSY}) in 2012 (Pr($H_{2012} > H_{MSY}$) < 1%).

Eastern Pacific Ocean Stock Status

For the EPO stock, exploitable biomass had a declining trend during 1969-1995 and increased from 31,000 t in 1995 to over 60,000 t in 2010, generally remaining above $B_{\rm MSY}$. Harvest rates were initially low, have had a long-term increasing trend, and likely exceeded $H_{\rm MSY}$ in 1998, 2002, 2003, as well as in 2012, the terminal year of the stock assessment.

The Kobe plot shows that overfishing likely occurred in a few years, but may have occurred from 2010 to 2012. There was a 55% probability that overfishing occurred in 2012, but there was a less than 1% probability that the stock was overfished relative to MSY-based reference points.

Western and Central North Pacific Ocean Stock Conservation Information

Stochastic projections for the WCNPO stock were conducted using eight harvest scenarios through 2016. Results relative to MSY-based reference points show that exploitable biomass would likely remain above 60,720 t (B_{MSY}) through 2016 under the status quo catch or status quo harvest rate scenarios. For the high harvest rate scenarios (i.e., maximum observed harvest rate, 150% of H_{MSY} , 125% of H_{MSY}), exploitable biomass was projected to decline below BMSY by 2016 with harvest rates exceeding H_{MSY} . In contrast, the stock would not be expected to experience any overfishing during 2014-2016 under the status quo catch and status quo harvest rate scenarios.

The risk analyses of harvesting a constant annual catch of WCNPO SWO during 2014-2016 showed that there would be less than 1% probability of the stock being overfished or

experiencing overfishing relative to MSY-based reference points in 2016 if current annual catches (2011-2012) of about 10,000 t were maintained.

The WCNPO SWO stock is healthy ($B_{2010-2012} > B_{MSY}$) and is above the level required to sustain recent harvest rates ($H_{2010-2012} < H_{MSY}$).

Eastern Pacific Ocean Stock Conservation Information

Stochastic projections for the EPO stock show that exploitable biomass will likely have a decreasing trajectory during 2014-2016 under the eight harvest scenarios examined. Under the high harvest rate scenarios (status quo catch, maximum observed harvest rate, 150% of H_{MSY}), exploitable biomass was projected to decline to 31,170 t (B_{MSY}) by 2016 with corresponding harvest rates above H_{MSY} . In comparison, under the status quo harvest rate scenario, exploitable biomass was projected to decline to only 40,000 t by 2016, well above the B_{MSY} level. Overall, the projections showed that if recent high catch levels (9,700 t) persist, exploitable biomass will decrease and a moderate risk (50%) of overfishing will continue to occur.

The risk analyses for harvesting a constant catch of EPO SWO during 2014-2016 showed that the probabilities of overfishing and becoming overfished increased as projected catch increased in the future. Maintaining the current (2010-2012) catch of EPO SWO of approximately 9,700 t would lead to a 50% probability of overfishing in 2016 and a less than 1% probability of the stock being overfished in 2016.

For the EPO SWO stock, overfishing may have occurred from 2010 to 2012, and the average yield of roughly 10,000 t in those years, or almost two times higher than the estimated MSY, is not likely to be sustainable in the long term. While biomass of the EPO stock appears to be nearly twice BMSY, any increases in catch above recent (3-year average 2010-2012) levels should consider the uncertainty in stock structure and unreported catch.

7.5 Western and Central North Pacific Striped Marlin

J. Brodziak noted that an MLS stock assessment was not conducted by the BILLWG in the past year.

Discussion

The Plenary agreed to forward the stock status and conservation information statements adopted at ISC16 for MLS, except for the omission of accompanying figures and tables and slight clarification modifications if necessary.

Stock Status and Conservation Information

The most recent MLS stock assessment was conducted in 2015 and the ISC Plenary endorsed the stock status and conservation information adopted at ISC16 (see pp. 53-55 in the <u>ISC16 Plenary Report</u>), which is reproduced here with slight clarifying modifications.

Stock Status

Estimates of population biomass of the WCNPO MLS exhibit a long-term decline. Population biomass (age-1 and older) averaged roughly 20,513 t, or 46% of unfished biomass during 1975-1979, the first 5 years of the assessment time frame, and declined to 6,819 t, or 15% of unfished biomass in 2013. Spawning stock biomass is estimated to be 1,094 t in 2013 (39% of SSBMSY, the spawning stock biomass to produce MSY). Fishing mortality on the stock (average F on ages 3 and older) is currently high and averaged roughly F = 0.94 during 2010-2012, or 49% above FMSY. The predicted value of the spawning potential ratio (SPR, the predicted spawning output at current F as a fraction of unfished spawning output) is currently SPR2010-2012 = 12% ,which is 33% below the level of SPR required to produce MSY. Recruitment averaged about 308,000 recruits during 1994-2011, which was 25% below the 1975-2013 average. No target or limit reference points have been established for the WCNPO striped marlin stock under the auspices of the WCPFC.

The WCNPO MLS stock is expected to be highly productive due to its rapid growth and high resilience to reductions in spawning potential. The status of the stock is highly dependent on the magnitude of recruitment, which has been below its long-term average since 2007, with the exception of 2010. Changes in recent size composition data in comparison to the previous assessment resulted in changes in fishery selectivity estimates and also affected recruitment estimates. These changes, in turn, affected the scaling of biomass and fishing mortality to reference levels.

When the status of MLS is evaluated relative to MSY-based reference points, the 2013 spawning stock biomass is 61% below SSBMSY (2819 t) and the 2010-2012 fishing mortality exceeds FMSY by 49%. Therefore, overfishing is occurring relative to MSY-based reference points and the WCNPO MLS stock is overfished.

Conservation Information

The stock has been experiencing overfishing since 1977, with the exception of 1982 and 1983, and fishing appears to be impeding rebuilding, especially if recent (2007-2011) low recruitment levels persist. Projection results show that fishing at FMSY could lead to median spawning biomass increases of 25%, 55%, and 95% from 2015 to 2020 under the recent recruitment, medium-term recruitment, and stock recruitment-curve scenarios. Fishing at a constant catch of 2,850 t could lead to potential increases in spawning biomass of 19% to over 191% by 2020, depending upon the recruitment scenario. In comparison, fishing at the 2010-2012 fishing mortality rate, which is 49% above FMSY, could lead to changes in spawning stock biomass of -18% to +18% by 2020, while fishing at the average 2001-2003 fishing mortality rate (F2001-2003=1.15), which is 82% above FMSY, could lead to spawning stock biomass decreases of -32% to -9% by 2020, depending upon the recruitment scenario.

7.6 Pacific Blue Marlin

J. Brodziak noted that a BUM stock assessment was not conducted by the BILLWG in the past year.

Discussion

The Plenary agreed to forward the stock status and conservation information statements adopted at ISC16 for BUM, except for the omission of accompanying figures and tables, and slight clarifying if necessary.

Stock Status and Conservation Information

The most recent BUM stock assessment was conducted in 2016 and the ISC Plenary endorsed the stock status and conservation information adopted at ISC16 (see pp. 46-52 in the <u>ISC16</u> Plenary Report), which is reproduced here with slight clarifying modifications.

Stock Status

Estimates of total BUM stock biomass show a long term decline. Population biomass (age-1 and older) averaged roughly 130,965 t in 1971-1975, the first 5 years of the assessment time frame, and has declined by approximately 40% to 78,082 t in 2014. Female spawning biomass was estimated to be 24,809 t in 2014, or about 25% above SSBMSY. Fishing mortality on the stock (average F, ages 2 and older) averaged roughly F = 0.28 during 2012-2014, or about 12% below FMSY. The estimated spawning potential ratio of the stock (SPR, the predicted spawning output at the current F as a fraction of unfished spawning output) is currently SPR2012-2014 = 21%. Annual recruitment averaged about 897,000 recruits during 2008-2014, and no long-term trend in recruitment was apparent. Overall, the time series of spawning stock biomass and recruitment estimates show a long-term decline in spawning stock biomass and a fluctuating pattern without trend for recruitment. The Kobe plot depicts the stock status relative to MSY-based reference points for the base case model and shows that spawning stock biomass decreased to roughly the MSY level in the mid-2000s, and has increased slightly in recent years.

Based on the results of the 2016 stock assessment update, the Pacific blue marlin stock is not currently overfished and is not experiencing overfishing relative to MSY-based reference points. Because Pacific blue marlin is mainly caught as bycatch, direct control of the annual catch amount through the setting of a total allowable catch may be difficult.

Conservation Information

Since the stock is nearly full exploited, the ISC recommends that fishing mortality remain at or below current levels (2012-2014).

7.7 Pacific Bluefin Tuna

H. Nakano, PBFWG Chair, summarized the recommendations of the PBFWG (ISC/17/ANNEX/07/11/14) as it pertains to stock status and conservation advice.

Discussion

The stock assessment of PBF was not conducted in the past year. The PBFWG proposed some revisions to the stock status and conservation information narrative, which prompted considerable discussion by the Plenary. Some of the revisions were based on new information

that had come forward and was discussed in the past year but has not been thoroughly documented and reviewed by the PBFWG. While the utility of this information was not disputed, concerns were raised about the lack of rigorous documentation and the potential for the endorsement of this information.

The Plenary agreed to forward the stock status and conservation information statements adopted at ISC16 with slight clarifying modifications. In addition, the Plenary agreed to incorporate the summary results from the stakeholders workshop (ISC/17/ANNEX/11) and recent real-time monitoring observations from the 2015 and 2016 cohorts into the conservation information statement in this Plenary Report, because it represented new information discussed by the PBFWG that will be valuable for managers to consider. Management measures by RFMOs referred to here may be succeeded by updated measures.

Stock Status and Conservation Information

Stock Status

The PBFWG conducted a bench mark assessment (base-case model) that used the best available fisheries and biological information. The base-case model fits the data that were considered to be more reliable well and was internally consistent among most of the sources of data. The 2016 base-case model was a substantial improvement compared to the 2014 assessment and fits all reliable data well. The base-case model indicates: (1) spawning stock biomass (SSB) fluctuated throughout the assessment period (fishing years 1952-2014); (2) the SSB steadily declined from 1996 to 2010; and (3) the decline appears to have ceased since 2010, although the stock remains near the historic low. The model diagnostics suggest that the estimated biomass trend for the last 30 years is robust although SSB prior to the 1980s is uncertain due to data limitations.

Using the base-case model, the 2014 (terminal year) SSB was estimated to be around 17,000 t, which is about 9,000 t below the terminal year estimated in the 2014 assessment (26,000 in 2012). This was because of improvements to the input data and refinements to the assessment model scaled down the estimated value of SSB and not because the SSB declined from 2012 to 2014.

Recruitment estimates fluctuate widely without an apparent trend. The 2014 recruitment was relatively low, and the average recruitment for the last five years may have been below the historical average level. Note that recruitments in terminal years in an assessment are highly uncertain due to limited information on the cohorts. However, two of the last three data points from the Japanese troll CPUE-based index of recruitment, which was consistent with other data in the model, are at their lowest level since the start of the index (1980). Most age-specific fishing mortalities (F) for intermediate ages (2-10 years) are substantially above $F_{2002-2004}$ while those for age 0 as well as ages 11 and above are lower.

Although no limit reference points have been established for the PBF stock under the auspices of the WCPFC and IATTC, the F2011-2013 exceeds all calculated biological reference points except for F_{MED} and F_{LOSS} despite slight reductions to F in recent years. The ratio of SSB in 2014

relative to the theoretical unfished SSB ($SSB_{2014}/SSB_{F=0}$, the depletion ratio) is $2.6\%^5$ and $SSB_{2012}/SSB_{F=0}$ is 2.1% indicating a slight increase from 2012 to 2014. Although the $SSB_{2014}/SSB_{F=0}$ for this assessment (2.6%) is lower than $SSB_{2012}/SSB_{F=0}$ from the 2014 assessment (4.2%), this difference is due to improvements to the input data and model structure rather than a decline in SSB from 2012 to 2014. Note that potential effects on Fs as a result of the measures of the WCPFC and IATTC that started in 2015 or by other voluntary measures were not yet reflected in the data used in this assessment.

Since reference points for PBF have yet to be identified, two examples of Kobe plots are presented. These versions of the Kobe plot represent two interpretations of stock status in an effort to prompt further discussion. In summary, if these were the reference points, overfishing would be occurring or just at the threshold in the case of F_{MED}; and the stock would be considered overfished. Plot B in **ISC/16/ANNEX/09** shows that the stock has remained in an overfished and overfishing status for the vast majority of the assessment period if F20% and SSB20% are the reference points. The ISC notes that the SSB estimates before 1980 are more uncertain and that the reason why the fishing mortality is estimated to be so high right after the WWII is not well understood. The low biomass level at the beginning of the assessment period (1952) could potentially have been the result of relatively high catches prior to the assessment period.

Historically, the WPO coastal fisheries group has had the greatest impact on the PBF stock, but since about the early 1990s the WPO purse seine fleets, in particular those targeting small fish (age 0-1), have had a greater impact, and the effect of these fleets in 2014 was greater than any of the other fishery groups. The impact of the EPO fishery was large before the mid-1980s, decreasing significantly thereafter. The WPO longline fleet has had a limited effect on the stock throughout the analysis period. This is because the impact of a fishery on a stock depends on both the number and size of the fish caught by each fleet; i.e., catching a high number of smaller juvenile fish can have a greater impact on future spawning stock biomass than catching the same weight of larger mature fish.

Conservation Information

The steady decline in SSB from 1996 to 2010 appears to have ceased, although SSB₂₀₁₄ is near the historic low and the stock is experiencing exploitation rates above all calculated biological reference points except for F_{MED} and F_{LOSS} .

The projection results are based on the base-case model under several harvest and recruitment scenarios and time schedules. Under all examined scenarios the initial goal of WCPFC, rebuilding to SSB_{MED} by 2024 with at least 60% probability, is reached and the risk of SSB falling below SSB_{LOSS} at least once in 10 years was low.

⁴ "Unfished" refers to what SSB would be had there been no fishing.

⁵ The unfished SSB is estimated based upon equilibrium assumptions of no environmental or density-dependent effects.

The projection results show that the probability of SSB recovering to the initial WCPFC target (SSB_{MED} by 2024, 41,000 t, calculated in the same manner as the previous assessment) is 61.5% or above the level prescribed in the WCPFC CMM if low recruitment scenario is assumed and WCPFC CMM 2015-04 and IATTC Resolution C-14-06 continue in force and are fully implemented (Scenario 2 with low recruitment). Scenario 2 with low recruitment has the lowest prospect of recovery among the examined harvest scenarios. The probability of achieving the WCPFC's initial target (SSB_{MED} by 2024) would increase if more conservative management measures were implemented. The projection results indicate that a 10% reduction in the catch limit for fish smaller than the weight threshold in CMM 2015-04 would have a larger effect on recovery than a 10% reduction in the catch limit for fish larger than the weight threshold. The ISC notes that the current assessment model uses a maturity ogive that assumes 20%, 50% and 100% maturity in age 3 (weight on July 1: 34kg), 4 (weight on July 1: 58kg) and 5 (weight on July 1: 85kg), respectively, while the WCPFC CMM 2015-04 specifies that catches of fish smaller than 30kg should be reduced. The weight threshold in the CMM needs to be increased to 85kg (weight of age 5) if the intent is to reduce catches on all juveniles according to the maturity ogive in the assessment.

The projections results assuming a stronger stock-recruitment relationship (where h=0.9) than in the assessment model are not necessarily more pessimistic than the low recruitment scenario.

The projection results assume that the CMMs are fully implemented and are based on certain biological or other assumptions. In particular, the ISC noted the implementation of size based management measures need to be monitored carefully. If conditions change, the projection results would be more uncertain. Given the low SSB, the uncertainty in future recruitment, and the influence of recruitment has on stock biomass, monitoring recruitment and SSB should be strengthened so that the recruitment trends can be understood in a timely manner.

It should be noted that recent data (2016) on the abundance of large fish from Japanese and Taiwanese longline fisheries are higher than those of 2014 and 2015.

Japan reported that recent data from Japanese real-time monitoring on cohorts of 2015 and 2016 suggests that they are above that of 2014 (see **ISC/17/ANNEX/07**). While this information is encouraging, it needs to be vetted to assess its utility.

ISC conducted further projections on various harvest and recruitment scenarios in accordance with the requests of RFMOs and the result was presented to the ISC PBF Stakeholder Meeting in April, 2017 (ISC/17/ANNEX/11). Notable points from the additional analysis were:

- Different recruitment scenarios forecast entirely different levels of SSB in the future;
- Under average recruitment conditions, all harvest scenarios achieve the initial rebuilding target of SSB_{MED1952-2014} by 2024;
- Under all recruitment conditions with zero removals (no fishing), SSB trajectories achieved all rebuilding targets by approximately 2020 and the initial rebuilding target, SSB_{MED1952-2014}, within 2-3 years;
- Achieving 20% SSB0 during the projection period was not possible in most of the low recruitment scenarios;

- The probability of SSB falling below the historical lowest level at any time during the projection period is low (< 2%) in all projections;
- Scenarios that do not have catch limits for large fish in the EPO and WPO, or has a higher catch limit for large fish in WPO, do not achieve the initial rebuilding target, SSB_{MED1952-2014}, by 2024 under low recruitment conditions;
- Reducing the catch of small fish results in positive impacts on SSB trajectories, even with increases in the catch of large fish in the WPO.

8 REVIEW OF STOCK STATUS OF SECONDARY STOCKS IN THE WESTERN PACIFIC

8.1 **EPO**

M. Dreyfus provided an overview of the status of Secondary stocks (YFT, BET, and SKJ) in the IATTC Convention Area (ISC/17/INFODOC/02/03/04/08).

Catches of YFT are above 200,000 t, after record catches in 2002-2003 that were associated with very high recruitment. Most of the catch is taken by the purse seine fishery, especially in sets associated with dolphins. Nevertheless, the impact is more evenly distributed within the three types of sets in the purse seine fishery (dolphin associated, free swimming schools, and log sets), because of the size composition of the catch. Fish aggregating device (FAD) sets catch the smaller individuals.

Total BET catch is around 90,000 t. Most of the catch comes from the purse seine FAD fishery. Before 1993 it was basically a longline fishery. The FAD fishery has by far the biggest impact on this stock.

For those two stocks, IATTC periodically conducts full assessments and SSB has been close to the level that produces MSY in the recent past. In addition, SKJ condition is analyzed with relative reference points and based on productivity and susceptibility criteria compared to BET; IATTC scientific staff considers the SKJ biomass to be above the B_{MSY} level.

For 2017, after three meetings, the IATTC agreed to a quota for dolphin sets and FAD sets (including natural logs) was agreed. Nevertheless, new measures must be discussed for the following years in the upcoming IATTC annual meeting. Time closures, quotas, and FAD measures are among the options considered. There is concern about increasing capacity in the EPO, so IATTC staff recommended an increase from 62 to 72 days in the purse seine closure period.

Discussion

It was explained that the IATTC has been trying to control purse seine fishing capacity since 2003; despite adopted measures active capacity has continued to increase.

8.2 WCPO

A. Beeching presented on the SC12 review of stock assessments for SKJ and South Pacific BSH (ISC/17/INFODOC/01/05/06). The SKJ stock is in a good state, currently considered to be in

the vicinity of the interim target reference point. The South Pacific BSH assessment was considered to be preliminary and management advice was not drafted by SC12. This was partly due to the lack reference of points for this species and also the lack of quality historical data to support a stock assessment. The indicator analyses for the stocks for which stock assessments were not conducted (BET, YFT, and SPALB) were then briefly reviewed. The presentation concluded with a slide detailing upcoming WCPFC meetings.

Discussion

It was noted that a full stock assessment was conducted for the WCPO SKJ stock whereas IATTC evaluates various indicators without conducting a full assessment. It was also noted that there was a difference of opinion in the SC with respect to weighting of different data sources (CPUE versus tagging). The weighting scheme affects interpretation of stock status.

Although there is a 19% risk of SPALB falling below the LRP in 2033, the recommended catch reduction is aimed at maintaining economic viability of the fishery and is also expected to reduce the risk of breaching the LRP by 2033. It was further noted that the WCPFC has not yet adopted an acceptable risk level for breaching the SPALB LRP.

It was noted that it may be difficult to conduct a stock assessment for SP BSH using a fully integrated model because of data limitations.

9 REVIEW OF STATISTICS

9.1 STATWG Report

R.-F. Wu, the STATWG Chair, provided a summary of STATWG activities since ISC16 (ISC/17/ANNEX/06/16).

The STATWG Steering Group convened an intercessional meeting in La Jolla, California, USA, in January 18-20, 2017, as well as May 30, 2013. A meeting of the entire STATWG was convened in Vancouver, Canada, in July 7, 2017, prior to ISC17; two Information Papers and one Working Paper were submitted for this meeting.

It was announced that Izumi Yamasaki has relinquished her role as the ISC Data Administrator (DA) after eight years of outstanding service to the ISC. Kirara Nishikawa will assume the role as the ISC DA, as well as retain her current role as the ISC Webmaster.

The STATWG also welcomed new member Zane Zhang (Canada) who will replace John Holmes as the Data Correspondent for Canada.

It was reported that four of six items of the 2016 STATWG Work Plan were completed since ISC16; the two outstanding items were completed by July 13 at ISC17. The two items were updating member metadata annually, and resolving discrepancies of the data comparison report.

At the STATWG meeting in July 7, 2017 the following topics were presented and discussed:

- 1. Reviewed and resolved ISC database issues presented at ISC16
- 2. Reviewed the ISC catch tables

- 3. Compiled information on member's shark data collection systems
- 4. Reviewed the duties and Deliverables for the Data Administrator
- 5. Reviewed improvements to the ISC website
- 6. Conducted elections for Chair and Vice Chair of the STATWG for 2017-2018

Accomplishments of the STATWG since ISC16 include:

- 1. Continuing with publishing of data inventories on the ISC website
- 2. Continuing with improvements to the online data submission system and the ISC database, including updating the User's Manual for members
- 3. Continuing with archiving of stock assessment files from the species Working Groups from 2014-2017
- 4. Provided member metadata for updating annually
- 5. Provided comparison report on Member's re-submitted data vs data in the ISC database
- 6. Provided report on discrepancies between ISC catch tables vs catch tables in National Reports of Members
- 7. Compiled information on member's shark data collection systems to provide available years of data by shark species
- 8. Drafted an updated description of the duties and deliverables of the Database Administrator
- 9. Continuing with improvements and updates to the ISC website

The 2017 Work Plan for the STATWG was presented, as well as recommendations to the ISC17 Plenary.

The recommendations were:

- 1. The STATWG recommends that the Plenary approve the publishing of ISC Category II data online as public domain data, consistent with the practices of other RFMOs.
- 2. The STATWG recommends that the Plenary consider revising the current data submission deadline from July 1 to June 1, in order to provide sufficient time for the DA to validate Category II data prior to publishing online on the ISC website.
- 3. The STATWG recommends that the Plenary formally review the ISC Operations Manual to correct inconsistent and inaccurate information as it pertains to data submissions, archiving, and access.

The current national contacts list for the STATWG was also provided, as well as the 2017 Member Report Card. Ren-Fen Wu and Darryl Tagami were elected as Chair and Vice Chair of the STATWG, respectively, for a 1 year term. The STATWG Steering Group will schedule their next meeting in January, 2018, and requested the scheduling of a 2-day STATWG meeting prior to the ISC18 Plenary.

Discussion

Mexico noted that it has increased staff capacity, which will enable it to provide required data to the ISC by established deadlines.

The process for preparing and publicly disseminating Category II data was discussed at length. The STATWG has been working toward making these data available through the ISC website. It

was clarified that only Category II data from high-seas fisheries (longline, purse seine, pole and line, and troll) will be published. According to the ISC Operations Manual, Category II public domain data should be summarized monthly for all flag states combined at 5° x 5° latitude-longitude grid resolution for longline fisheries and 1° x 1° latitude-longitude grid resolution for surface fisheries. The STATWG proposed disaggregating data by flag but this is likely to present issues with respect to confidentiality rules, which prohibit publishing data coming from any spatio-temporal stratum in which fewer than three vessels are reporting. The alternatives of excluding data or aggregating data on a case-by-case basis were discussed. As a first step, the Plenary agreed that the STATWG should endeavor to make monthly Category II data available by ISC18 with fishery-level data aggregated by all countries at the appropriate resolution latitude-longitude grid resolution. Japan noted it may need approval from its relevant authorities if the definition of public domain data is changed from that in the ISC Operations Manual. It was noted that Japan already releases these data as part of the data sharing agreements with the IATTC and WCPFC.

The Plenary also discussed the STATWG's request to shift the data submission deadline from July 1 to June 1. The STATWG outlined the Database Administrator's workload and short time window to prepare all the data for Plenary given the current deadline. The Plenary concluded that the current July 1 deadline should stand although Members are encouraged to submit their data before the deadline if possible.

The Plenary endorsed the STATWG's proposal to convene a two-day meeting prior to ISC18.

9.2 Total catch tables

K. Nishikawa, the Database Administrator, presented the annual catch tables for ISC Member countries for countries for 2015-2016. The catch tables include the following ISC species of interest: albacore, blue shark, blue shark, Pacific bluefin tuna, striped marlin, swordfish, blue marlin, and shortfin make shark. Graphs of Graphs of the historical catch by country were also presented for each species. The catch tables were were generated from the ISC database, and are based on Category I data submitted by Data Correspondents Correspondents for the major fisheries in the North Pacific Ocean of the member countries. These catch These catch tables are included in the ISC Plenary Report as Table 15-1 to

Table 15-7 and serve as the official ISC catch tables.

Discussion

Canada noted an error in its reported NPALB landings for the period 1968-71. (During the meeting Canada subsequently updated its Category I data for the 1939-1993 period.) Canada will work with the Database Administrator to correct this error.

Mexico noted it is working on methods to provide more accurate estimates of its shark landings. It hopes to provide more accurate data in the future.

The omission of catch data in the ISC database for the historical Korean high seas driftnet fishery was flagged. This is especially an issue for billfish. Korea agreed to provide data from its high seas driftnet fisheries.

Japan noted that the lack of pre-1971 longline catch data in the ISC database for BUM is because the catch of both BUM and BLM were combined.

It was reiterated that WG input on data needs or issues is effected through WG chair membership on the STATWG.

10 REPORT OF THE SEMINAR

J. Holmes provided an overview of the ISC17 Seminar, "HMS Tagging and Lessons Learned" (ISC/17/ANNEX/15). David Itano, American Fishermen's Research Foundation, Honolulu, U.S.A., presented on "Tagging Tuna: Lessons learned, focusing on conventional large scale tuna tagging studies in the Pacific Ocean." Chuck Farwell, Monterey Bay Aquarium, U.S.A., then made a presentation on "Tagging Approaches for Highly Migratory Fish", which described four types of electronic tags (archival tags, acoustic tags, pop-up satellite archival tags, and SPOT tags) deployed by the aquarium as part of its research program. Finally, Dr. Erin Rechisky, Kintama Research, Nanaimo, Canada, presented on "Acoustic-tracking Juvenile Pacific Salmon in the Coastal Ocean: Results and Lessons Learned" in which she described tracking experiments of salmon smolts migrating down the Columbia and Fraser Rivers and north along the continental shelf.

Five types of tags and programs were discussed: dart tags, pop-up satellite archival tags, archival tags, acoustic tags, and acoustic tags with tracking. Each tag type provides useful information, but some (PSAT, archival) potentially provide richer datasets than others. Dart tags and archival tags are largely fisheries dependent while pop-up satellite tags, archival tags and acoustic tags with tracking are independent of fisheries for information return. When considering tagging studies/experiments, four themes emerged for future programs: (1) planning and experimental design are important to overall success as it links the project to specific assessment/management needs and develops a scientifically defensible tagging plan; (2) develop and enforce standardized tagging and release protocols to minimize differences between groups conducting tagging operations, and develop criteria for selecting fish to tag and identifying fish that should not be tagged; (3) data management is important, particularly with electronic tags since the amount of information provided by each tag is potentially large; and (4) publicize the program to the fishing industry (fishers, processors, buyers), the public, and other stakeholders before it begins and

provide feedback as the program progresses to ensure ongoing support (which can be translated into funding and long-term commitment) and the return of tags as necessary.

Discussion

The Plenary agreed to form an ad hoc working group to explore the development of a coordinated tagging program under the auspices of the ISC. It was agreed that it would be important for appropriate scientists, such as statisticians and stock assessment scientists, to participate in the ad hoc WG, so that any products proposed would be useful for enhancing stock assessments. The first task of this working group would be to inventory and describe existing tagging programs for HMS in the Pacific with the objective of reporting this information to ISC18. Given the commitment of Baja Aqua Farms to provide PBF in the EPO for tagging, the initial focus of the working group should be PBF. Based on further guidance from ISC18, the working group would then develop a framework for a possible ISC tagging program to be reviewed at ISC19. This framework would identify the scientific objectives to be achieved for each species under consideration, the likely costs for any such program, and possibly a prioritization of the species for which tagging programs would be supported. The outgoing ISC Chair agreed to work with the new Chair to organize this working group in the near future.

11 REVIEW OF MEETING SCHEDULE

11.1 Time and Place of ISC18

It was agreed that ISC18 will be held 11-16 July 2018 in the Republic of Korea.

11.2 Time and place of Working Group intercessional meetings

A draft schedule of proposed intersessional meetings was reviewed and amended, see Table 11-1.

Table 11-1. Schedule of ISC working group and Plenary meetings.

Date	Meeting	Contact
	2017	
	BILLWG	
	SWO Data Prep & Modeling (SS3	Jon Brodziak
Sept	version)- Webinar	jon.brodziak@noaa.gov
		Mikihiko Kai (Acting
	SHARKWG	Chair)
Sept	Mako Ageing- Webinar	kaim@affrc.go.jp
	ALBWG	
	MSE (unconfirmed)- Vancouver,	Hidetada Kiyofuji
Oct	Canada	hkiyofuji@affrc.go.jp
		Mikihiko Kai (Acting
	SHARKWG	Chair)
Oct 19-24	Mako ageing workshop- Japan	kaim@affrc.go.jp

	BILLWG	
	SWO Data Prep & Modeling (Data	Jon Brodziak
Nov	Plan)- Webinar	jon.brodziak@noaa.gov
		Mikihiko Kai (Acting
	SHARKWG	Chair)
Nov	Mako Data Prep-Webinar	kaim@affrc.go.jp
1101	Mario 2 marriop (Vocamar	Mikihiko Kai (Acting
Nov 27-Dec	SHARKWG	Chair)
1	Mako Data Prep Meeting- Japan	kaim@affrc.go.jp
	· · · · · · · · · · · · · · · · · · ·	kann@amc.go.jp
Dec 3-8	WCPFC	
	BILLWG	
	SWO Data Prep & Modeling (Data	Jon Brodziak
Dec	Process)- Webinar	jon.brodziak@noaa.gov
Dec	<i>'</i>	Jon.brodziak@noaa.gov
	2018	
	BILLWG	Jon Brodziak
Jan 17-23	SWO Data Prep- Honolulu, HI, USA	jon.brodziak@noaa.gov
	STATWG	<u> </u>
	Steering Group Meeting- Shimizu,	Ren-Fen Wu
Jan 23-25	Japan	fan@ofdc.org.tw
3dii 23 23	Japan	Mikihiko Kai (Acting
	SHARKWG	, ,
T /IC 1		Chair)
Jan/Feb	Data Prep- Webinar	kaim@affrc.go.jp
	PBFWG	
	Update Stock Assessment- La Jolla,	Hideki Nakano
Mar 5-12	CA, USA	hnakano@affrc.go.jp
		<u> </u>
	SHARKWG	Mikihiko Kai (Acting
	Mako Assessment Meeting- Location	Chair)
Mar/Apr	TBD	kaim@affrc.go.jp
	ALBWG	Hidetada Kiyofuji
Apr	Workshop- Location TBD	hkiyofuji@affrc.go.jp
	Workshop Location 1DD	inkryoruji e urii e.go.jp
	BILLWG	
	SWO Assessment Modeling Workshop-	Jon Brodziak
Apr/May	Location TBD	jon.brodziak@noaa.gov
	ALBWG	Hidetada Kiyofuji
Jul	1 Day Pre-Plenary Meeting- Korea	hkiyofuji@affrc.go.jp
	•	
_	PBFWG	Hideki Nakano
Jul	1 Day Pre-Plenary Meeting- Korea	hnakano@affrc.go.jp
	BILLWG	Jon Brodziak
Jul	1 Day Pre-Plenary Meeting- Korea	jon.brodziak@noaa.gov

Jul	SHARKWG 1 Day Pre-Plenary Meeting- Korea	Mikihiko Kai (Acting Chair) kaim@affrc.go.jp
Jul	STATWG 2 Days Pre-Plenary Meeting- Korea	Ren-Fen Wu fan@ofdc.org.tw
Jul 11-18	ISC18 Plenary- Korea	

12 ADMINISTRATIVE MATTERS

12.1 Formalization of ISC

Plenary discussed formalization of ISC. ISC currently does not operate under a formal agreement or Memorandum of Understanding (MoU). Concern for this arrangement was noted several Plenary sessions ago. At the time, it was noted that ISC Members may be able to better participate in ISC scientific working groups if a formal agreement existed. A formal agreement may also allow the creation of a budget as well as a formal secretariat that would support ISC operations. ISC Members were asked to review a draft MoU as a model to share with the people in their governments that would be responsible for furthering a formal multi-country agreement. ISC Members were also asked to send to the ISC Chair by the end of ISC17 the name of the person in their governments that might be responsible for addressing this issue.

Discussion

Members agreed with the requests of the ISC Chair to provide the MoU to the appropriate contact person in their respective governments for review and provide him with the names of these points of contact by the close of ISC17. Finally, he emphasized the need to conclude this process as quickly as possible.

12.2 Peer Review of Function and Process for Stock Assessments

ISC is required to conduct a peer review of its function and process every five years. At ISC16, it was agreed to solicit a panel to recommend an independent peer-review process for ISC. Independent peer review is critical to rigorous science and a clear process is an issue faced by many science providers that provide stock assessments and other science products to tuna RFMOs. At ISC16, the U.S.A. and Japan agreed to solicit funding for one reviewer each. A Member country to support a third reviewer was not identified. Draft terms of reference (TOR) were developed and made available for review (ISC/17/PLENARY/12).

Discussion

The ToR were agreed to but it was noted that they should be updated with regard to the dates and locations of meeting that reviewers may attend. It was agreed that one reviewer would observe the benchmark SWO stock assessment and the other would observe the PBF update assessment. If a third reviewer is available, then the SMA benchmark assessment should be observed assuming an assessment is conducted.

It was noted that reviewers can be selected from government agencies as long as they have no direct involvement in the ISC or other Pacific tuna RFMOs.


The outgoing Chair will continue working with Members to solicit a third reviewer, if at all possible.

12.3 Work Group Election Results

Hidetada Kiyofugi (Japan) was elected Chair of the ALBWG and Steve Teo (USA) Vice Chair. Jon Brodziak (USA) was reelected as Chair of the BILLWG and Hirotaka Ijima (Japan) as Vice Chair. Mikihiko Kai (Japan) was elected as Vice Chair of the SHARKWG and will serve as Chair until the November 2017 SHARKWG Workshop when the election for Chair is scheduled. Ren-fan Wu (Chinese Taipei) was reelected as Chair of the STATSWG and Darryl Tagami (USA) as Vice Chair. All will serve in this capacity for three years, spanning 2017-2020.

12.4 Election of the Chair and Vice Chair

Dr. John Holmes, Canada, was elected as ISC Chair and Dr. Eric Chang, Chinese Taipei, was elected as the ISC Vice Chair for three-year terms (2017-2020). **Organizational Chart and Contact Persons**



Working Group Key:

1 Canada 2 China 3 Chinese-Taipei 4 Korea 5 Japan 6 Mexico 7 USA 8 PICES 9 SPC 10 IATTC 11 FAO 12 WCPFC VC Vice Chair DM Database Manager

This is not a comprehensive list but the main points of contact.

12.5 Other Business

12.5.1 Template for Stock Status and Conservation Advice

At ISC15, Plenary agreed to develop a template for WGs to present information on stock status and conservation advice in order to produce greater consistency across stock assessments and the scientific information ISC provides and to facilitate Plenary deliberations. The U.S.A. provided a draft template (ISC/17/PLENARY/13) for ISC to consider adopting noting that as a science organization, ISC documents should be results-focused.

Discussion

It was noted that the proposed template was used as a guide for presenting stock status and conservation advice for the new NPALB and BSH assessments. More generally, the way in which the framework can be adapted to the specifics of a particular stock assessment was discussed. The template is proposed as a minimum set of contents for such reporting while striving to be flexible enough to accommodate different approaches and needs of WGs. The purpose is to agree upon some level of consistency in the way in which ISC stock assessment results are reported.

The U.S.A. reported that in their search they could not find any other tuna RFMO that had formally adopted such a detailed template (WCPFC, IATTC, IOTC, ICCAT). But it was noted that the RFMOs tend to have consistent formats among their own reports. The U.S. noted that the terms "overfished" and "overfishing" have specific legal definitions within its domestic policy framework related to domestically required biological reference points. Plenary discussions provided many useful comments for improving the template, all of which were incorporated into the revised version of the template (ISC/17/PLENARY/13).

After considerable discussion and input from the Members, it was agreed that further review is needed before the template can be adopted. An important aspect of this further review is to involve fishery managers since the objective is to guide the production of consistent and concise statements about stock status and conservation information for managers. To this end it was agreed that the revised template should be presented to the NC and Members should also discuss the template with fishery managers in their respective governments.

Although not formally adopted, working groups were encouraged to consult the draft template when formulating stock status and conservation information in future stock assessment reports.

12.5.2 Candidate Reference Points

The Chair drew the Plenary's attention to **ISC/10/PLENARY/04**, Biological Reference Points for Tuna and Tuna-Like Species in the North Pacific Ocean. He recommended that this document be revisited and updated. The need for timely advice on candidate reference points is only growing as the WCPFC develops harvest strategies for HMS stocks under its purview. The ISC should be prepared to respond. An update of this document, both to narrow the suite of candidates and evaluate candidates based on the latest information, would serve that purpose.

13 ADOPTION OF REPORT

The draft report of the meeting was adopted and will be forwarded to the WCPFC-SC. The ISC Chair will finalized the plenary report for posting on the ISC website.

14 CLOSE OF MEETING

G. DiNardo thanked the ISC Membership for their support and commitment over the past year and throughout his 7 year tenure as ISC Chair. The scientific structure and competency of ISC has increased significantly, owing to the dedication and support of its scientists. I step down from the leadership role knowing that the organization is strong and in capable hands. The meeting was closed at 2:15 PM on 17 July 2017.

15 CATCH TABLES

Table 15-1. North Pacific albacore catches (in metric tons) by fisheries, 1952-2012. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional.

		CA	AN				JPN					K()R	MEX		
Catch dispositi on	Year	Troll	CAN Total	Set-net	Drift gill- net	Longline	Pole and line	Troll	Others	Purse seine	JPN Total	Longline	KOR Total	Others	Purse seine	MEX Total
Retain ca	1936															
	1937															
	1938															
	1939	1,290	1,290													
	1940	20	20													
	1941	350	350													
	1942															
	1943	130	130													
	1944	2,100	2,100													
	1945	6,480	6,480													I
	1946	1,960	1,960													I
	1947	360	360													I
	1948	9,840	9,840													
	1949	10,120	10,120													
	1950	9,610	9,610													
	1951	860	860													1
	1952	710	710	55	-	26,687	41,787	-	237	154	68,920			-	-	-
	1953	50	50	88	-	27,777	32,921	-	132	38	60,956			-	_	-
	1954			6	-	20,958	28,069	_	38	23	49,094			-	_	-
	1955			28	-	16,277	24,236	-	136	8	40,685			-	_	-
	1956	170	170	23	-	14,341	42,810	_	57		57,231			-	_	-
	1957	80	80	13	-	21,053	49,500	-	151	83	70,800			_	_	_
	1958	740	740	38	-	18,432	22,175	-	124	8	40,777			_	_	_
	1959	2,120	2,120	48	_	15,802	14,252	_	67		30,169			_	_	_
	1960	50	50	23	-	17,369	25,156	_	76		42,624			_	_	-
	1961	40	40	111	_	17,437	18,639	-	268	7	36,462			39	2	41
	1962	10	10	20	-	15,764	8,729	-	191	53	24,757			0	0	0
	1963	50	50	4	-	13,464	26,420	-	218	59	40,165			0	31	31
	1964	30	30	50	-	15,458	23,858	-	319	128	39,813			_	0	_
	1965	150	150	70	_	13,701	41,491	_	121	11	55,394			_	0	_
	1966	440	440	64		25,050	22,830	_	585	111	48,640			_	0	_
	1967	1,610	1,610	43	-	28,869	30,481	_	520	89	60,002			_		_
	1968	10,280	10,280	58		23,961	16,597	_	1,109	267	41,992			_	_	_
	1969	13,650	13,650	34		18,006	31,912	_	925	521	51,398			_	0	_
	1970	3,900	3,900	19		16,222	24,263	_	498	317	41,319			_	0	
	1971	17,460	17,460	5		11,473	52,957		354	902	65,691	0	0		0	
	1972	39,210	39,210	6	1	13,022	60,569	_	638	277	74,513	0	0	0	100	100
	1973	14,000	14,000	44	39	16,760	68,767	_		1,353	87,449	4	4		0	100
	1973	13,310	13,310	13	224	13,384	73,564	_	891	1,333	88,237	91	91	0	1	1
	1974	1,110	1,110	13	166	10,303	52,152	_	230	159	63,023	7,050	7,050	0	1	1
	1976	2,780	2,780	15	1,070	15,812	85,336		270	1,109	103,612	2,212	2,212	5	36	41
	1976	530	530	5	688	15,681	31,934	-	365	669	49,342	500	500	0	30	3
	1977	230	230	21	4,029	13,007	59,877	-	2,073	1,115	80,122	669	669	0	1	1
	1978	5,210	5,210	16	2,856	14,186	44,662	-	1,139	1,113	62,984	009	009	0	1	1
	1979	2,120	2,120	10	2,836	14,180	46,742	-	1,139	329	65,925	592	592	0	31	31
	1980	2,000	2,000	8	10,348	17,878	27,426	-	699	252	56,611	0	0	0	8	8
	1981	1,040	1,040	11	10,348	16,714	29,614	-		561	59,893	4,874	4,874	0	0	0
	1982	2,250	2,250	22	6,852	15,094	21,098	_	99	350	43,515	366	366	0	0	0
	1983	500	500	24				-	494	3,380	53,952	1,925	1,925	6	107	113
	1984	560	560	68	8,988 11,204	15,053 14,249	26,013 20,714	-	339	1,533	53,952 48,107	2,789	2,789	35	107	49

2006 2007 2008 2009 2010 2011	5,833 6,040 5,464 5,693 6,527 5,385	5,833 6,040 5,464 5,693 6,527 5,385	55 30 101 33 42 50	221 226 1,531 149 24	21,168 22,381 19,092 21,995 21,167 20,956	15,400 37,768 19,060 31,172 19,561 25,704	460 519 549 410 588 443	42 44 34 43 37 78	364 5,682 825 2,076 330 480	37,710 66,650 41,192 55,878 41,749 47,723	134 136 400 95 107	134 136 400 95 107	0 0 -	109 40 10 17 25	0 109 40 10 17 25
2006 2007 2008 2009	5,833 6,040 5,464 5,693	5,833 6,040 5,464 5,693	55 30 101 33	221 226 1,531 149	21,168 22,381 19,092 21,995	15,400 37,768 19,060 31,172	460 519 549 410	42 44 34 43	364 5,682 825 2,076	37,710 66,650 41,192 55,878	134 136 400 95	134 136 400 95	0	109 40 10 17	109 40 10 17
2006 2007 2008	5,833 6,040 5,464	5,833 6,040 5,464	55 30 101	221 226 1,531	21,168 22,381 19,092	15,400 37,768 19,060	460 519 549	42 44 34	364 5,682 825	37,710 66,650 41,192	134 136 400	134 136 400	0	109 40 10	109 40 10
2006 2007	5,833 6,040	5,833 6,040	55 30	221 226	21,168 22,381	15,400 37,768	460 519	42 44	364 5,682	37,710 66,650	134 136	134 136	0	109 40	109 40
2006	5,833	5,833	55	221	21,168	15,400	460	42	364	37,710	134	134	0	109	109
										,	-				-
2005	4,829	4,829	07	154	20.465	16 122	665	224	950	38,598	419	410	0	0	
	.,	.,	30	61	-		772	54	.,	57,713	77	77	0	104	104
													0		29
										76,655	111	111			28
2001	5,248	5,248		117	28,801	29,430	416			59,851		63		0	18
			136				645								103
										,				0	57
1998	4,177	4,177	41	206	35,755	22,926	1,190	104	1,554	61,776	225	225	0	8	8
1997	2,166	2,166	40	359	38,899	32,238	1,585	135	1,068	74,324	404	404	0	53	53
1996	3,321	3,321	43	116	32,440	20,272	815	127	581	54,394	157	157	0	21	21
1995	1,761	1,761	28	282	29,050	20,981	856	78	1,177	52,452	13	13	0	5	5
1994	1,998	1,998	11	263	29,565	26,389	-	823	2,026	59,077	2	2	0	6	6
1993	4,940	4,940	3	287	-		-	897	2,889	46,806	2	2	0	11	11
1992	3,630	3,630	12	2,721	19,042	13,888	-		4,104	41,289	1	1	0	10	10
1991	1,390	1,390	4	3,401	17,039	7,103	-	395	2,652	30,594	4	4	0	2	2
1990	3,020	3,020	5	6,064	15,785	8,532	-	248	1,995	32,629	29	29	0	2	2
1989	1,400	1,400	33	7,437	13,031	8,629	-	433	2,521	32,084	561	561	0	2	2
	,	,			-		_			,-	, .	, -	0	15	15
							-						0	7	7
	1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000	1987 1,040 1988 1,550 1989 1,400 1990 3,020 1991 1,390 1992 3,630 1993 4,940 1994 1,998 1995 1,761 1996 3,321 1997 2,166 1998 4,177 1999 2,734 2000 4,531 2001 5,248 2002 5,379 2003 6,847 2004 7,857	1987 1,040 1,040 1988 1,550 1,550 1989 1,400 1,400 1990 3,020 3,020 1991 1,390 1,390 1992 3,630 3,630 1993 4,940 4,940 1994 1,998 1,998 1995 1,761 1,761 1996 3,321 3,321 1997 2,166 2,166 1998 4,177 4,177 1999 2,734 2,734 2000 4,531 4,531 2001 5,248 5,248 2002 5,379 5,379 2003 6,847 6,847 2004 7,857 7,857	1987 1,040 1,040 16 1988 1,550 1,550 7 1989 1,400 1,400 33 1990 3,020 3,020 5 1991 1,390 1,390 4 1992 3,630 3,630 12 1993 4,940 4,940 3 1994 1,998 1,998 11 1995 1,761 1,761 28 1996 3,321 3321 43 1997 2,166 2,166 40 1998 4,177 4,177 41 1999 2,734 2,734 90 2000 4,531 4,531 136 2001 5,248 5,248 78 2002 5,379 5,379 109 2004 7,857 7,857 30	1987 1,040 1,040 16 6,698 1988 1,550 1,550 7 9,074 1989 1,400 1,400 33 7,437 1990 3,020 3,020 5 6,064 1991 1,390 1,390 4 3,401 1992 3,630 3,630 12 2,721 1993 4,940 4,940 3 287 1994 1,998 1,998 11 263 1995 1,761 1,761 28 282 1996 3,321 3,321 43 111 1997 2,166 2,166 40 359 1998 4,177 4,177 41 206 1999 2,734 2,734 90 289 2000 4,531 4,531 136 67 2001 5,248 5,248 78 117 2002 5,379 5,379 109 332 <td>1987 1,040 1,040 16 6,698 14,668 1988 1,550 1,550 7 9,074 14,688 1989 1,400 1,400 33 7,437 13,031 1990 3,020 3,020 5 6,064 15,785 1991 1,390 1,390 4 3,401 17,039 1992 3,630 3,630 12 2,721 19,042 1993 4,940 4,940 3 287 29,933 1994 1,998 1,998 11 263 29,565 1995 1,761 1,761 28 282 29,050 1996 3,321 3,321 43 116 32,440 1997 2,166 2,166 40 359 38,899 1998 4,177 4,177 41 206 35,755 1999 2,734 2,734 90 289 33,339 2000 4,531</td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 1988 1,550 1,550 7 9,074 14,688 6,216 1989 1,400 1,400 33 7,437 13,031 8,629 1990 3,020 3,020 5 6,064 15,785 8,532 1991 1,390 1,390 4 3,401 17,039 7,103 1992 3,630 3,630 12 2,721 19,042 13,888 1993 4,940 4,940 3 287 29,933 12,797 1994 1,998 1,761 28 282 29,050 20,981 1996 3,321 3,321 43 116 32,440 20,272 1997 2,166 2,166 40 359 38,899 32,238 1998 4,177 4,177 41 206 35,755 22,926 1999 2,734 2,734 90</td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 - 1988 1,550 1,550 7 9,074 14,688 6,216 - 1989 1,400 1,400 33 7,437 13,031 8,629 - 1990 3,020 3,020 5 6,064 15,785 8,532 - 1991 1,390 1,390 4 3,401 17,039 7,103 - 1992 3,630 3,630 12 2,721 19,042 13,888 - 1993 4,940 4,940 3 287 29,933 12,797 - 1994 1,998 1,998 11 263 29,565 26,389 - 1995 1,761 1,761 28 282 29,050 20,981 856 1996 3,321 3,321 43 116 32,440 20,272 815 1997 2,166 40</td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 1993 4,940 4,940 3 287 29,933 12,797 - 897 1994 1,998 1,998 11 263 29,565 26,389 - 823 1995 3,321 3,321 43 116 32,440 20,272 815 127 1997 2,166 2,166 40<</td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 1994 1,998 1,998 11 263 29,565 26,389 - 823 2,026 1995 1,761 1,761 28 282</td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 44,842 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 46,806 1994 1,998 1,998 11 263 29,565 26,389 -</td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 29 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 46,806 2 1994 1,998</td> <td> 1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1,624 1,988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 799 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 561 561 590 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 29 29 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 44 4992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1 1 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 46,806 2 2 2 1994 1,998 1,998 11 263 29,565 26,389 - 823 2,026 59,077 2 2 2 1995 1,761 1,761 28 282 29,050 20,981 856 78 1,177 52,452 13 13 1396 3,321 3,321 43 116 32,440 20,272 815 127 581 54,394 157 157 1997 2,166 2,166 40 359 38,899 32,238 1,585 135 1,068 74,324 404 404 404 1998 4,177 4,177 41 206 35,755 22,926 1,190 104 1,554 61,776 225 225 1999 2,734 2,734 90 289 33,339 50,369 891 62 6,872 91,912 98 98 2000 4,531 4,531 136 67 29,995 21,550 645 86 2,408 54,887 15 15 2001 5,248 5,248 78 117 28,801 29,430 416 35 974 59,851 63 63 2002 5,379 5,379 109 332 23,585 48,454 787 85 3,303 76,655 111 111 2003 6,847 6,847 69 126 20,907 36,114 922 85 627 58,850 146 146 2004 7,857 7,857 30 61 17,341 33,255 772 54 7,200 57,713 77 77 </td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1,624 0 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 799 0 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 561 0 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 29 29 29 0 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 4 0 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1 0 1993 4,940 4,940</td> <td>1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1,624 0 7 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 799 0 15 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 561 0 2 1990 3,020 3,020 5 6,604 15,785 8,532 - 248 1,995 32,629 29 29 0 2 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 4 0 2 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1 0 0</td>	1987 1,040 1,040 16 6,698 14,668 1988 1,550 1,550 7 9,074 14,688 1989 1,400 1,400 33 7,437 13,031 1990 3,020 3,020 5 6,064 15,785 1991 1,390 1,390 4 3,401 17,039 1992 3,630 3,630 12 2,721 19,042 1993 4,940 4,940 3 287 29,933 1994 1,998 1,998 11 263 29,565 1995 1,761 1,761 28 282 29,050 1996 3,321 3,321 43 116 32,440 1997 2,166 2,166 40 359 38,899 1998 4,177 4,177 41 206 35,755 1999 2,734 2,734 90 289 33,339 2000 4,531	1987 1,040 1,040 16 6,698 14,668 19,082 1988 1,550 1,550 7 9,074 14,688 6,216 1989 1,400 1,400 33 7,437 13,031 8,629 1990 3,020 3,020 5 6,064 15,785 8,532 1991 1,390 1,390 4 3,401 17,039 7,103 1992 3,630 3,630 12 2,721 19,042 13,888 1993 4,940 4,940 3 287 29,933 12,797 1994 1,998 1,761 28 282 29,050 20,981 1996 3,321 3,321 43 116 32,440 20,272 1997 2,166 2,166 40 359 38,899 32,238 1998 4,177 4,177 41 206 35,755 22,926 1999 2,734 2,734 90	1987 1,040 1,040 16 6,698 14,668 19,082 - 1988 1,550 1,550 7 9,074 14,688 6,216 - 1989 1,400 1,400 33 7,437 13,031 8,629 - 1990 3,020 3,020 5 6,064 15,785 8,532 - 1991 1,390 1,390 4 3,401 17,039 7,103 - 1992 3,630 3,630 12 2,721 19,042 13,888 - 1993 4,940 4,940 3 287 29,933 12,797 - 1994 1,998 1,998 11 263 29,565 26,389 - 1995 1,761 1,761 28 282 29,050 20,981 856 1996 3,321 3,321 43 116 32,440 20,272 815 1997 2,166 40	1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 1993 4,940 4,940 3 287 29,933 12,797 - 897 1994 1,998 1,998 11 263 29,565 26,389 - 823 1995 3,321 3,321 43 116 32,440 20,272 815 127 1997 2,166 2,166 40<	1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 1994 1,998 1,998 11 263 29,565 26,389 - 823 2,026 1995 1,761 1,761 28 282	1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 44,842 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 46,806 1994 1,998 1,998 11 263 29,565 26,389 -	1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 29 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 46,806 2 1994 1,998	1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1,624 1,988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 799 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 561 561 590 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 29 29 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 44 4992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1 1 1993 4,940 4,940 3 287 29,933 12,797 - 897 2,889 46,806 2 2 2 1994 1,998 1,998 11 263 29,565 26,389 - 823 2,026 59,077 2 2 2 1995 1,761 1,761 28 282 29,050 20,981 856 78 1,177 52,452 13 13 1396 3,321 3,321 43 116 32,440 20,272 815 127 581 54,394 157 157 1997 2,166 2,166 40 359 38,899 32,238 1,585 135 1,068 74,324 404 404 404 1998 4,177 4,177 41 206 35,755 22,926 1,190 104 1,554 61,776 225 225 1999 2,734 2,734 90 289 33,339 50,369 891 62 6,872 91,912 98 98 2000 4,531 4,531 136 67 29,995 21,550 645 86 2,408 54,887 15 15 2001 5,248 5,248 78 117 28,801 29,430 416 35 974 59,851 63 63 2002 5,379 5,379 109 332 23,585 48,454 787 85 3,303 76,655 111 111 2003 6,847 6,847 69 126 20,907 36,114 922 85 627 58,850 146 146 2004 7,857 7,857 30 61 17,341 33,255 772 54 7,200 57,713 77 77	1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1,624 0 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 799 0 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 561 0 1990 3,020 3,020 5 6,064 15,785 8,532 - 248 1,995 32,629 29 29 29 0 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 4 0 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1 0 1993 4,940 4,940	1987 1,040 1,040 16 6,698 14,668 19,082 - 173 1,205 41,842 1,624 1,624 0 7 1988 1,550 1,550 7 9,074 14,688 6,216 - 170 1,208 31,363 799 799 0 15 1989 1,400 1,400 33 7,437 13,031 8,629 - 433 2,521 32,084 561 561 0 2 1990 3,020 3,020 5 6,604 15,785 8,532 - 248 1,995 32,629 29 29 0 2 1991 1,390 1,390 4 3,401 17,039 7,103 - 395 2,652 30,594 4 4 0 2 1992 3,630 3,630 12 2,721 19,042 13,888 - 1,522 4,104 41,289 1 1 0 0

Table 15-1. Continued

		TV	VN							USA					l
Set-net	Gill-net (not specified	Longline	Others	Purse seine	TWN Total	Drift gill- ne t	Handline	Longline	Pole and line	Troll	Others	Purse seine	Sport	USA Total	Total
										442				442	442
										1,681				1,681	1,681
										8,594				8,594	8,594
										8,586				8,586	9,876
										6,603				6,603	6,623
										5,412				5,412	5,762
										10,678				10,678	10,678
										17,071				17,071	17,201
										23,957				23,957	26,057
										17,886				17,886	24,366
										10,955				10,955	12,915
										12,235				12,235	12,595
						1		45		22,457				22,502	32,342
								33		24,901				24,934	35,054
								27		32,746				32,773	42,383
								24		15,629				15,653	16,513
								46		23,843			1,373	25,262	94,892
								23		15,740			171	15,934	76,940
								13		12,246			147	12,406	61,500
								9		13,264			577	13,850	54,535
								6		18,751			482	19,239	76,640
								4		21,165			304	21,473	92,353
								7		14,855			48	14,910	56,427
								5		20,990				20,995	53,284
								4		20,100			557	20,661	63,335
								5	2,837	12,055	1		1,355	16,253	52,796
								7	1,085	19,752	1		1,681	22,526	47,293
								7	2,432	25,140	1		1,161	28,740	68,986
								4	3,411	18,388			824	22,627	62,470
								3	417	16,542	1		731	17,694	73,238
								8	1,600	15,333	1		588	17,529	66,609
		330	189		519			12	4,113	17,814			707	22,646	84,777
	_	216	283		499			11	4,906	20,434			951	26,302	79,073
	_	65	423		488			14	2,996	18,827			358	22,195	87,731
		34	59		93			9	4,416	21,032			822	26,279	71,591
	_	20	52		72			11	2,071	20,526			1,175	23,783	107,006
	-	187	32		187	1		8	3,750	23,600			637	23,783	142,005
	_	10/	-		10/	1		14	2,236	15,653			84	17,987	119,440
	_	486	-		486			9	4,777	20,178			94	25,058	127,183
	-	1,240	-		1,240	-		33	3,243	18,932	10		640	22,858	95,282
	_	686	-		686			23	2,700	15,905	4		713	19,345	128,676
	_		-		572			37	1,497	9,969	4		537	19,345	62,987
	-	572						54			15		810		99,470
-	-	6 81	-		6 81			34	950 303	16,613	15		74	18,442 7,158	75,434
	1	249	20		270	-			303	6,781			168	7,158 8,106	75,434
- 1						-		25		7,556					
1	-	143	12		156	-		25	748	12,637	21		195	13,605	72,380
-	-	38	9		47	-		105	425	6,609	21		257	7,417	73,271
-	-	8	1		9			6	607	9,359		2.720	87	10,059	56,199
	1	-	_		1	ļ .		2	1,030	9,304		3,728	1,427	15,491	71,982
1	-	-	2		3	2				6,422	118	26	1,176	7,744	59,252

(15)	(46,226)	(120,202)	(1,091)		(167,546)	800	797	17,370	52,932	1,072,843	1,003	4,197	47,022	1,196,964	5,179,06
															2
(10)	(20,227)	(120,202)	(1,077)	(0)	(107540)	000	121	27,570	52,752	2,072,073	1,003	7,177	17,022	1,170,704	3,177,04
- ' '		(120,202)	(1,099)	(0)	(167546)	800	797	17,370	52,932	1,072,843	1,003	4,197	47,022	1,196,964	5,179,04
(1)	(2)	(3406)	(4)	(0)	(3,413)	0.8	31	250		10,687	7		675	11,650	53,54
1	2	3,020	4	0	(3,027)	0.5	62	243		11,560	2		926	12,794	64,2
1	1	2,617	0	0	2,619	0	49	208		13,369	0		1,045	14,671	74,0
1	0	4,427	0	0	4,428	5	46	317		12,312	0		839	13,519	79,1
2	2	2,643	0	0	2,647	8	253	660		14,152	2		1,212	16,287	83,1
+	1	2,818 3,434	2	0	3,437	5	84	708		10,147	0		421	11,365	67,9
-	-	2,378	-	-	2,378	5	53	421		12,343	0	31	862	13,622	64,2
-	-	3,069 2,378	-	-	3,069 2,378	1 4	28 97	203		11,762 12,343	+	31	944	12,563 13,622	62,6 77,6
-	+	2,916	-	-	2,916	4	94 28	250 354		11,911	+	77	461 418	12,797	88,5
-	-	4,317	-	-	4,317	3	0.	270		12,642	+		385	13,300	61,4
-	-	4,472	-	-	4,472	20		296		8,654	1		1,719	10,690	59,0
-	-	4,988	-	-	4,988	12		361		13,630	3	1	1,506	15,513	86,2
-	-	7,166	-	-	7,166	16		524		14,246	8	44	2,214	17,052	90,0
-	-	7,965	-	-	7,965	30		525		11,003	13	4	2,357	13,932	104,0
-	1	8,684	+	-	8,685	94		1,295		11,543	39	51	1,635	14,657	88,
-	-	8,842	-	-	8,842	55		940		9,834	24	4	1,798	12,655	81,
-	-	8,393	-	-	8,393	149		1,542		10,451	61	48	3,621	15,872	119,0
-	-	8,810	-	-	8,810	80		1,120		14,577	15	33	1,208	17,033	92,0
-	-	9,456	-	-	9,456	60		1,653		14,458	12	2	1,018	17,203	103,0
-	-	7,596	-	-	7,596	83		1,185		17,150	10	11	88	18,527	84,0
-	-	4,280	-		4,280	52		882		8,302	4		102	9,342	67,8
-	-	83	-		83	38		544		11,068	66		106	11,822	72,9
-	-	5	-		5			438		6,325	194		25	6,982	58,
-	7,866	-	_		7,866			334		4,626	40		2	5,002	57,
-	3,398	12	37		3,410	17		312		1,917	20	/1	6	2,272	37,
-	16,701	40	39		16,744	29		177		2,733	20	71	24	3,054	55,4
6	7,389 8,350	40	-		7,395 8,390	15		307 248		4,221 1,896	76 10	17	64 160	4,700 2,319	45,3 44,7
2	2,514	-	-		2,516	5		150		2,772	139	1	74	3,141	50,1
-						_									= o .

Table 15-2. Pacific bluefin tuna catches (in metric tons) by fisheries, 1952-2012. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional.

					JPN						KC)R			MEX		
Catch disposition	Year	Set-net	Longline ¹	Pole and line	Troll ²	Others	Purse seine	JPN Total	Set-net	Longline	Purse seine	Trawl	Troll	KOR Total ³	Others	Purse seine	MEX Total
Retain catch	1952	2,145	2,694	2,198	667	1,700	7,680	17,084							-	-	-
	1953	2,335	3,040	3,052	1,472	160	5,570	15,629							-	-	-
	1954	5,579	3,088	3,044	1,656	266	5,366	18,999							-	-	-
	1955	3,256	2,951	2,841	1,507	1,151	14,016	25,722							-	-	-
	1956	4,170	2,672	4,060	1,763	385	20,979	34,029							-	-	-
	1957	2,822	1,685	1,795	2,392	414	18,147	27,255							-	-	-
	1958	1,187	818	2,337	1,497	215	8,586	14,640							-	-	-
	1959	1,575	3,136	586	736	167	9,996	16,196							32	171	203
	1960	2,032	5,910	600	1,885	369	10,541	21,337							-	-	-
	1961	2,710	6,364	662	3,193	599	9,124	22,652							-	130	130
	1962	2,545	5,769	747	1,683	293	10,657	21,694							-	294	294
	1963	2,797	6,077	1,256	2,542	294	9,786	22,752							-	412	412
	1964	1,475	3,140	1,037	2,784	1,884	8,973	19,293							-	131	131
	1965	2,121	2,569	831	1,963	1,106	11,496	20,086							-	289	289
	1966	1,261	1,370	613	1,614	129	10,082	15,069							-	435	435
	1967	2,603	878	1,210	3,273	302	6,462	14,728							-	371	371
	1968	3,058	500	983	1,568	217	9,268	15,594							-	195	195
	1969	2,187	878	721	2,219	195	3,236	9,436							-	260	260
	1970	1,779	607	723	1,198	224	2,907	7,438							-	92	92
	1971	1,555	697	938	1,492	317	3,721	8,720		0				0	-	555	555
	1972	1,107	512	944	842	197	4,212	7,814		0				0	-	1,646	1,646
	1973	2,351	838	526	2,108	636	2,266	8,725		0				0	-	1,084	1,084
	1974	6,019	1,177	1,192	1,656	754	4,106	14,904		0				0	-	344	344
	1975	2,433	1,061	1,401	1,031	808	4,491	11,225		3				3	-	2,145	2,145
	1976	2,996	320	1,082	830	1,237	2,148	8,613		5				5	-	1,968	1,968
	1977	2,257	338	2,256	2,166	1,052	5,110	13,179		0				0	-	2,186	2,186
	1978	2,546	648	1,154	4,517	2,276	10,427	21,568		3				3	-	545	545
	1979	4,558	729	1,250	2,655	2,429	13,881	25,502		0				0	-	213	213
	1980	2,521	811	1,392	1,531	1,953	11,327	19,535		0				0	-	582	582
	1981	2,129	590	754	1,777	2,653	25,422	33,325		0				0	-	218	218
	1982	1,667	718	1,777	864	1,709	19,234	25,969		0	31			31	-	506	506
	1983	972	217	356	2,028	1,117	14,774	19,464		0	13			13	_	214	214
	1984	2,234	142	587	1,874	868	4,433	10,138		1	4			5	-	166	166
	1985	2,562	105	1,817	1,850	1,175	4,154	11,663		0	1			1		676	676

) Catch of s	et net in 2013 w	ere updated	based on the .	Japanese o	official statis	stics of an	nual catch.										
	istics of Korea v				tatistics for	1982-199	9.										
	troll catch since				oio was fer	viscu as a	result of de	acting dodoic coul	innig and C	manging th	c data sourc	c (13C13/3)	MINC	J/ 17 F -4).			_
	paranthesis are p		itch data from	2007 to 2	013 was ray	viced ac a	result of de	leting double cou	nting and	hanging th	e data source	e (ISC15/97	ΓΔΤΨ	5/WP_4)			+
otal	oronthosis s	129,470	(90,163)	54,708	129,155	46,975	559,777	(1,010,248)	11	25	24,973	75	0	25,083	245	97,865	98,11
	ain catch total		(90,163)		129,155		559,777	(1,010,248)	11	25	24,973	75	0	25,083	245	97,865	98,11
	2016	1,225	(654)	37	779	508	5,096	8,299	3	0	1,024	2	0	1,029		2,706	2,70
	2015	1,267	619	4		407	3,645	6,354	1	0	676	+	0	676		3,082	3,08
	2014	1,907	683	5	1,023	499	5,456	9,573	6	0	1,305	+	0	1,311		4,862	4,86
	2013	1,415 #	784.12	8	904	529	2,771	6,411	1	0	604	+	+	605		3,154	3,1
	2012	1,932	673	113	570	343	2,462	6,093		0	1,421	2		1,423	1	6,667	6,6
	2011	1,651	837	63	1,820	283	8,340	12,994		0	670	14	+	684	1	2,730	2,7
	2010	1,603	890	83	1,583	495	3,742	8,396		0	1,196	16		1,212	-	7,746	7,7
	2009	2,236	1,080	50	2,003	575	8,077	14,021		0	936	4		940	-	3,019	3,0
	2008	2,358	1,390	64	2,377	770	10,221	17,180		0	1,866	10		1,876	15	4,392	4,
	2007	1,503	1,762	236	2,823	657	6,840	13,821		0	1,281	4		1,285	-	4,147	4,
	2006	1,421	1,121	108	1,860	777	8,880	14,167		0	1,012	3		1,015	-	9,927	9,
	2005	2,182	1,925	549	3,633	548	12,817	21,654		0	1,318	9		1,327	_	4,542	4,
	2004	896	1,847	132	2,445	514	8,257	14,091		0	773	0		773	14	8,880	8,
	2003	839	1,237	44	2,060	446	5,768	10,394		0	2,601	0		2,601	43	3,211	3,2
	2002	1,100	841	99	2,607	631	8,903	14,181		0	932	1		933	2	1,708	1,
	2001	1,366	791	180	3,466	782	7,620	14,205		0	1,176	10		1.186		863],
	2000	1,125	953	102	5,217	689	16,486	22,611 24,572		0	2,401	0		2,401	99	3,019	3,
	1998 1999	874 1,097	1,255 1,157	85 35	2,876 3,440	700 709	5,391 16,173	11,181		0	188 256			188 256	35	2,369	2,4
	1997	803	1,300	34	2,740	811	13,152	18,840		0	1,054			1,054	-	367	3
	1996	1,149	901	94	3,640	570	7,644	13,998		0	102			102		3,700	3,7
	1995	1,859	678	270	4,778	586	18,909	27,080		0	821			821		11	
	1994	1,158	1,206	162	4,111	398	8,131	15,166		0	50			50	2	63	
	1993	848	812	129	546	365	6,600	9,300		0	40			40	-	-	
	1992	1,208	513	166	915	1,081	7,162	11,045		0	288			288	-	0	
	1991	1,485	218	286	2,069	650	9,808	14,516		0	265			265	-	9	
	1990	716	309	536	1,250	455	2,989	6,255		0	132			132	-	50	
	1989	748	209	754	903	283	6,190	9,087		0	71			71	-	57	
	1988	843	157	907	1,124	498	3,605	7,134		0	32			32	1	447	
	1987	2,198	211	1,565	880	445	8,653	13,952		13	89			102	_	119	

⁷²

Table 15-2. Continued

			TWN								USA					
t-net	Gill-net (not specified)	Drift gill-ne t	Longlin e	Others	Purse seine	TWN Total	Drift gill-net	Longlin e	Pole and line	Troll	Hook and Line	Others	Purse seine	Sport	USA Total ⁶	Total
													2,076	2	2,078	19,162
													4,433	48	4,481	20,110
													9,537	11	9,548	28,547
													6,173	93	6,266	31,988
													5,727	388	6,115	40,144
													9,215	73	9,288	36,543
													13,934	10	13,944	28,584
									56				3,506	13	3,575	19,974
									+				4,547	1	4,548	25,885
									16				7,989	23	8,028	30,810
									+				10,769	25	10,794	32,782
									28				11,832	7	11,867	35,031
									39				9,047	7	9,093	28,517
			54			54			11	+		66	6,523	1	6,601	27,030
			-			-			12				15,450	20	15,482	30,986
			53			53			+				5,517	32	5,549	20,701
			33			33			8				5,773	12	5,793	21,615
			23			23			9				6,657	15	6,681	16,400
			-			-			+				3,873	19	3,892	11,422
			1			1			+				7,804	8	7,812	17,088
			14			14			3			42	11,656	15	11,716	21,190
			33			33			5	+		20	9,639	54	9,718	19,560
			47	15		62			+	+		30	5,243	58	5,331	20,641
			61	5		66			83			1	7,353	34	7,471	20,910
			17	2		19			22	+		3	8,652	21	8,698	19,303
			131	2		133			10			3	3,259	19	3,291	18,789
			66	2		68			4			2	4,663	5	4,674	26,858
			58	-		58			5			1	5,889	11	5,906	31,679
			114	5		119			+			24	2,327	7	2,358	22,594
			179	-		179	4		+	10		+	867	9	890	34,612
		2	207	-		209	9		1			+	2,639	11	2,660	29,375
		2	175	-	9	186	31		59			2	629	33	754	20,631
			477	8	5	490	6	1	5			18	673	49	752	11,551
		11	210		80	301	8	-				20	3,320	89	3,437	16,078

		13	70	-	16	99	16					41	4,851	12	4,920	19,252
		14	365	-	21	400	2					18	861	34	915	15,488
		37	108	25	197	367	4					46	923	6	979	8,960
		51	205	3	259	518	3					18	1,046	112	1,179	10,912
		299	189	16	149	653	11					81	1,380	65	1,537	8,627
		107	342	12	-	461	4	2				+	410	92	508	15,759
		3	464	5	73	545	9	38				14	1,928	110	2,099	13,977
			471	3	1	475	32	42				29	580	283	966	10,781
			559	-		559	28	30				1	906	86	1,051	16,891
			335	2		337	20	29				+	657	245	951	29,200
-	-		956	-	-	956	43	25		2		+	4,639	40	4,749	23,505
-	-		1,814	-	-	1,814	58	26		1		48	2,240	131	2,504	24,579
-	-		1,910	-	-	1,910	40	54		128		59	1,771	422	2,474	15,754
-	-		3,089	-	-	3,089	22	54		20		88	184	408	776	29,136
-	1		2,780	1	-	2,782	30	19		1		11	693	319	1,073	33,946
-	2		1,839	2	-	1,843	35	6		6		1	292	344	684	18,781
-	3		1,523	1	-	1,527	7	2		1		2	50	613	675	19,026
-	10		1,863	11	-	1,884	14	1				3	22	355	395	18,528
-	1		1,714	2	-	1,717	10	1				+		50	61	25,536
1	-		1,368	1	-	1,370	5	1				1	201	73	281	29,174
1	-		1,149	-	-	1,150	1	1				+		94	96	26,355
2	8		1,401	-	-	1,411	2	+				+	42	12	56	20,720
1	1		979	-	-	981	1	+				+		63	64	24,508
1	10		877	-	-	888	3	1		0		2	410	156	572	19,440
29	7		373	-	-	409	1	0				0		88	89	17,852
16	7		292	1	0	316	18	0		0		100		225	343	17,068
2	0		210	2	-	214	4	0		0		38		400	442	14,840
2	1		331	-	-	334	7	1		0		3		809	820	11,324
38	4		483	-	-	525	5	0		0		2	401	436	844	17,115
25	1		552	-	-	578	4				26	0	68	382	480	11,171
25	1		454	-	-	480	8				30	1	316	298	653	(13,167)
143	57	539	31,018	126	810	32693	506	334	376	169	56	839	242,062	7,986	252,328	(1,418,462)
143	57	539	31,018	126	810	32254	506	334	376	169	56	839	242,062	7,986	252,328	(1,418,462)

Table 15-3. Annual catch of swordfish (*Xiphias gladius*) in metric tons for fisheries monitored by ISC for assessments of North Pacific Ocean stocks, 1951-2010. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional.

				JI	PN			KC	R		MEX	
Catch disposition	Year	Set-net	Drift gill- net	Longline	Others	Not specified	JPN Total	Longline	KOR Total	Others	Sport	MEX Total
Retain catch	1951	78	10	7,246	4,246	98	11,678					
	1952	68	-	8,890	2,721	12	11,691					
	1953	21	_	10,796	1,484	107	12,408					
	1954	18	-	12,563	909	121	13,611					
	1955	37	-	13,064	850	160	14,111					
	1956	31	-	14,596	786	73	15,486					
	1957	18	_	14,268	895	70	15,251					
	1958	31	_	18,525	1,111	67	19,734					
	1959	31	-	17,236	956	44	18,267					
	1960	67	1	20,058	1,243	30	21,399					
	1961	15	2	19,715	1,386	30	21,148					
	1962	15	-	10,607	1,449	44	12,115					
	1963	17	-	10,322	845	59	11,243					
	1964	16	4	7,669	1,097	66	8,852					
	1965	14	0	8,742	2,027	208	10,991					
	1966	11	0	9,866	1,841	45	11,763					
	1967	12	0	10,883	1,075	38	12,008					
	1968	14	0	9,810	1,775	50	11,649					
	1969	11	0	9,702	1,567	56	11,336					
	1970	9	0	7,715	1,784	39	9,547					
	1971	37	1	7,369	491	48	7,946	0	0			
	1972	1	55	7,316	293	22	7,687	0	0			
	1973	23	720	7,564	131	29	8,467	0	0			
	1974	16	1,304	6,523	336	29	8,208	0	0			
	1975	18	2,672	7,659	223	60	10,632	0	0			
	1976	14	3,488	8,786	372	182	12,842	0	0			
	1977	7	2,344	9,255	247	73	11,926	0	0			
	1978	22	2,475	9,022	177	111	11,807	0	0			
	1979	15	983	9,627	226	49	10,900	0	0			
	1980	15	1,746	6,873	423	30	9,087	135	135			
	1981	9	1,848	7,789	181	61	9,888	0	0			
	1982	7	1,257	6,963	230	59	8,516	166	166			
	1983	9	1,033	8,708	210	32	9,992	47	47			
	1984	13	1,053	8,375	153	98	9,692	27	27			
	1985	10		10,368	210	69	11,790	12	12			

Total		re provisio	` ' '	(575,961)	(41,085)	(6,323)	(673,213)	11,174	11,174	3,430	-	3,430
	lease total	(0.00.0	(40 == 0	(*** • • • • • • • • • • • • • • • • • •	(44.00=)	// aa=:	//=a a 4 5 i	44.4=:	44.4	2 420		2.45*
	2011											
Release	2010											
Retain	catch total	(933)	(44,041)	(580,152)	(41,493)	(6,594)	(673,213)	11,756	11,756	3,430	0	3,430
	2016	(3)	(277)	(4,119)	(205)	(281)	(4,885)	582	582	-		
	2015	3	277	5,024	205	281	5,790	776	776	-	-	
	2014	7	269	4,367	2	291	4,936	829	829	0	0	C
	2013	13	290	4,240	168	291	5,002	1,071	1,071	0	0	(
	2012	8	371	4,030	63	288	4,760	856	856	0	0	(
	2011	2	193	4,019	12	233	4,460	962	962	-	-	
	2010	8	494	5,398	112	230	6,241	662	662	222	-1	222
	2009	3	682	6,006	240	249	7,180	993	993	394	-1	394
	2008	3	648	6,187	175	349	7,362	794	794	242	-	242
	2007	2	829	8,123	124	368	9,446	453	453	172	_	17
	2006	5	796	7,687	247	343	9,078	465	465	328	_	32
	2005	3	956	6,647	193	337	8,136	403	403	234		23
	2003	4	1,062	6,900	233	30	8,229	410	410	268	-	26
	2002	4	1,104	6,434	154	29	7,818	380	380	237	-	23
	2001	15	1,164	7,831 7,185	239	30	8,847 8,600	438	438	516 215	-	51 21
	2000	5 15	808 732	7,292	506 239	49	8,660	202 438	202 438	602	-	60
	1999	5	724	6,604	418	47	7,798	131	131	602	-	
	1998	2	471	7,419	480	68	8,440	153	153		-	
	1997	5	365	8,215	347	62	8,994	99	99		-	
	1996	4	428	7,979	601	45	9,057	15	15		-	
	1995	7	561	7,808	424	34	8,834	9	9		-	
	1994	4	421	8,723	308	37	9,493	4	4		-	
	1993	4	292	9,777	309	44	10,426	27	27		-	
	1992	6	887	8,497	389	16	9,795	32	32		-	
	1991	5	498	5,682	161	20	6,366	37	37		-	
	1990	4	1,074	6,595	131	13	7,817	46	46		-	
	1989	10	1,596	7,482	372	21	9,481	7	7			
	1988	8	1,234	9,304	186	19	10,751	27	27			
	1987	11	1,051	10,370	128	45	11,258 11,605	50	50			

Table 15-3. Continued.

			TWN								U	SA					
Set-net	Gill-net (not specified	Harpoon		Others	Purse seine	TWN Total	Drift gill- net	Harpoon	Handline	Longline	Pole and line	Troll	Others	Purse seine	Sport	USA Total	Total
																	11,678
																	11,691
			-			-											12,408
			-			-											13,611
			-			-											14,111
			-			-											15,486
			-			-											15,251
			-			-											19,734
			427			427											18,694
			520			520											21,919
			318			318											21,466
			494			494											12,609
			343			343											11,586
			358			358											9,210
			331			331											11,322
			489			489											12,252
		5	646	30		681											12,689
	- 8	3	763	1		775											12,424
	- 1	6	843	-		850											12,186
	- 1	5	904	-		910		612		5						617	11,074
		3	992	_		995		99		1						100	9,041
		12	862			874		171								171	8,732
		113	860	6		979		399								399	9,845
		98	881	38		1,017		406								406	9,631
		152	928	1		1,081		557								557	12,270
		159	636	35		830		42								42	13,714
	- 2	139	578	- 33		719		318		17						335	12,980
	. 3	10	546			559		1,699		9						1,708	14,074
	-	24	668	4		701		329		7						336	11,937
		72	613	1		690	160	566		5						731	10,643
	3	18	658	4		683	473	271		3	2					749	11,320
	. 3	46	856	-		905	945	156		5	3	6	1			1,116	10,703
	3	164	783	_		950	1,693	58		5	2	3		1		1,763	12,752
43	5	259	733	-		1,040	2,647	104		15	49		1	26		2,841	13,600
3		166	566	61		825	2,990	305	4	2	47		104	20		3,405	16,032

3	1	201	456	6		667	2,069	291	4	2			109			2,475	14,418
-	-	187	1,331	3		1,521	1,529	235	4	24			31			1,823	14,999
-	1	80	777	183		1,041	1,376	198	6	24			64			1,668	13,487
3	2	61	1,541	35		1,642	1,243	62	7	218			56			1,586	12,716
4	2	118	1,452	88		1,664	1,131	64	5	2,437			43			3,680	13,207
4	2	205	1,430	56		1,697	944	20	6	4,535			44			5,549	13,649
12	1	287	1,494	33		1,827	1,356	75	1	5,762			47			7,241	18,895
13	3	194	1,228	100		1,538	1,412	168	4	5,936			161			7,681	19,672
12	3	211	1,155	9		1,390	792	157	4	3,807			24			4,784	15,671
6	2	14	1,185	203		1,410	771	97	6	2,981			29			3,884	14,137
10	2	19	710	1	-	742	761	81	5	2,848			15			3,710	13,524
8	1	27	1,397	1	-	1,434	708	84	7	3,393			11			4,203	14,730
15	9	17	1,198	-	-	1,239	931	48	7	3,681			19			4,686	14,518
5	5	51	1,455	+	-	1,516	606	81	9	4,329			27			5,052	14,497
5	6	74	3,716	-	-	3,801	649	90		4,834			33			5,606	18,871
8	18	64	4,853	-	-	4,943	375	52		1,969			19			2,415	17,159
16	8	1	5,400	1	-	5,426	302	90		1,524			3			1,919	16,598
8	3	-	4,771	-	-	4,782	216	107	10	1,958			11			2,302	15,519
7	6	1	4,248	2	-	4,264	182	69	7	1,185			44			1,487	14,658
5	3	16	3,964	2	-	3,990	220	77	5	1,622			5			1,929	14,692
7	2	49	4,382	3	-	4,443	443	71	4	1,211			5			1,734	16,048
2	2	20	4,099	2	-	4,125	490	59	5	1,735		1				2,290	16,486
3	6	39	3,745	+	-	3,793	405	48	6	2,014			19			2,492	14,683
83	7	31	3,550	-	-	3,671	253	50	5	1,817		0	0			2,125	14,363
6	4	42	2,844	-	-	2,896	62	37	3	1,676			18			1,796	11,817
8	17	52	3,577	1	+	3,655	119	24	5	1,623			90			1,861	10,938
3	15	30	3,746	0	0	3,794	118	5	6	1,395		1	1			1,526	10,936
2	8	0	2,846	1	0	2,857	95	6	6	1,270		1	7			1,385	10,315
4	4	0	2,817	0	0	2,825	124	5	7	1,665		1	+			1,802	10,392
4	4	0	3,199	0	0	3,207	97	5.3	5	1,516		1	14			1,640	11,413
(4)	(4)	(0)	(2,054)	(0)	(0)	(2,062)	176	25	4	1,092			36			1,339	8,867
(316)	(218)	(3545)	(98216)	(911)	(0)	(103,206)	28,863	8,573	157	70,157	56	14	1,091	27	+	(108,939)	(900,543)
. /	. /			. /						·				0		0	0
														0		0	0
														0		0	0
(312)	(214)	(3,545)	(96,162)	(911)	(0)	(101,144)	28,656	8,548	153	69,064	56	14	1,042	27	+	107,560	(900,543)

Table 15-4. Annual catch of striped marlin (*Kajikia audax*) in metric tons for fisheries monitored by ISC for assessments of North Pacific Ocean stocks, 1951-2011 "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recents years are provisional.

				J	PN		•		KOR		M	EX
Catch disposition	Year	Set-net	Drift gill-net	Longline	Others	Not specified	JPN Total	Longline	Purse seine	KOR Total	Sport	MEX Total
Retain catch	1951	92	-	2,494	1,822	39	4,447					
	1952	203	-	2,901	2,043	40	5,187					
	1953	126	-	2,138	840	36	3,140					
	1954	82	-	3,068	990	67	4,207					
	1955	106	-	3,082	878	82	4,148					
	1956	133	-	3,729	1,881	41	5,784					
	1957	71	-	3,189	2,431	76	5,767					
	1958	82	3	4,106	2,981	127	7,299					
	1959	87	2	4,152	3,061	200	7,502					
	1960	161	4	3,862	1,790	87	5,904					
	1961	161	2	4,420	1,707	98	6,388					
	1962	197	8	5,739	1,717	108	7,769					
	1963	92	17	6,135	1,590	292	8,126					
	1964	81	2	14,304	2,264	41	16,692					
	1965	81	1	11,602	2,659	73	14,416					
	1966	226	2	8,419	1,425	31	10,103					
	1967	82	3	11,698	1,521	75	13,379					
	1968	71	0	15,913	1,144	58	17,186					
	1969	71	3	9,144	2,519	81	11,818					
	1970	55	3	13,686	1,005	153	14,902					
	1971	61	10	11,632	1,933	307	13,943	0		0		
	1972	72	243	7,843	972	94	9,224	0		0		
	1973	80	3,265	6,989	594	146	11,074	0		0		
	1974	90	3,112	7,027	630	104	10,963	0		0		
	1975	105	6,534	5,567	530	89	12,825	0		0		
	1976	37	3,561	5,380	475	107	9,560	0		0		
	1977	103	4,424	3,275	352	107	8,261	0		0		
	1978	93	5,593	4,200	237	243	10,366	0		0		
	1979	66	2,532	5,927	348	133	9,006	0		0		
	1980	80	3,467	6,985	402	59	10,993	73		73		
	1981	88	3,866	4,365	397	69	8,785	0		0		
	1982	52	2,351	5,653	489	128	8,673	102		102		
	1983	124	1,867	4,042	557	156	6,746	49		49		
	1984	144	2,333	3,892	407	177	6,953	39		39		
	1985	81	2,363	4,608	523	153	7,728	13		13		

Fotal Numbers in pa	oronthocus		(80,614)	(314,069)	(49,584)	(6,454)	(454,166)	6,246	0	6,246		
	ase total	(4.050)	(00.61.5)	(21.1.0.00)	(40.50.0)	(C. 4 = 1)	(454465)	6.046	0	0		
	2016								0	0		
	2011											
Release	2010											
Retain ca	atch total	(4,972)	(80,614)	(314,069)	(49,584)	(6,454)	(455,693)	6,246		6,246	-	
	2016	(37)	(287)	(889)	(80)	(27)	(1,320)	61		61	_	
	2015	(37)	(287)	(1,277)	(80)	(27)	(1,708)	44		44	-	
	2014	35	173	1,111	35	22	1,376	82		82	-	
	2013	39	336	1,457	52	34	1,918	65		65	-	
	2012	52	597	1,307	62	34	2,052	33		33	-	
	2011	30	347	1,251	87	26	1,741	48		48	-	
	2010	20	913	1,019	68	36	2,056	18		18	-	
	2009	17	821	788	55	39	1,720	22		22	-	
	2008	26	1,302	999	53	28	2,408	29		29	-	
	2007	21	970	1,166	25	38	2,220	46		46	-	
	2006	30	1,190	1,161	34	32	2,447	55		55	-	
	2005	28	1,214	1,230	36	35	2,543	136		136	_	
	2004	23	1,339	1,699	35	34	3,130	75		75	_	
	2002	41	1,064	1,855	138	27	3,125	205		205	_	
	2002	80	1,264	1,560	187	28	3,119	187		187	_	
	2001	51	1,002	2,311	161	36	3,636	253		253	_	
	2000	41	1,062	2,261	173	91	3,628	536		536		
	1998	28	1,126	3,159	70	90	4,473	433		433	-	
	1997	34	1,092	3,759	125	147	5,157	635		635	-	
	1996	34	813	3,787 3,520	64	68	4,499	1,017		1,017	_	
	1995 1996	39	970 703	5,334	142 29	88	6,590 4,646	307 429		429	-	
	1994	73 58	1,443	4,195	219	92 86	6,022	556		556 307	-	
	1993	27	828	5,199	309	373	6,736	568		568	-	
	1992	71	1,204	4,202	147	95	5,719	53		53	-	
	1991	56	1,329	4,042	323	131	5,881	7		7	-	
	1990	65	1,945	3,493	290	193	5,986	1		1	-	
	1989	47	1,664	5,821	358	145	8,035	24		24		
	1988	63	2,211	7,023	407	205	9,909	16		16		
	1987	102	1,888	8,725	250	167	11,132	15		15		
	1986	131	3,584	7,303	376	103	11,497	14		14		

Table 15-4. Continued.

			TWN					-		USA				
Set-net	Gill-net (not specified	Harpoon	Longline	Others	Purse seine	TWN Total	Handline	Longline	Troll	Others	Purse seine	Sport	USA Total	Total
														4,447
												23	23	5,210
				-		-						5	5	3,145
				-		-						16	16	4,223
				-		-						5	5	4,153
				-		-						34	34	5,818
				-		-						42	42	5,809
			543	387		930						59	59	8,288
			391	354		745						65	65	8,312
			398	350		748						30	30	6,682
			306	342		648						24	24	7,060
			332	211		543						5	5	8,317
			560	199		759						68	68	8,953
			392	175		567						58	58	17,317
			355	157		512						23	23	14,951
			370	180		550						36	36	10,689
-	-	141	387	63		591						49	49	14,019
-	40	134	333	34		541						51	51	17,778
-	5	159	573	28		765						30	30	12,613
-	. 8	175	495	6		684						18	18	15,604
-	16	101	449	18		584						17	17	14,544
-	1	124	389	1		515						21	21	9,760
-	4	115	569	20		708						9	9	11,791
-	7	53	674	58		792						55	55	11,810
-	7	86	796	3		892						27	27	13,744
-	. 9	61	379	70		519						31	31	10,110
-	. 9	207	541	3		760						41	41	9,062
-	7	70	618	1		696						37	37	11,099
2	18	104	458	-		582						36	36	9,624
-	39	92	284	1		416						33	33	11,515
-	25	70	508	-		603						60	60	9,448
-	20	112	404	-		542						41	41	9,358
		144	555	39		769						39	39	7,603
-	16	314	965	-		1,295						36	36	8,323
1	6	152	513	23		695			18			42	60	8,496

(90)	(870)	(6,454)	(23,785)	(2,967)	(3)		8	12,192	862	0	2	1,484	14,144	(496,746
					1	1					1		1	2
					1	1					0		0	1
											0		0	0
(34)	(013)	(3,170)	(43,103)	(4,307)	(2)	(34,007)	0	14,174	002	U	1	1,704	14,143	1
(92)	(873)	(5,148)	(23,785)	(2,967)	(2)	(32,867)	8	12,192	862	0	1	1,484	14,143	(496,744
(1)	(4)	(28)	(214)	(0)	(1)	(248)		392	11	0			404	(2,033
1	4	28	228	0	0	210		426	11	0		1	504	(2,517
0	5	64	140	0	1	210		398 426	8			1	406	2,982 2,107
0	24	114 197	443 372	0	0	592 593		398	11				293 406	2,970
1	34		343	4				362 282			0			2,637
- 1	42 27	131 95	253	3	0	429 470		165 362	19 16		0		185 378	2,688
-	46	92	262	-	-	400		258	10		1		268	2,410
-	43	168	270	2	-	483		427	14				441	3,361
-	29	141	301	-	-	471		276	13			+	289	3,026
-	30	117	741	+	-	888	+	611	21			+	632	4,022
9	9	5	783	9	-	815	+	511	20			+	531	4,025
5	10	7	522	2	-	546	2	376	31			+	409	4,160
5	27	11	922	+	-	965	+	538	29			+	567	4,862
8	15	6	477	-	-	506	+	226	30			+	256	4,068
5	16	30	250	-	-	301		351	44			+	395	4,585
6	18	29	315	1	-	369		200	15			10	225	4,758
5	8	26	197	-	-	236	1	364	27			12	404	5,546
6	16	19	304	-	-	345	+	378	26			23	427	6,564
3	9	33	351	-	-	396	1	352	37			21	411	6,323
3	8	30	188	6	-	235	1	418	53			20	492	5,802
4	5	67	110	6		192	+	543	52			14	609	7,698
4	5	176	138	11		334	+	326	35			17	378	7,290
3	7	132	226	-		368	1	471	67			11	550	8,222
-	11	163	220	24		418	1	459	37			25	522	6,712
-	81	173	290	32		576	+	663	41			12	716	7,180
12	16	207	139	21		395	+	538	27			23	588	6,970
-	23	124	192	10		349	+	612	24			52	688	9,090
7	12	70	464	80		633	_	504	54			30	588	11,140
1	13	119 132	179 414	16 16		327 565	1	272	19 29			19 28	38	11,870 12,042

Table 15-5. Retained catches (metric tons, whole weight) of ISC Members of blue marlin (*Makaira nigricans*) by fishery in the North Pacific Ocean, north of the equator. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional.

		JP	N		KOR		M	EX
Catch disposition	Year	Longline	JPN Total	Longline	Purse seine	KOR Total	Sport	MEX Total
Retain catch	1953							
	1954							
	1955							
	1956							
	1957							
	1958							
	1959							
	1960							
	1961							
	1962							
	1963							
	1964							
	1965							
	1966							
	1967							
	1968							
	1969							
	1970							
	1971	5,461	5,461	0		0		
	1972	6,772	6,772	0		0		
	1973	6,453	6,453	0		0		
	1974		6,545	0		0		
	1975	4,374	4,374	0		0		
	1976		5,018	0		0		
	1977	4,780	4,780	0		0		
	1978		5,900	0		0		
	1979		5,949	0		0		
	1980		5,613	155		155		
	1981	5,518	5,518	0		0		
	1982	6,051	6,051	351		351		
	1983		4,796	82		82		
	1984		6,248	155		155		
	1985		5,164	45		45		

Total Numbers in p	o mo m th o cay		(189,507)	8,837	1	8,838	-	-
	ase total	(400 F0=)	(400 505)	0.025	1	1		
	2016				1	1		
	2015				0	0		
	2013							
	2011							
Release	2010							
		(189,507)	(189,507)	8,837		8,837	-	-
	2016		(1,722)	1,116		1,116	-	-
	2015	(1,642)	(1,642)	531		531	-	-
	2014	1,898	1,898	801		801	-	-
	2013	2,176	2,176	963		963	-	-
	2012	2,019	2,019	587		587	-	
	2011	2,211	2,211	684		684	-	
	2010		2,457	257		257	-	
	2009	1,840	1,840	224		224	-	-
	2008	2,096	2,096	219		219	-	-
	2007	2,016	2,016	120		120	-	-
	2006		2,414	217		217	-	-
	2005	2,506	2,506	303		303	-	
	2004	2,977	2,977	226		226	-	
	2003	2,836	2,836	158		158	-	-
	2002	2,976	2,976	152		152	-	
	2001	3,594	3,594	166		166	-	-
	2000	3,606	3,606	96		96	-	-
	1999	3,751	3,751	164		164	-	_
	1998	3,438	3,438	335		335	-	-
	1997	3,697	3,697	145		145	-	-
	1996	3,260	3,260	10		10	-	
	1995	5,907	5,907	0		0	-	-
	1994	5,832	5,832	2		2	-	_
	1993	4,600	4,600	36		36	-	_
	1992	3,721	3,721	60		60	-	_
	1991	4,094	4,094	75		75	_	
	1990		4,116	44		44	_	_
	1989	5,117	5,117	50		50		
	1988	5,054	5,054	133		133		
	1986 1987		5,922 5,370	86 89		86 89		

Table 15-5. Continued.

			TWN				USA							
Set-net	Gill-net (not specified	Harpoon	Longline	Others	Purse seine	TWN Total	Handline	Longline	Troll	Others	Purse seine	USA Total	Total	
			-			-								
			-			-								
			-			-								
			-			-								
			-			-								
			887			887							88'	
			781			781							78	
			948			948							94	
			703			703							703	
			628			628							623	
			691			691							69	
			934			934							93	
			1,016			1,016							1,01	
			957			957							95'	
-	-	317	898	167		1,382							1,38	
-	30	649	1,433	120		2,232							2,23	
-	- 58	465	1,232	103		1,858							1,85	
1	21	604	1,385	70		2,081							2,08	
-	- 13	473	1,331	118		1,935							7,39	
-	14	490	1,205	50		1,759							8,53	
-	12	275	1,650	265		2,202							8,65	
1	6	355	2,144	146		2,652							9,19	
-	. 3	421	2,638	207		3,269							7,64	
-	. 9	511	1,315	162		1,997							7,01	
-	- 11	391	1,183	110		1,695							6,47	
1	15	364	1,633	7		2,020							7,92	
3	19	362	1,646	164		2,194							8,14	
_	35	444	1,185	170		1,834							7,602	
-	35	313	1,840	69		2,257							7,77	
-	_	306	2,139	120		2,572							8,97	
-	26	741	2,122	127		3,016							7,89	
-		960	1,789	111		2,882							9,28	
9	11	747	1,187	43		1,997			145			145	7,351	

					5	5					6	6	6 5
(= = 3)	(-,)	(10,000)	(22 ,22 0)	(= ;= ;=)	(-1)	-,			- , 0		1	1	1
(368)	(1,451)		(169,398)	(3,599)		(191,738)	21	10,961	7,338	6		18,326	(408,408)
(3)	(14)	(177)	(3,398)	(0)	(4)	(3,596)	2	564	157			723	7,157
3	14	177	4,306	0	3	4,503	3	631	196			830	7,506
4	11	124	4,861	5	7	5,012	4	535	159			698	8,409
2	6	16	4,607	0	9	4,640	3	406	137			546	8,325
6	13	16	4,023	0	12	4,144	2	298	141			441	7,191
3	18	17	4,625	9	2	4,674	2	373	199			574	8,143
5	27	15	4,959	1		5,007	2	306	148			456	8,177
9	12	9	4,638	1		4,669	1	360	15			376	7,109
10	16	15	5,477	1	_	5,519	1	349	198			548	8,382
3	17	20	5,117	+	_	5,157	1	262	127			390	7,683
12	15	30	5,729	-	-	5,786		409	160			569	8,986
8	65	48	7,630	16	-	7,767		337	187			524	11,100
14	93	36	6,672	9	-	6,824		283	188	5		476	10,503
20	107	52	7,685	4	-	7,868		363	210			573	11,435
13	63	32	4,448	6	-	4,562		264	225	1		490	8,180
18	57	229	4,733	-	-	5,037		399	291			690	9,487
24	40	186	4,131	2	-	4,383		314	235			549	8,634
21	32	135	3,417	-	-	3,605		357	293			650	8,170
8	59	91	3,624	1	-	3,783		395	242			637	8,193
5	48	194	3,683	-	_	3,930		487	378			865	8,637
13	35	270	3,337	10	_	3,665		467	409			876	7,811
15	36	895	3,896	5		4,847		570	315			885	11,639
12	30	206	3,007	7		3,262		362	298			660	9,756
44	41	400	4,135	82		4,702		339	311			650	9,988
25	40	432	3,786	110		4,393		347	260			607	8,781
4	50	338	2,288	152		2,832		297	346			643	7,644
3	24	427	1,749	143		2,346		378	295			673	7,179
10	14	640	2,691	9		3,364		356	326			682	9,213
20	8	658	2,822	589		4,097		102	266			368	9,652
12	90	839 973	1,723 4,627	107		2,763 5,622		51	220 261			220 312	8,991 11,393

Table 15-6. Retained catches (metric tons, whole weight) of ISC Members of blue sharks (*Prionace glauca*) by fishery in the North Pacific Ocean, north of the equator. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional.

				JP	'n			KO)R	MI	EX
Catch disposition	Year	Set-net	Drift gill- net	Longline	Others	Not specified	JPN Total	Longline	KOR Total	Others	MEX Total
Retain catch	1985										
	1986										
	1987										
	1988										
	1989										
	1990										
	1991										
	1992										
	1993										
	1994			20,062			20,062				
	1995			18,427			18,427				
	1996			21,251			21,251				
	1997			26,105			26,105				
	1998			23,988			23,988				
	1999			26,541			26,541				
	2000			27,511			27,511				
	2001			28,126			28,126			-	-
	2002			26,345			26,345			-	-
	2003			26,278			26,278			-	-
	2004			22,470			22,470			-	-
	2005			21,887			21,887			-	-
	2006			19,063			19,063			-	-
	2007			15,190			15,190			2,073	2,073
	2008			21,773			21,773			3,531	3,531
	2009	6	908	19,260	730	1	20,905			3,261	3,261
	2010	9	733	22,633	802	1	24,177			3,700	3,700
	2011	7	438	18,780	862	3	20,090			3,366	3,366
	2012	2	1,076	13,132	762	3	14,975			4,108	4,108
	2013	6	1,103	16,581	626	2	18,319	75	75	4,494	4,494
	2014	4	1,060	15,644	598	2	17,306	100	100	5,513	5,513
	2015	21	697	12,083	387	2	13,190	53	53	-	-
	2016	(21)	(697)	(12,083)	(387)	(2)	(13,190)	0	0	-	-
Retain cat	ch total	(32)	(5,344)	(466,171)	(4,989)	(14)	(473,979)	228	228	30,046 2	30,046
Release	2015							0	0		
	2016							8	8		
	se total							8	8		
Total		(32)	(5344)	(466171)	(4989)	(14)	(473979)	236	236	30,046	30,046

TV	VN			U	SA			
Longline	TWN Total	Drift gill-net	Longline	Troll	Others	Sport	USA Total	Total
		+			1		1	1
		1			1		2	2
		1			1		2	2
		+			3		3	3
					6		6	6
		+			20		20	20
		+			1		1	1
		1			1		2	2
		+			+		0	0
		+			12		12	20,074
		+			5		5	18,432
		+			+		0	21,251
		+			+		0	26,105
		+			1		1	23,989
		+			+		0	26,541
		+			+		0	27,511
					+		+	28,126
					+		+	26,345
		+			+		0	26,278
					+		+	22,470
					+		+	21,887
					+		+	19,063
		9	8		+		17	17,280
			7				7	25,311
11,541	11,541	1	9		1		11	35,718
7,670	7,670	+			0		7	35,554
13,117	13,117		13		0		13	36,586
10,606	10,606		16		0		16	29,705
6,321	6,321		1	0	0		1	29,210
8,151	8,151		0		0	0	0	31,070
8,551	8,551				1		1	21,795
9,444	9,444				(0)		0	(22,634)
(65,678)	(65,678)	13	61	0	54	0	128	(570,059)
								+
								8
(65678)	(65678)	13	61	0	54	0	128	(570067)
(050/ð)	(050/8)	13	01	U	54	U	128	(570067)

Table 15-7. Retained catches (metric tons, whole weight) of ISC Members of shortfin make sharks (*Isurus oxyrhinchus*) by fishery in the North Pacific Ocean, north of the equator. "0"; Fishing effort was reported but no catch. "+"; Bellow 499kg catch. "-"; Unreported catch or catch information not available. *: Data from the most recent years are provisional.

Catch		KO	R	M	EX		TWN	
disposition	Year	Longline	KOR Total	Others	MEX Total	Longline	Purse seine	
Retain catch	1985		10441	43	43		Seme	10441
	1986			84	84			
	1987			197	197			
	1988			248	248			
	1989			135	135			
	1990			288	288			
	1991			228	228			
	1992			376	376			
	1993			442	442			
	1994			336	336			
	1995			333	333			
	1996			413	413			
	1997			401	401			
	1998			386	386			
	1999			439	439			
	2000			539	539			
	2001			491	491			
	2002			488	488			
	2003			471	471			
	2004			865	865			
	2005			609	609			
	2006			641	641			
	2007			689	689			
	2008	-	-	609	609			
	2009	-	-	653	653	78		78
	2010	-	-	760	760	54		54
	2011	-	-	758	758	208		208
	2012	0	0	715	715	74		74
	2013	8	8	711	711	107		107
	2014	8	8	0	0	119		119
	2015			-	-	322		322
	2016	0	0	-	-	327		327
Retain cat	tch total	16	16	13,348	13,348	1,289		1,289
Release	2011							
	2012						0	0
	2016	1	1					
Relea	se total	1	1				0	0
Fotal		17	17	13,348	13,348	1,289	0	1,289
Numbers in par								
Sharks catch is	all retain	ed, and no o	discard dat	a.				

				A	US			
Total	USA Total	Sport	Purse seine	Others		Troll	Harpoon	Drift gill- net
192	149			19			1	129
394	310			59			1	250
596	399			188			3	208
571	323			214			3	106
390	255			137			1	117
661	373			141			3	229
445	217			91			1	125
516	140			19			3	118
562	120			32			1	87
463	127			46			1	80
427	94			14			1	79
508	95			9			1	85
533	132			11			3	118
484	98			12			1	85
500	61			9			+	52
615	76			12			+	64
532	41			10			1	30
569	81			12			+	69
537	66			9			+	57
917	52			13			1	38
643	34			8			1	25
686	45			7			+	38
732	43			6			+	37
642	33			5			1	27
760	29			7		0	1	21
834	20			10			0	10
982	16			8			0	8
809	20			11		0	0	9
854	28			12			0	16
152	25	9		9			0	7
337	15	2		4	1			7
345	18	0		4	1	0	1	12
14,824	171	11		1,148	2	0	30	2,343
0	0		0					
0								
1								
1	0		0					
14,825	171	11	0	1,148	2	0	30	2,343